**12455-30** -3240

Printed at 100% size on this 11" x 8.5" sheet







MOLES & POCKET GOPHERS



## USE RESTRICTIONS (Continued from other panel)

- This product may only be used to control pocket gophers\*\* and moles\*
- in manual, below-ground applications in lawns.

  Bait must be applied directly into the burrow or tunnel systems.

- Bat must be applied directly into the burrow of utunel systems.
   Don drapply in garders and areas where food or feed may become contaminated.
   Any person who retrieves carcasses or unused bit following application of this product must wear waterproof gloves.
   Do not apply this product by the use of an artificial burrow builder.
   Not to be sold in individual containers containing more than 1 lb. of product.

APPLICATION DIRECTIONS FOR POCKET GOPHERS\*\*: Apply bait

Plug Removal Baiting: Burrowing pocket gophers (Thomomys spp. and Fig hemova batting, continuing pocuse gopties (Triticianys size and Geomys spp) frow out fow, lan shaped mounds on either side of their underground turnel. These lateral funnels coming to the surface are on the flat side of the fan and are plugged with soil. Using a long-handled spoon, carefully remove the plug, Insert one teaspoon of bat as far down as possible. Reclose opening, being careful not to cover balt with soil.

Main Runway Baiting: To locate main runways, probe about 6 to 8 inches away from mounds by use of an iron bar, a strong stick, or other suitable implement. Resistance on probe will drop suddenly and probe will fall about two inches when main runway is entered. Carefully enlarge the opening to accommodate insertion of the bait. Insert one teaspoon of bait into each active tunnel. Carefully close the hole with a stone, sod, or soil. Take care not to cover bait with soil.

APPLICATION DIRECTIONS FOR MOLES\*: The presence of moles

APPLICATION DIRECTIONS FOR MOLES\*: The presence of moles way be indicated by a network of surface ridges in the Lift, by a series of conical mounds of earth pushed up from deep burrows. Surface Ridges: Pirio to treatment, determine which burrows are active according to one of the following two membrds. Either press down as mail section of the tunnel or remove a one-inch section of the tunnel's root. Mark these sections and recheck in 24 hours. The burrow is considered active if the fillathered runway has been raised up or if the roof has been repaired. Only treat active burrows.



\*Not for mole use in California or Indiana. Not for use against star-nosed mole in North Carolina. \*No usar para topos en California o Indiana No usar para star-nosed moles en North Carolina \*\*Not for use against gophers in North Carolina.
\*\*No usar para gophers en North Carolina

Carefully punch a hole in the top of active tunnels, drop a teaspoonful amount of TOMCAT MOLE\* & GOPHER\*\* BAIT and carefully close the hole with sod or stone. Do not collapse the runway again or allow loose soil to cover the bait. After a few days

MOLES & POCKET GOPHERS

runway again or allow lose so lit o lover the ball. After a lew days, check to see if the burrows are still active according to one of the two methods described above. If so, repeat treatment.

Conical Mounds: Remove center plug from the conical mounds with a long-handled spoon and drop at leaspoorful amount of TOMCAT MOLE\* & GOPHER\*\* BAIT in the underground tunnel. Close opening with rock or sod. Do not allow losse dirt to cover bait.

FOLLOW-UP: Wearing waterproof gloves, collect and dispose of dead, exposed animals. Dispose of lettover balt in accordance with disposal instructors. Use deteigent and not water to wash spoors used for application into burrows. Do not use spoon for mixing, holding or transferring food or feed. If reinfestation does occur, repeat treatment.

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HAZARUS IO HUMANS AND UDMESTIC ANIMALS
Harmful if swallowed or inhaled, causes moderate eye irritation, avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long sleeved shirt and long pants, shoes plus socks and gloves. Remove and wash contaminated clothing before reuse. Any person who retrieves carcasses or unused balt following application of this product must wear gloves.

## ENVIRONMENTAL HAZARDS

This product is extremely took to birds, fish and other wildlife. Wildlife feeding on treated bait may be killed. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply

PHYSICAL-CHEMICAL HAZARDS Do not use with or store near any oxidizing agents

## STORAGE AND DISPOSAL

STORAGE: Store only in original container in a dry place inaccessible to children and pets.

Infacessing to crudined and pels.

DISPOSAL: Nonrefillable container. Do not reuse or refill this container. More fillable container. If empty: Place in trash or offer for recycling if available. If partially filled: Place in trash or call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DISCLAIMER: To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with product other than indicated on the label. To the extent consistent with applicable law, the buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

#### KEEP OUT OF REACH OF CHILDREN CAUTION

See back and side panels for First Aid and additional precautionary statements. EPA REG. NO. 12455-30-3240 EPA EST. NO. 12455-WI-1

ACTIVE INGREDIENT: Zinc Phosphide

**NET WT 6 oz** (168g)

## FIRST AID

MOLES & POCKET GOPHERS

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-854-2494 for emergency medical treatment information. If you experience signs and symptoms such as nausea, abdominal pain, lightness in ohest, or weakness, see a physician immediately. For information on pesticide products (including health concerns, mediciael emergencies, or pesticide incidents), call the National Pesticide Information Conter at 1-000-85-7278. Pesticide Information Center at 1-800-858-7378.

If swallowed . Call a poison control center or doctor immediately for treatment advice or transport the patient to the neares Do not drink water Do not administer anything by mouth or induce vomiting unless told to do so by the poison control center or doctors.

 Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If in eyes

If person is not breathing, call 911 or ambulance, ther give artificial respiration, preferably mouth-to-mouth, i

Call a poison control center or doctor for further nt advice

# TREATMENT FOR PET POISONING

## NOTE TO PHYSICIAN OR VETERINARIAN

Contains the phosphilite-photouring active, and in tribugation in mucosal damage may contraindicate the use of gastric lavage. For animals ingesting bait and/or showing poisoning signs, induce vomitting by using hydrogen peroxide. Sodium bicarbonate can be given orally to neutralize the stomach acidity. The stomach and intestinal tract can be evacuated, oxygen administered and cardiac and circulatory stimulants given.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions and use precautions

## USE RESTRICTIONS:

This product may only be applied against the pests listed on this label, at the use sites listed on this label, and by the application methods listed on this label.

