



Bug-No-More

LAWN & GARDEN INSECT CONTROL

► **FOR USE ON FLOWERS, FRUITS AND VEGETABLES, TREES AND SHRUBS**



► **PREVENTS NEW INFESTATIONS** ► **KILLS OVER 200 LISTED INSECTS BY CONTACT**

Treats up to 8,000 sq. ft. of gardens

ACTIVE INGREDIENT:	By Wt.
Zeta-Cypermethrin*	0.35%
OTHER INGREDIENTS:	99.65%
	TOTAL 100.00%

*See attached booklet for complete
Precautionary Statements and Directions
for Use (including First Aid and
Storage and Disposal).*

THIS PRODUCT CONTAINS:
0.03 pounds of active ingredient per gallon.
*Cis/trans ratio: Max. 75% (+/-) cis and
min. 25% (+/-) trans.

**Shake Well
Before Using**

**KEEP OUT OF
REACH OF CHILDREN**

For Outdoor Residential and Farmstead
Uses Including Lawns, Ornamentals,
Perimeter and Home Gardens

**NET CONTENTS:
40 FL. OZ.**

724/G-2015



Bug-No-More

LAWN & GARDEN INSECT CONTROL

→ Getting Started

→ **Where To Use:** For flowers, fruits, vegetables, fruit and nut trees, and trees

Fruits, Nuts, and Vegetables Mixing Chart			
	Pump-Style Sprayers	For Backpack and 12-Volt Spray Equipment	
Amount of Water in Sprayer	1 gallon	4 gallons	5 gallons
Amount of Product to Add to Sprayer	5 fl. oz.	20 fl. oz.	25 fl. oz.
Area to Treat in Square Feet	1,000	4,000	5,000

For ornamentals, including flowers, roses, trees and shrubs: Apply 15 fl. oz. per gallon of water per 1,000 sq. ft.



For questions or comments call toll-free **800.884.3179**

Mon.-Fri. 8:00 a.m. - 4:30 p.m. CST



or visit: **GordonsUSA.com**



See inside pages for complete Precautionary Statements and Directions for Use (including First Aid and Storage and Disposal).

724/6-2015 AP081414
EPA REG. NO. 279-3472-2217
Lot numbers beginning with 02:
EPA EST. NO. 2217-KS-2
Lot numbers beginning with B:
EPA EST. NO. 57144-GA-1



(40 FL. OZ.)

Plants	Pests	Use Before Harvest
FRUITS and NUTS (cool.)		
Pome-Fruit Trees Apple, Crabapple; Loquat; Mayhaw; Pear; Oriental Pear; Quince	Apple Maggot, Codling Moth, European Apple Sawfly, Green Fruitworm, Japanese Beetle, Lesser Appleworm, Ribwort, Leafroller, Oriental Fruit Moth, Panicle Leafroller, Pear Psylla, Plum Curculio, Pear Leafhopper, Red-banded Leafroller, Rusty Apple Worm, Spina Aphid, Spotted Tentiform Leafminer, Stink Bugs, Tarnished Plant Bug, Tulle's Apple Bud Moth, Variegated Leafroller, White Apple Leaf-opper	Spray up to 14 days before harvesting.
Grapes	Eastern grape leafhopper, Variegated leafhopper, Western grape leafhopper	Spray up to 14 days before harvesting.
Nut Trees Almond; peach nut; Shellnut; hickory nut; chestnut; chestnut; hazelnut; Walnut (hazelnut); Hickory nut; macadamia nut; pecan; walnut (black and English)	Black Pecan Aphid, Codling Moth, Flower Weevil, Pecanery Shuckworm, Leafroller II, etc., Navel Orangeworm, Oak-leaf-miner Leafroller, Peach Twig Miner, Pecan Leaf Chalcid, Pecan Nut Casebearer, Pecan Phylloxera, Pecan Weevil, Pear Bud Bug, Saw Bug, Walnut Aphid, Walnut Husk Fly, Yellow Pecan Aphid	Spray up to 7 days before harvesting. Do not spray more than 1-2 times per season.

ORNAMENTALS, INCLUDING FLOWERS, ROSES, SHRUBS and TREES

For treatment of listed pests in annual and perennial flowers, roses, bedding plants, ground covers, shrubs and deciduous and evergreen trees:

- Spray 15 fl.oz. per gallon of water per 1000 sq.ft. up to the point of run-off making sure to cover all sides of leaves and tree trunks.
- For ground covers, bed plants, foliage, shrubs and general use, apply 15 fl.oz. per gallon of water as a uniform spray. 32 fl.oz. will treat 2-3 sq.ft. Repeat treatment as necessary to achieve control using higher application on roses as pest pressure & foliage area increases.

Plant Safety Note: Certain plants may be sensitive to the residual spray solution. A initial area should be treated and observed for one week prior to application to the entire planting.

LAWNS

For treatment of Ants, Fire Ants, Beetles, Chalcid Bugs, and Weevils, Weevils and Other Listed Pests in Lawns:

- To treat bare pest lawns, uniformly apply 15 fl.oz. per gallon of water per 1000 sq.ft. over lawn. Contents will treat up to 2666 sq.ft.
- Lawn should be no longer than 2 inches at time of application.
- Repeat application only if there are signs of renewed insect activity.

Ant Mound Treatment

- To treat ant mounds, including mounds of fire ant mounds, dilute 0.8 fl.oz. in 1-2 gallons of water and spray to fully soak each mound and the immediate surrounding area up to 4 ft. in diameter around the mound. Use the higher volume for mounds greater than 12 inches in diameter. 32 fl.oz. will treat up to 40 mounds.
- For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.

OUTDOOR TREATMENT

To kill ants, cockroaches, and other listed pests on outside surfaces and around perimeter of buildings (except for spiders, scorpions, crickets, mole crickets, grasshoppers and blacklegged and brown dog ticks) - use specific instructions below:

- NOTE: The application of this product to vinyl and aluminum siding (particularly lightly colored, aged, weathered or otherwise damaged) may result in staining, bleaching or discoloration. Factors such as extreme heat and direct sunlight can promote damage. Before applying this product to vinyl or aluminum siding, treat a small area and evaluate 24 hours later to allow any discoloration to occur. Regardless of the test results, do not apply to vinyl or aluminum siding which appears to discolor or during the heat of the day.
- Apply as a perimeter treatment to provide a barrier to listed insects. For best results, also apply to the outside surfaces of buildings (instructions) up to a maximum height of 3 feet above grade, and to surrounding bare ground, vegetation (such as bushes and trees), and turf-covered areas.
 - To prevent pests from entering buildings, apply 1 fl.oz. per gallon of water to a band of soil and vegetation 5 to 10 feet wide around and adjacent to the building. Also, treat the building foundation up to a maximum height of 3 feet above grade where pests are active and adjacent to the building. Thoroughly and uniformly wet the band area, using 1 gallon of spray solution per 1,000 sq.ft.
 - The surface to be treated should be dry at the time of application.
 - When applying to vegetation, such as bushes and trees, spray to the point of run-off.
 - To kill ants, apply directly to mounds, around doors, windows and other places where ants may be found.

For Outdoor Treatment of Carpenter Bees

- To kill carpenter ants, apply as a perimeter treatment as described above.
- Locate and treat carpenter ant nests for best results. Also apply to trails around doors and windows and other places where carpenter ants have been observed or are expected to travel.
- DO NOT directly treat finished wood with this product. To protect finished wood from carpenter ants (and termites), spray the soil beneath where the finished wood is situated. 32 fl.oz. will treat an area of up to 640 sq.ft. Not intended to replace structural treatments. Not recommended as a sole preventive against termites. Applications of Spot-Treat Lawn & Garden Insect Control are not a substitute for mechanical alterations, soil or foundation treatment but merely as a supplement. For active infestations, seek professional inspection and treatment.

For Outdoor Treatment of Bees and Wasps

- To kill bees and wasps, thoroughly spray 1 fl.oz. per gallon of water into nest entrance and any surrounding areas where these pests are found.
- Apply late in the evening when insects are at rest.

For Outdoor Treatment of Blacklegged and Brown Dog Ticks

- To kill blacklegged ticks and brown dog ticks and to prevent infestation of buildings, apply 15.0 fl.oz. per gallon of water per 1000 sq.ft. to soil and vegetation around the exterior perimeter of the building as described in the instructions for use in perimeter treatment.
- Spray to the point of runoff, applying spray thoroughly and uniformly on treated areas.
- Also treat areas of taller vegetation (7-9 ft.) around the exterior of the property and soil and vegetation which border wooded areas of the property where blacklegged and brown dog ticks may be harboring.

For Outdoor Treatment of Spiders, Scorpions, Mole Crickets, Crickets and Grasshoppers

- To kill spiders, scorpions, crickets, mole crickets, and grasshoppers and to prevent reinfestation of buildings, apply 15 fl.oz. per gallon of water per 1,000 sq.ft. to soil and vegetation around the exterior perimeter of the building as described in the instructions for use in perimeter treatment.
- Spray to the point of runoff, applying spray thoroughly and uniformly on treated areas.

PESTS KILLED

Caterpillars, Butterflies and Moths	Leaf-minning caterpillars, common fruiting caterpillars, chrysalis-forming caterpillars, leaf feeding caterpillars including larvae of brown moth, tentacles and other common caterpillars, vegetables and trees), leaf eating caterpillars, common fruiting caterpillars, vegetable feeding caterpillars, stem and trunk boring caterpillars, and leaf-minning caterpillars, Colorado of the Yucca Moth family, Girdler Moth family, Bagworm moth family, Girdler moth family, Eastern Moth family, Oak Wooded moth family, Tortricid Moth family (moths) such as Codling moth, Sawfly family, Whites, Sphagnum and Orange Tree family, Gummy-winged B. Herbivore family, Metalmark Moth family, Small Tortricid family, Dark-footed Butterfly family, Milkweed and Gummy Butterfly family, Satyr, Rymidae and Arctia family, Milkweed Butterfly family, Smoky Moth family, Pyralid Moth family such as Saw Worms Moth, Phoenix Moth family, Mues, tobacco Moth family, Tiger Moth family, Ctenuchid Moth family, Tussock moth family, Owllet Moth family such as Eight-spined Forest
Common Burdettia	Silver-sh Family
Springtails	Polined Springtail family such as Snow-Pea, Sashwort Springtail, Erioceroid Springtail family such as Andrew's Springtail, Golden Springtail etc.
Cockroaches	Nest of Cockroach Family such as Oriental Cockroach (Habit Cockroach, Shed Cockroach) and American Cockroach (Waterbug), Blackhead Cockroaches family such as German Cockroach (Droten Bug)
Termites	Rotting-wood termite family such as Pacific Coast Termite, Subterranean termite family such as subterranean termites. Not intended to replace structural treatments. For termites - not recommended as a sole preventive against termites. Such applications are not a substitute for mechanical alterations, soil or foundation treatment but merely as a supplement. For active infestations, get a professional inspection.
Earwigs	Long-Horned Earwig family such as Ring-legged Earwig, and Hymen Earwig; Common Earwig family such as European Earwig
Grasshoppers & Crickets	Pygmy Grasshopper family, Long-horned grasshoppers and Killyd-d family, Canal Crickets family, Tree Crickets, fly such as House Cricket
True Bugs	Plant Bug family, Assassin bug family, Lace Bug family, Seed Bug family, Salt B. g. family, Leaf-roller Bug family, Scumless Plant Bug family such as Eastern Bordered Bug and Western Bordered Bug, Sawk Bug family
Cicadas and Kix	Cicada family such as Periodical Cicadas, Dogday Hares, and Grand Western Cicadas
Treehoppers	Treehopper family, such as Threemass Treehopper, Oak Treehopper, Bull's Treehopper, and Local Treehopper
Leafhoppers	Leafhopper family such as Edge Leafhopper, Scarlet-and-green Leafhopper, Spangle-leaver
Thrips	Winged Thrips family such as Broad-winged Thrips, Common Thrips family
Beetles	Tiger Beetle family, Ground Beetle family such as Blunt-bell Beetle - Pesticidous Diving Beetle family, White-spotted Beetle family, Water Beetle family, Blue Beetle family, Common Beetle family, Stag Beetle, Beetle, grass family, Scarab Beetle family, Click Beetle family, Soldier Beetle family, Not-winged Beetle family, Golden Beetle family, Branch and Tong Beetle family, Characene Beetle family, Bark Beetle family, Ladybug, Beetle, Diving Beetle, Two-colored Beetle family, Water Beetle family, Long-horned Beetle family, Sand Beetle family, Leaf Beetle family, Primitive Wren's family, Ground Beetles and Weevil family, Shaw and Ambrosia Beetle family
Bees, California Carpenter Bees, Ants, Wasps, Sawflies	Dumbbell Sawflies family, Common Sawfly family, Hairtail family, Stem Sawflies family, Braconids family, Ichneumonids family, Tomyid Wasps family, Chalcid family, Gall Wasps family, Pelecinid family, Tenthredinid Wasps family, Vespid, Ant, Stink Wasps family, Ant family, Vespe Wasps family, Soldier Wasps, Scheuch Wasps family, Bee families
Flies	Crane fly family, Phorid Crane Flies, Net-winged Midges, Black Flies, March Fly family, Fungus Gnats family, Scorpion Fly family, Silblot Fly family, Flower-loving Fly family, Mydas Fly family, Robber Fly family, Bee Fly, Dance Flies, Long-legged Flies, Hopper Fly family, Thick-headed Fly family, Pygmy Fly family, Fruit Fly family, Sawwood Fly family, Marsh Fly family, Shore Fly family, Phoenix Fly family, Anthomyid Fly family, Muscid Fly family, Loose Fly family, Blow Fly family, Flesh Fly family, Leached Fly family
Spiders	Folding webster spider family, Tarantula family, Functore Spider family, Spitting spider family, Wolf Spider family, Comb-footed Spider family, Sheet-web Weaver family, Old Weaver family, Large-bodied Orb Weaver family, Funnel Web Weaver family, Nursery Web Spider family, Wolf Spider family, Lynx Spider family, Sac Spider family, Wandering spider family, Giant Crab spider family, Suberoid Crab Spider family, Crab Spider family, Phidippid family, Daddy-long-legs family
Ticks	Blacklegged tick, Stream-dog tick, Deer tick
Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects	Psyllid family, Whitefly family, Aphid family, Woolly and Gall Making Aphid family, Giant Scale Insects family, Armored Scale insects family, Mealybug family, Cottonmeel Bug family
Millipedes	Millipedes

IMPORTANT - Directions for Storage and Disposal

Pesticide Storage	Protect from freezing. Store this product in its original container in a cool, dry (preferably locked) area that is inaccessible to children and animals. After partial use, replace original lid and store tightly.
Container Disposal	Never flush contents. Do not reuse or refill this container. If empty: Place in trash or other for recycling, if available. If partially filled: Call your local soil & waste agency for disposal instructions. Never place unused product down any incoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or otherwise.

2/14/15-2015 AP01414
EPA REG. NO. 579-3474-2/11
Lot numbers beginning with 02 EPA EST. NO. 2217-KS-2
Lot numbers beginning with B EPA EST. NO. 57144-GA-1

MANUFACTURED BY
PAGCORPORATION
1217 WEST 18TH STREET
KANSAS CITY, MISSOURI 64101
www.gardens.com

**GORDON'S® BUG-NO-MORE®
LAWN & GARDEN INSECT CONTROL**

SEE HERE TO OPEN

KEEP OUT OF REACH OF CHILDREN

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-877-800-5556, twenty-four (24) hours per day, seven (7) days per week.	
NOTE TO PHYSICIAN: Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive.	
For Information Regarding the Use of this Product Call 1-877-800-5556.	

724/6-2015 AP081414
EPA REG. NO. 279-3472-2217
EPA EST. NO. 2217-KS-2
Lot numbers beginning with 02:
EPA EST. NO. 2217-KS-2
Lot numbers beginning with B:
EPA EST. NO. 57144-GA-1

**MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com**