

SPREAD IT & FORGET IT

SPREAD IT & FORGET IT Plus Dimension 0.15% Herbicide

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

This bag covers up to 6,280 sq ft

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, this product may be applied only by commercial application at no more than 232 lb per 3 in active ingredient per acre (7.3 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Active Ingredient: 37.00% Urea Nitrogen, 5.00% Soluble Potash (K2O), 58.00% Inert Ingredients

CAUTION

Keep Out of Reach of Children

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation - Harmful if Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Rinse contaminated clothing and wash before reuse. Preclude all frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

First Aid

If in eyes: Hold eye open and flush slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue flushing. Call a poison control center or doctor for treatment advice.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance. Give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Environmental Hazards: This product is toxic to fish and highly toxic to other aquatic organisms including crayfish and shrimp. Use with care when applying to turf areas adjacent to any body of water. Oil and drift from treated turf may adversely affect aquatic organisms at adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to rice field areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washings.

Other Use Precautions: Do not use cloppings from treated turf for mulching around vegetables or fruit trees. Do not allow other people or pets to enter treated areas until dust has settled. Do not cut grass immediately or feed large old farm animals treated with this product.

Directions for Use

Storage and Disposal: Do not contaminate water, foodstuffs, feed or feed by storage or disposal. Pesticide Storage: Store this product only in its original container in a dry, cool, unventilated storage area. Pesticide Disposal: Waste resulting from the use of this product that cannot be used or otherwise repackaged should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local jurisdiction. Contained Handling: Retainable container. Do not reuse or refill this container. Other: If recycling is available, it should be recycled. If not, dispose of empty bag in a sanitary landfill, or by extension, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.
Use Directions for Turf: This product provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs, and tee boxes.

Use Precautions for Turf

This product may be used on existing sodded or sprigged lawns and ornamental lawns that are well-established, well irrigated and have received at least 10 irrigations following seeding, sodding or sprigging. Use of this product on lawngresses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may reduce the chances of turf injury.

Apply this product only to lawngresses specified on the label. The following lawngresses species are tolerant to this product when applied as directed:

Table with 4 columns: Grass Species Group, Turfgrass Species, Warm Season Grasses, and Cool Season Grasses. Rows include: Bermuda grasses (Bermuda, Common, Hybrid, Dimple), Fescue (Hard, Fine-leaved), Ryegrass (Perennial, Annual), and Zoysia grasses.

Like the product on certain varieties of creeping bentgrass, such as 'Columbus', 'Canton', 'Seaside', and 'Wheatfield'. May result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colored bentgrass (Agrostis ssp.).

Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Alondra', 'Banner', 'Beauvois', 'Blazer', 'CS-7', 'Eony', 'HF-22', 'Mighty', 'Natic', 'Sunsetworth', 'Tortol', 'Majestic', 'May', 'Paradise', 'Tanager', 'Tanger', 'Worship', and 'Widdow' spp. All varieties of fine fescue have been tested.

Do not use this product on seedling tallgrasses in the spring of the first year of establishment until the tallgrass is fully green and has established new roots. Note: It is not possible to evaluate all lawngresses varieties for tolerance to this product. It is suggested that the user evaluate the response of lawngress varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

Precautions for Re-seeding, Overseeding, or Sprigging

- Re-seeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 15 weeks after a split application program totaling 7.7 to 1000 sq ft 10.5 lb active ingredient per acre) or more may inhibit the establishment of desirable tallgrasses.
- When re-seeding or overseeding, follow established cultural practices for seedbed preparation, seedling emergence and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (see dealers) designed to provide good contact between soil and seed.

Precautions for Control of Crabgrass

Early post-emergence weed control with this product is limited to crabgrass. Post-emergence application effectively controls crabgrass, but only if application is made prior to the 3rd stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core, spike, or hydro-mulching, and weeding, should be done before application of this product. Do not apply this product until crabgrass has received from the mechanical injury caused by time cultural practices.

Other Use Precautions

- Do not use cloppings from treated turf for mulching around vegetables or fruit trees.
- Do not allow other people or pets to enter treated areas until dust has settled.
- Do not cut grass immediately or feed large old farm animals treated with this product.

Product label for 37-0-5 fertilizer. Includes guaranteed analysis (37.00% Urea Nitrogen, 5.00% Soluble Potash), derived from polymer coated urea, and application instructions for Florida lawns and golf courses.

