



# Fairway Formula **GREENVIEW**<sup>®</sup> Spring Fertilizer with Crabgrass Preventer

**Prevents crabgrass for up to 120 days**

**Extended time release feeding of nitrogen greens  
into summer without excess growth**

<b>ACTIVE INGREDIENT:</b>	
Dithiopyr.....	0.13%
<b>OTHER INGREDIENTS:</b> ..... 99.87%	
<b>TOTAL</b> ..... 100.00%	

CAS. No. 97886-45-8  
EPA Reg. No. 961-392

(L) (D)  
EPA Est. No. 961-PA-01, 961-IL-01  
Superscript used is first letter of lot number.

**Covers 5,000 sq. ft.**

# 26-0-5

**Net Weight 17 lbs. (7.7 kg.)**

## **KEEP OUT OF REACH OF CHILDREN CAUTION**

See back panel for First Aid and additional Precautionary Statements

**READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.**

**USE ONLY ACCORDING TO LABEL INSTRUCTIONS.**

If terms are not acceptable, return the unopened container at once.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year. Do not apply more than once per year to the same area.

**ACTIVE INGREDIENT:**

Dithiopyr..... 0.13%

**OTHER INGREDIENTS:**..... 99.87%

**TOTAL** ..... 100.00%

CAS. No. 97886-45-8

EPA Reg. No. 961-392

(L) (D)

EPA Est. No. 961-PA-01, 961-IL-01

Superscript used is first letter of lot number.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid and additional Precautionary Statements

**READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.**

**USE ONLY ACCORDING TO LABEL INSTRUCTIONS.**

If terms are not acceptable, return the unopened container at once.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year. Do not apply more than once per year to the same area.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

First Aid	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of an emergency involving this product, call day or night (888) 208-1368.	

#### Environmental Hazards:

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

## PRODUCT INFORMATION FOR TURF USE

### Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This product contains Dimension<sup>®</sup>, a herbicide that can be applied up to 4 weeks later than other crabgrass preventers and still provides season-long control. Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer prevents crabgrass and other grass and broadleaf weeds up to 4 months, when applied prior to their germination in lawns and ornamental turfgrasses. It also provides postemergence control of seedling crabgrass for up to 4 weeks after it germinates.

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer should be applied to lawns and ornamental turfgrasses that are well established (See "Precautions and Restrictions" section below for a list of the tolerant turfgrasses). It will not harm nearby established ornamentals when used according to directions, given below.

### Precautions and Restrictions

**NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.**

**NOT FOR USE ON NEWLY SEEDED LAWNS.**

Use of Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Do not apply to lawns or turfgrasses that are under stress.

Reseeding, overseeding, or sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until after 8 weeks from the time of application. Reseeding, overseeding, or sprigging before 8 weeks or earlier after application may injure or prevent the establishment of the newly reseeded, overseeded or sprigged turfgrasses.

Do not make an application of this product later than 4 weeks after crabgrass germination. Do not expect control with applications made later than 4 weeks after crabgrass germination.

**DO NOT** apply more than 8.8 lbs. of this product per 1,000 sq. ft. per application, and no more than 26.4 lbs. of this product per 1000 sq. ft. per year.

In the State of New York, do not exceed 4.4 lbs. product per 1000 sq. ft. (0.25 lb. active ingredient per acre) per year.

**DO NOT** use clippings from treated turf for mulching, vegetables, vines, or fruit trees from which feed or food products may be harvested.

Keep people and pets off treated areas until dust has settled.

### Weeds Controlled and/or Suppressed

**Grass Weeds** - Annual Bluegrass, Annual Ryegrass, Barnyardgrass, Crabgrass, Crowfootgrass, Goosegrass, Green Foxtail, Kikuyugrass (seedling), Perennial Ryegrass, Smutgrass (seedling), Yellow Foxtail.

**Broadleaf Weeds** - Bittercress, Buttercup Oxalis, Carolina Geranium, Chickweed, Common Lespedeza, Common Purslane, Corn Speedwell, Creeping Oxalis, Henbit, Parsley-piert, Pineappleweed, Prostrate Spurge, Spotted Spurge, Yellow Woodsorrel.

# APPLICATION DIRECTIONS

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer can be applied to the following turfgrasses when used as directed:

**Cool-Season Turfgrasses** - Creeping Bentgrass, Fine Fescues, Kentucky Bluegrass, Perennial Ryegrass, Tall Fescues.

**Warm-Season Turfgrasses** - Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Centipedegrass, St. Augustinegrass, Zoysiagrass.

## Application Timing

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer should be applied in the spring or in the fall prior to weed seed germination for control of weeds in established lawns and ornamental turfgrasses. When this product is applied in the fall, a sequential or second application must be made the following spring to provide season-long control of weeds. If this product is applied in the fall, DO NOT reseed, overseed or sprig turfgrasses before 12 weeks after application due to potential turfgrass injury to the newly reseeded, overseeded or sprigged turfgrasses.

When establishing a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass, wait until the grass is well established (2 mowings) before applying this product. Do not apply this product within an 8 week period prior to seeding, reseeding, overseeding or sprigging.

Improved weed control may occur when this product is applied within a few days after mowing and further mowing is delayed until the turfgrass has been watered or received rainfall to dissolve the fertilizer granules.

Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

## Length of Control

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer will provide up to 4 months of weed control if applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, and intensity of weed pressure.

## Post Emergence Crabgrass Control

For post emergent crabgrass control, apply Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer when crabgrass is small (3-4 leaf). It will not control large tillered crabgrass.

## Product Use Rate Table (Lbs. Per 1,000 Sq. Ft.)

Region	2-3 Months Control	3-4 Months Control	Early Post Emergence Control
North	2.2	3.1	3.9 - 8.8
South	3.4	3.9 - 4.5	3.9 - 8.8

Use the higher rate in areas of high weed pressure

## Use Regions within the United States



## Application Equipment and Instructions

Greenview Fairway Formula Spring Fertilizer with Crabgrass Preventer can be applied with a drop or rotary type spreader designed to apply granular fertilizers. For best results, apply this product evenly and uniformly over the turfgrass area, avoiding streaking, skips or overlaps. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader settings so the spreader delivers the appropriate application rate as specified above.

A more uniform application can be made by spreading one-half of the specified amount of this product over the turfgrass area and then applying the remaining one-half at a right angle to the previous direction. Check spreader frequently to insure it is functioning properly and distributing the granules uniformly.

Preventer when crabgrass is small (3-4 leaf). It will not control large tiered crabgrass.

frequently to insure it is functioning properly and distributing the granules uniformly.

## SPREADER SETTINGS

### Drop Spreader Settings

Apply in a single pass at a rate of 3.4 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.



#### Drop Spreader

#### Setting (3.4 lbs. / 1000 Sq. Ft.)

Greenview Drop Spred XP1	15
Greenview Drop Spred III	6
Agri-Fab Drop	6 1/2
Red Devil Drop (1-20)	9
Republic Drop	6 1/2
Scotts Drop PF	6

To calibrate your spreader: 1 lb. of material should cover approximately 294 sq. ft.

If your spreader model is not listed, go to our website [www.greenviewfertilizer.com](http://www.greenviewfertilizer.com), or call toll-free 800-233-1067,

8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

### Broadcast Spreader Settings

Apply in a single pass at a rate of 3.4 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.



#### Broadcast Spreader

#### Setting (3.4 lbs. / 1000 Sq. Ft.)

Greenview Jet-Spred SS1	14
Greenview Jet-Spred III	3 1/2
Agri-Fab® Rotary	3 1/2
Earthway Rotary	14
Lesco Rotary (letter/numeric)	H/15
Red Devil Rotary (1-20)	6
Republic Rotary	3
Sears Craftsman Rotary	3 1/2
Scotts Speedy Green	4
Scotts Edgeguard	4

