

**Shaklee®**

Concentrated Germicide

**Basic-G®**

Disinfects • Cleans • Deodorizes

**ACTIVE INGREDIENTS**

DIDECYL DIMETHYLAMMONIUM CHLORIDE .....	10.14%
n-ALKYL (C <sub>14</sub> 50%, C <sub>12</sub> 40%, C <sub>16</sub> 10%) DIMETHYL BENZYLAMMONIUM CHLORIDE .....	6.76%
INERT INGREDIENTS .....	83.10%
TOTAL .....	100.00%

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

SEE BACK PANEL FOR  
PRECAUTIONARY STATEMENTS  
AND FIRST AID

**1 Qt. (946 mL)**

NA167A

## Basic-G®

**DILUTION:** 1:256 (660 ppm quat)  
1 ounce per 2 gallons of water, 1/2 ounce per gallon of water.

For use in households. Disinfects, cleans, and deodorizes the following hard, nonporous, inanimate surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.). Effective against the following pathogens: Salmonella enterica, Bordetella bronchiseptica, Chlamydia psittaci, Legionella pneumophila, Listeria monocytogenes, Salmonella enteritidis, Salmonella typhi, Salmonella typhimurium, Staphylococcus aureus—Methicillin Resistant (MRSA), Staphylococcus aureus—Vancomycin Intermediate Resistant (VISA), Streptococcus pyogenes (Strep), Trichophyton mentagrophytes. See reference sheet for a complete list of organisms claimed for this product.

**DIRECTIONS FOR USE:** It is a violation of federal law to use this product in a manner inconsistent with its labeling. This product is not for use on medical-device surfaces.

**DISINFECTION/CLEANING/DEODORIZING DIRECTIONS:** Remove heavy soil deposits from surface. Then thoroughly wet surface with a use-solution of 1/2 oz. of concentrate per gallon of water or equivalent. (Use 2 oz. per gallon of water to kill Adenovirus Type 7). The use-solution can be applied with a cloth, mop, sponge, or coarse spray, or by soaking. For sprayer applications, use a coarse spray device. Spray 6–8 inches from the surface, rub with a brush, cloth, or sponge. Do not breathe spray. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Food contact surfaces must be thoroughly rinsed with potable water. This product must not be used to clean the following food contact surfaces: utensils, glassware, and dishes. Prepare a fresh solution daily or more often if the solution becomes visibly dirty or diluted.

**FUNGICIDAL DIRECTIONS:** For use in shower and bath areas, follow disinfection directions.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS—DANGER. CORROSIVE.** Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), protective clothing, and protective gloves (rubber or chemical resistant). Harmful if inhaled. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**FIRST AID: IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or when going for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

**STORAGE AND DISPOSAL:** Store in original container in areas inaccessible to children. Nonrefillable container. Do not reuse or refill this container. Wrap container and put it in trash or offer for recycling if available.

 **Shaklee®**  
Creating Healthier Lives®

#00525

DISTRIBUTED BY:  
SHAKLEE CORPORATION  
4747 WILLOW ROAD  
PLEASANTON, CA 94588

EPA Registration No. 47371-129-7294  
EPA Est. No. 10508-OH-001

★ NC981B



## Shaklee® Concentrated Germicide Basic-G® - Reference Sheet

This product is a multipurpose neutral pH germicide and deodorant for use in households. Disinfects, cleans, and deodorizes the following hard, nonporous, inanimate surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.).

**Dilution:** 1:256 (660 ppm quaternary strength) - 1 ounce per 2 gallons of water; ½ ounce per gallon of water. Use 2 oz. per gallon of water to kill Adenovirus Type 7.

**EPA Registration Number:** 47371-129-7294

### Directions For Use:

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read this product label affixed to the container before using and applying.
- Use of this product according to the label is subject to the usage precautions and limitations imposed by the product label affixed to the container.

### *Basic-G® Germicide is effective against the following pathogens:*

#### Bacteria

Pseudomonas aeruginosa<sup>1</sup>  
Staphylococcus aureus<sup>1</sup>  
Salmonella enterica  
Acinetobacter calcoaceticus  
Bordetella bronchiseptica  
Chlamydia psittaci  
Community Associated Methicillin-Resistant Staphylococcus aureus (NRS384) (USA300)  
Community Associated Methicillin-Resistant Staphylococcus aureus (NRS123) (USA400)  
Enterobacter aerogenes  
Enterobacter cloacae  
Enterococcus faecalis - Vancomycin Resistant (VRE)  
Escherichia coli<sup>1</sup>  
Fusobacterium necrophorum  
Klebsiella pneumoniae<sup>1</sup>  
Legionella pneumophila  
Listeria monocytogenes  
Pasteurella multocida  
Proteus mirabilis  
Proteus vulgaris  
Salmonella enteritidis  
Salmonella typhi  
Salmonella typhimurium  
Serratia marcescens  
Shigella flexneri  
Shigella sonnei

Staphylococcus aureus - Methicillin resistant (MRSA)  
Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA)  
Staphylococcus epidermidis<sup>2</sup>  
Streptococcus faecalis<sup>1</sup>  
Streptococcus pyogenes (Strep)

#### Viruses

\*Adenovirus Type 4  
\*Adenovirus Type 7 (mixed 2 oz. Basic-G/gallon of water)  
\*Hepatitis B Virus [HBV]  
\*Hepatitis C Virus [HCV]  
\*Herpes Simplex Type 1<sup>1</sup>  
\*Herpes Simplex Type 2<sup>2</sup>  
\*Human coronavirus  
\*HIV-1 [AIDS virus]  
\*Influenza A / Hong Kong Flu Virus\*  
\*Pandemic 2009 H1N1 Influenza A virus [formerly swine flu]  
\*Respiratory Syncytial Virus [RSV]  
\*Rotavirus [WA]  
\*Rubella [German Measles]  
\*SARS Associated Coronavirus  
\*Vaccinia [Pox virus]

## Animal Viruses

---

- \*Avian Influenza (H5N1)
- \*Avian polyomavirus
- \*Canine distemper
- \*Equine Herpes Virus Type 1
- \*Feline coronavirus
- \*Feline leukemia
- \*Feline panleukopenia virus
- \*Feline picornavirus
- \*Infectious bovine rhinotracheitis
- \*Infectious bronchitis (Avian IBV)
- \*Newcastle Disease
- \*Porcine parvovirus
- \*Pseudorabies (PRV)

- \*Rabies
- \*Transmissible gastroenteritis virus (TGE)

## Fungi

---

- Aspergillus niger
- Candida albicans
- Trichophyton mentagrophytes (Athlete's Foot Fungus)

- 
- 1 ATCC & antibiotic-resistant strain
  - 2 Antibiotic-resistant only
  - \* Virucidal

For effective use of this product, always follow label instructions, paying special attention to the dilution rate, precautions, and contact time. Please note that this product is not to be used on the skin or to be taken orally.

Revised February 2012