



ferti·lome®

Weed-Out™

Lawn Weed Killer

Contains TRIMEC® Herbicide

- Kills troublesome Broadleaf Weeds: Dandelion, Chickweed, Clover, Henbit & Many Others As Listed

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See back panel for complete Precautionary
Statements, including First Aid, and Directions for
Use, including Storage and Disposal.

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	5.45%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	5.88%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	1.21%

OTHER INGREDIENTS:	87.46%
TOTAL:	100.00%

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.

Isomer Specific by AOAC Methods.


TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

NET CONTENTS: ONE GALLON (128 FL. OZS. / 3.785 LITERS)

FIRST AID

If In Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15–20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If On Skin Or On Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15–20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

In the event of a medical or chemical emergency, contact  **North America 1-800-255-3924.**

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

Manufactured By:



230 FM 87
 BONHAM, TX 75418
 EPA Reg. No. 2217-597-7401
 EPA Est. No. 7401-TX-01
 Visit Us At: www.fertilome.com
 Product Questions? 855-270-4776



7 32221 10519 5

KEEP FROM FREEZING.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets. Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not exceed specified dosages for any area.
- Do not spray Carpetgrass, Dichondra, or lawn where desirable clovers are present. Do not apply this product to “Floratam” St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- Do not spray when this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Failure to follow above precautions may result in injury.

PRODUCT INFORMATION

ferti•lome® Weed-Out™ Lawn Weed Killer controls troublesome common lawn weeds, including Poison Ivy, in both Northern and Southern Grasses.

WHEN TO USE

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

BROADCAST TREATMENTS

Limitations on broadcast applications to residential turfgrass: The maximum application rate is 4 fl. oz. of product per 1,000 sq. ft. per application (0.72 lb 2,4-D ae, 0.57 lb MCPP-p ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 fl. oz. of product per 1,000 sq. ft. (1.14 lb 2,4-D ae, 1.06 lb MCPP-p ae and 0.24 lb dicamba ae per acre).

BROADCAST TREATMENTS WITH HOSE-END SPRAYERS WITH DILUTION:

Spray preparation for hose-end dilution sprayers:

Add the required amount of **ferti•lome® Weed-Out™ Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Table 1. RATES OF ferti•lome® Weed-Out™ Lawn Weed Killer FOR VARIOUS LAWN GRASSES				
Gallons Of Water Spray	Area to Treat (\$q. Ft.)	St. Augustinegrass, Centipede ¹ grass, Bahiagrass (fl. oz.)	Bermudagrass, Buffalograss, Bentgrass (fl. oz)	Bluegrass Zoysiagrass, Fescue (fl. oz)
Hose End Sprayer with dilution				
1	250	½	2/3	2/3 to 1
4	1,000	2	2 2/3	2 2/3 to 4
8	2,000	4	5 1/3	5 1/3 to 8
20	5,000	10	13 1/3	13 1/3 to 20
32	8,000	16	21 1/3	21 1/3 to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup
8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
32 fluid ounces = 1 quart (qt.)

SPOT TREATMENT WITH PRESSURE SPRAYERS

Limitations on spot treatments to residential turfgrass: Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4 fl. oz. per 1,000 sq. ft. per application (0.57 lb MCPP-p acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND BAHIAGRASS: Mix 2/3 fl. oz. (4 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. Do not apply to St. Augustinegrass, Centipede¹grass, or Bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on “Floratam” St. Augustinegrass in Florida.

BERMUDAGRASS, BUFFALOGRASS, AND BENTGRASS: Mix 1 fl. oz. (2 Tbs) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

KENTUCKY BLUEGRASS, FESCUE, PERENNIAL RYEGRASS, AND ZOYSIAGRASS: Mix 1-1/3 fl. oz. (8 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

BROADLEAF WEEDS					
Annual Yellow Sweetclover Aster Austrian Fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, Field Bird Vetch Bitter Wintercress Bittercress, Hairy Bitterweed Black-Eyed Susan Black Medic Black Mustard Blackseed Plantain Blessed Thistle Bloodflower Milkweed Blue Lettuce Blue Vervain Bracted Plantain Bristly Oxtongue Broadleaf Dock Broadleaf Plantain Broomweed Buckhorn Buckhorn Plantain Bulbous Buttercup Bull Thistle Bullnettle Burclover Burdock Burning Nettle Burweed Buttercup	Buttonweed Canada Thistle Carolina Geranium Carpetweed Catchweed Bedstraw Catnip Catsear Chickweed, Common Chickweed, Mouseear Chicory Cinquefoil Clover, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Cockle Cocklebur Common Mullein Creeping Jenny Cudweed Curly Dock Daisy, English Daisy Fleabane Daisy, Oxeye Dandelion Dichondra Dogfennel Dollarweed False Dandelion Falseflax False Sunflower	Fiddleneck Florida Pusley Frenchweed Galinsoga Goldenrod Ground Ivy Gumweed Hairy Fleabane Hawkweed Healall Heartleaf Drymary Heathaster Hedge Bindweed Hedge Mustard Henbit Hoary Cress Hoary Plantain Hoary Vervain Horsenettle Jimsonweed Knawel Knotweed Kochia Lambsquarters Lespedeza Mallo Matchweed Mexicanweed Milk Vetch Morningglory Mouseear Hawkweed Mugwort Musk Thistle Mustard Narrowleaf Plantain	Narrowleaf Vetch Nettle Orange Hawkweed Oriental Cocklebur Oxalis Parsley-Piert Parsnip Pearlwort Pennycress Pennywort Pepperweed Pigweed Pinewoods Bedstraw Plains Coreopsis Plantain Poison Ivy Poison Oak Pokeweed Poorjoe Prairie Sunflower Prickly Lettuce Prickly Sida Prostrate Knotweed Prostrate Pigweed Prostrate Spurge Prostrate Vervain Puncturevine Purslane, Common Ragweed Red Sorrel Redroot Pigweed Redstem Filaree Rough Cinquefoil Rough Fleabane Russian Pigweed Russian Thistle	Scarlet Pimpernel Scotch Thistle Sheep Sorrel Sheeperspurse Slender Plantain Smallflower Galinsoga Smooth Dock Smooth Pigweed Sorrel Sowthistle Spanishneedles Speedwell Spiny Amaranth Spiny Cocklebur Spiny Sowthistle Spotted Catsear Spotted Spurge Spurweed Stinging Nettle Strawberry, India Mock Tall Nettle Tall Vervain Tansy Ragwort Tansy Mustard Tanweed Thistle Trailing Crownvetch Tumble Mustard Tumble Pigweed Velvetleaf Venice Mallo Virginia Buttonweed	Virginia Creeper Virginia Pepperweed Wavyleaf Bullthistle Western Clematis Western Salsify White Mustard Wild Aster Wild Buckwheat Wild Carrot Wild Four-O’Clock Wild Garlic Wild Geranium Wild Lettuce Wild Marigold Wild Mustard Wild Onion Wild Parsnip Wild Radish Wild Rape Wild Strawberry Wild Sweet Potato Wild Vetch Woodsorrel Woolly Croton Woolly Morning-glory Woolly Plantain Wormseed Yarrow Yellow Rocket Yellowflower Pepperweed

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Manufactured By:



230 FM 87 • BONHAM, TX 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-01
Product Questions? 855-270-4776

10519-1-0620-CL

