



FIFRA Section 24(c) Special Local Need (SLN) Label

For Distribution and Use Only in the State of Oregon

JMS Stylet-Oil

EPA Reg. No. 65564-1 EPA SLN No. OR-120011

For Control of Rust (*Puccinia* spp.) on Winter and Spring Wheat

This label is valid until December 31, 2029 or until otherwise amended, withdrawn, canceled, or suspended.

Active ingredient:

Mineral oil* 97.1%

Inert ingredients:..... 2.9%

Total:.....100.0%

*Contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Environmental Hazards: Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of JMS Stylet-Oil.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This SLN label and the label affixed to the main container of JMS Stylet-Oil must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the main container of JMS Stylet-Oil.

Disease	Application Rate	Application Timing	Restrictions
Rust (<i>Puccinia</i> spp.)	Ground: 1-2 pints JMS Stylet-Oil/acre in minimum 15 gals water/acre Aerial: 1-2 pints JMS Stylet-Oil/acre in 5 to 10 gals water/acre	Feekes Growth Stage: 6 through 10.5.	Maximum 3 applications per year. No preharvest interval.

USE INSTRUCTIONS:

For Ground Application: Apply JMS Stylet-Oil in a minimum of 15 gallons of water per acre by ground equipment. Use a spray volume that assures uniform coverage of the infested portions of the treated crop. Calibrate equipment to the desired spray volume. Optimum results are received when higher spray volumes are used.

Aerial Application: For best results, apply JMS Stylet-Oil in 5-10 gallons of water per acre. Higher gallonage will provide better coverage and performance.

Thorough coverage and proper equipment calibration are critical to achieve good control when aerially applying JMS Stylet-Oil. Aerial applications may not provide the same level of efficacy that can be obtained with ground equipment.

Chemigation: Do not apply this product through any type of irrigation system.

Tank Mixing with Other Fungicides:

Combining JMS Stylet-Oil with other fungicides labeled for use on wheat may increase the efficacy of both products. In limited trials, phytotoxicity has not been evident; however, it is always recommended to test products for compatibility and possible phytotoxicity. Follow all applicable directions, restrictions, and precautions in the Chemical and Physical Compatibility and Phytotoxicity section of the label affixed to the main container of JMS Stylet-Oil.

Section 24(c) Registrant:

JMS Flower farms, Inc.
4423 5th PL SW
Vero Beach, FL 32968
772-562-8770