

Clinafarm[®] EC

Emulsifiable Concentrate

For control of *Aspergillus fumigatus* contamination in poultry hatchery equipment

ACTIVE INGREDIENT:

Imazalil (enilconazole): 1-(2-(2, 4-Dichlorophenyl)-2-(2-propenyloxy)ethyl)-1H-imidazole..... 13.8% w/w

INERT INGREDIENTS..... 86.2%
100.0% w/w

KEEP OUT OF REACH OF CHILDREN
DANGER

See Back Panel for Additional Precautionary Statements

131696 R2



TM

FIRST AID

If In Eyes •Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for further treatment advice. **If Swallowed** •Call a poison control center or doctor immediately for treatment advice. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person. •Promptly drink either large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. **If On Skin Or Clothing** •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice. **If Inhaled** •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. •Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling the poison control center or doctor or going for treatment. You may also contact the Rocky Mountain Poison Center at 1-303-595-4869 for emergency medical treatment information.



Intervet Inc, a subsidiary of Merck & Company, Inc.,
Summit, NJ 07901. All rights reserved.

Net Contents: 750 mL (1.58 US Pints)

Product of Janssen Research
EPA Reg. No. 773-55, EPA Est. No. 618-LA-1
Rev. 9/12

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER**

Corrosive, causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if absorbed through skin or inhaled. Avoid breathing spray mist and contact with skin. Do not get in eyes, wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

133827 R2

TAKE TIME



**OBSERVE LABEL
DIRECTIONS**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For the control of *Aspergillus* organisms in cleaned poultry and turkey hatchery equipment prior to introduction of the eggs. Equipment would include empty hatcher cabinets, setters, coolers, storerooms, handling equipment, etc. CLINAFARM® EC will dramatically reduce the levels of infectious organisms and spores in treated areas. To make 1 gallon of 1:100 dilution, add 1.25 fl. oz. (36mL) of CLINAFARM® EC to one gallon of clean water to provide an antifungal concentration of 0.15% active ingredient. Do not use product undiluted. To assure proper efficacy, the area or equipment to be treated should be cleaned and closed off from outside air currents before treatment. Ventilation ducts may be treated but fans must be shut off. Fog or spray area to be treated at the rate of ½ fl. oz. diluted product per 150 cu. ft. (1 mL per 10 cu. ft.). Fog or spray to ensure good contact with surface areas. Do not re-enter treatment areas for two hours. CLINAFARM® EC is compatible with other commonly used disinfectants such as quaternary ammonium compounds, glutaraldehyde, formaldehyde, and phenolic compounds.

PHYSICAL/CHEMICAL HAZARDS: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store at room temperature.

PESTICIDE DISPOSAL: Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CLINAFARM is a registered trademark of Johnson & Johnson CF-15

Rev. 9/12

batching legends
to be printed online