



Soap-Shield® Flowable Liquid Copper Fungicide

**Controls powdery mildew,
black spot & rust**



Intended For Residential Use Only

ACTIVE INGREDIENT:

Copper Octanoate (Copper Soap)..... 10.0%

CAS Reg. No. 20543-04-8

OTHER INGREDIENTS 90.0%

TOTAL 100.0%

metallic copper equivalent 1.8%

EPA Reg. No. 67702-2-56872, EPA Est. No. 50591-GA-01

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panels for first aid
and directions for use.

NET CONTENTS: 16 Fl. Oz.

64432

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals.

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the poison control center at 1-800-222-1222.	

Environmental Hazards: This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

STORAGE AND DISPOSAL

Pesticide Storage: Store in a secure place, away from open fire or flame. Keep container closed and reseal after use. Product may be damaged by freezing. Do not store product below 4° C. If spilled, use absorbent materials and dispose of in an approved manner.

Pesticide Disposal and Container Handling:

Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Sold under a license of W. Neudorff GmbH KG, Germany

U.S. Patent No.: 5,246,716

Distributed by Gardens Alive!®, Inc.
706 W. Cumberland St.
Lewisburg, OH 45338
Phone: 513-354-1483
www.GardensAlive.com



Soap-Shield®
Flowable Liquid
Copper Fungicide
Controls powdery mildew,
black spot & rust



FOR ORGANIC
GARDENING

- May be used up to the day of harvest.
- For hydroponic gardening.
- Where to use: Listed vegetables, fruits, nuts, ornamentals and turf.
- Use for early and late blight on tomatoes and potatoes.
- Where to use: Listed Vegetables: Controls powdery mildew, downy mildew, Botrytis, Alternaria leaf blight and Septoria leaf spot.
- Controls leaf blight, leafspot, dollar spot and rust on lawns.
- Use to control a wide range of listed plant diseases: powdery mildew, rusts, blackspot, leaf & fruit spot, downy mildew, fruit rot and late blight.

FIRST AID

IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the poison control center at 1-800-222-1222.

KEEP OUT OF REACH
OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals.

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards: This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Peel back tab for Precautionary Statements, Directions for Use and Storage and Disposal instructions.



Sold under a license of W.
Neudorff GmbH KG, Germany

U.S. Patent No.: 5,246,716

BATCH CODE:

Distributed by Gardens Alive!®, Inc.
706 W. Cumberland St. Lewisburg, OH 45338
Phone: 513-354-1483 www.GardensAlive.com



1

92665

04956

9

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions and precautions on this label before using.

Do not allow adults, children or pets to enter the treated area until sprays have dried. Do not apply this product in a way that will contact adults, children or pets, either directly or through drift.

Application Directions

Shake well before use. Most conventional liquid pesticide plant sprayers can be used to apply Soap-Shield® Flowable Liquid Copper Fungicide to plants. Mix 0.5 to 2.0 fluid ounces of Soap-Shield® Flowable Liquid Copper Fungicide in a gallon of water. Unless otherwise stated in specific application notes, apply 1.1 to 2.3 gallons of diluted spray per 1,000 ft².

For best control, start spraying before the disease is visible or when disease is first visible on the plant. Spray all plant parts thoroughly (top and bottom of leaves), and unless otherwise directed in the crop table, repeat every 7 to 10 days. Use the higher rate to control diseases that may go dormant and overwinter. See application notes for the specific crop application interval. If possible, time applications so that at least 12 hours of dry weather follows application. Reapply after rain, following crop specific application interval.

Ornamentals

Do not use more than 373 fl. oz. of product/1000 sq. ft. per year on ornamentals. Soap-Shield® Flowable Liquid Copper Fungicide may cause some copper toxicity on some plant species. Before spraying a specific plant species consult your State Experiment Station or make a test spray.

Hydroponic use: Apply as soon as disease appears, or as a preventive spray 2 weeks before disease normally appears. Apply as a foliar spray only. Do not apply directly to the water in hydroponic growing systems. Discarded water from hydroponic growing systems may be used in greenhouses and to water indoor and outdoor container plants and garden areas. Water from these systems is prohibited from being discarded directly into a water source.

When **powdery mildew** presence is expected on a plant, spray the plants at the minimum application interval during the first 2 weeks after emergence.

To control **downy mildews, leaf and fruit spots, blights, and rust**, begin treatment 2 weeks before disease normally appears or when weather forecasts predict a long period of wet weather. Alternatively, begin treatment when disease first appears, and repeat as long as needed.

To prevent **fruit rots**, apply at the start of flowering and unless otherwise directed in the crop table, repeat every 7 to 10 days until harvest. Fungicidal sprays are especially warranted when weather forecasts predict a long period of wet weather.

EPA Registration No. 67702-2-56872

EPA Establishment No. 50591-GA-01

Crop	Diseases Controlled	Application Notes
Pine	Needle blight	Spray until needles are thoroughly wet with spray. Apply when new needles are just emerging.
Rose and Ornamental Shrubs (such as Crape Myrtle, Forsythia, Hydrangea, Willow, Mock-Orange, Deutzia, Pyracantha, Japanese quince, Abelia Summersweet)	Blackspot, Downy mildew, Gray mold (Botrytis), Leaf spots, Powdery mildew, Rust	May cause copper toxicity on some rose varieties. Copper toxicity appears as purple spots. For Black spot, mix 1.44 fl. oz. of Soap-Shield® Flowable Liquid Copper Fungicide per gallon of water. For Powdery Mildew, mix 1.08 fl. oz. of Soap-Shield® Flowable Liquid Copper Fungicide per gallon of water. In damp cool conditions (below 60°F), phytotoxicity is likely to occur.

Fruit and Vegetables

Crop	Diseases Controlled	Maximum Annual Rate (fl. oz. product/ 1000 sq. ft.)	Application Notes
Bean, Pea	Anthrachnose leaf and fruit spot, Ascochyta leaf and pod spot, Bacterial blights (halo, common and brown spot), Downy mildew, Gray mold (Botrytis), Powdery mildew, White mold (Sclerotinia)	For beans: 88 fl. oz. For peas: 73 fl. oz.	Repeat application every 7-10 days if needed.
Beet, Sugar Beet, Chard, Spinach	Cercospora leaf spot, Downy mildew, White rust, Powdery Mildew, Anthracnose blue mold	For beets: 146 fl. oz. For spinach and chard: 73 fl. oz.	Repeat application every 7-10 days if needed.
Carrots	Alternaria leaf blight, Bacterial leaf blight, Cercospora leaf blight	93 fl. oz.	Repeat application every 7-10 days if needed.
Citrus (Grapefruit, Lemon, Lime, Kumquat, Orange, Pummelo, Tangerine)	Melanose spot, greasy spot, citrus scab, Alternaria brown spot, citrus canker, Phytophthora brown rot, and Septoria	235 fl. oz.	Repeat application every 14 days if needed. May cause phytotoxicity if conditions are conducive, when mixed with other products, or when applied to citrus seedlings grown in greenhouses or shadehouses.
Corn	Alternaria blight, Anthracnose, Ascochyta leaf and pod spot, Bacterial blights (halo, common, and brown spot), Bacterial leaf spot, Downy mildew, Gray mold, Southern leaf blight, Cercospora leaf blight, Common or Southern Rust, Gray Leaf Spot, Stewart's Wilt*, Bacterial Stalk Rot*	78 fl. oz.	Repeat application every 7-10 days if needed.
Crucifer Crops (Bok Choy, Broccoli, Brussels sprouts, Canola, Cauliflower, Cabbage, Kale, Kohlrabi, Mustard, Pak-Choi, Rape, Rutabaga, Turnip)	Alternaria blight, Bacterial leaf spot, Downy mildew, Powdery mildew, White mold (Sclerotinia), Black rot	49 fl. oz.	Repeat application every 7-10 days if needed.

Crop	Diseases Controlled	Maximum Annual Rate (fl. oz. product/1000 sq. ft.)	Application Notes
Cucurbits (Cucumbers, Honeydew, Muskmelon, Watermelon, Cantaloupe, Melons, Squash, Pumpkin, Zucchini)	Alternaria blight, Angular leaf spot, Anthracnose leaf and fruit spot, Downy mildew, Gray mold, Scab, Ulocladium leaf spot, Powdery mildew, Watermelon bacterial fruit blotch, Fruit blotch (suppression)	97 fl. oz.	For cucumbers grown in a greenhouse, apply every 5 days in the first 2 weeks after emergence, then reapply every 7 days.
Currant and Gooseberry	Anthracnose leaf and fruit spot, Phyllosticta, Septoria leaf spots, Powdery mildew	298 fl. oz.	Repeat application every 7-10 days if needed.
Herbs (basil, chives, coriander, dill, lavender, mint, parsley, rosemary)	Anthracnose, Alternaria blight, Bacterial Blight, Botrytis, Downy mildew, Leaf scorch, Leaf spot, Rhizoctonia Leaf blight	Dill: 73 fl. oz. Parsley: 37 fl. oz. Other listed herbs: 49 fl. oz.	Repeat application every 10-14 days if needed.
Grapes	Downy mildew, Black rot, Phomopsis Cane and Leaf Spot, Powdery mildew, Gray mold (Botrytis), ripe rot*	373 fl. oz.	Repeat application every 7-10 days if needed. Do not mix product with lime. Certain Vinifera and French Hybrid varieties may be sensitive to copper sprays resulting in marginal leaf burn. Before spraying these varieties, consult your State Experiment Station or make test sprays.
Hop	Anthracnose leaf and fruit spot, Cercospora leaf spot, Downy mildew, Powdery mildew	49 fl. oz.	Repeat application every 10 days if needed.
Lettuce, Chicory, Endive	Downy mildew, Septoria leaf spot, Powdery mildew, Bacterial soft rot and bottom rot	149 fl. oz.	Repeat application every 10 days if needed. Use lower rate when disease pressure is low or on copper sensitive varieties of lettuce.

Crop	Diseases Controlled	Maximum Annual Rate (fl. oz. product/ 1000 sq. ft.)	Application Notes
Onion, Garlic, Leek, Shallot	Botrytis leaf blight, Downy mildew, Neck rot, Bacterial soft rot, Purple blotch	112 fl. oz.	Repeat application every 7-10 days if needed.
Peanuts	Sclerotinia blight, Leaf spots (early and late), web blotch	88 fl. oz.	Repeat application every 7-10 days if needed.
Pome Fruits (Apples, Pears, Quince)	Anthrachnose, Cedar Apple Rust, Fireblight, Scab, Sooty Blotch, Flyspeck, Quince Rust, Blossom Blast, European Canker, Collar Rot, Crown Rot	298 fl. oz.	For fireblight, apply in the dormant period, during bloom, or as in-season cover spray. Do not exceed one application during the fall, late dormant period. Do not exceed one application between silver tip and green tip growth stages. May cause russetting of susceptible apple varieties. Do not exceed 1 gallon of product per 100 gallons of water. Repeat application every 7-10 days during bloom and growing season if needed.
Blackberry, Blueberry, Raspberry	Gray mold (Botrytis), Mucor fruit rot, Rhizopus fruit rot, Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust, Bacterial Canker	Blueberry: 156 fl. oz. Blackberry and Raspberry: 186 fl. oz.	Apply at the start of flowering and continue every 7-10 days until harvest.
Stone Fruit Trees (Almond, Apricot, Cherry, Nectarine, Peach, Plum)	Bacterial canker (Pseudomonas syringae), <i>Monolinia</i> brown rot and blossom blight, leaf and fruit spot, Bacterial leaf spot, Bacterial Blast (Pseudomonas), black knot* (plum only)	335 fl. oz.	For bacterial canker, apply as a dormant spray as buds begin to swell, repeating at the bud burst stage, and weekly thereafter as needed, up to six sprays. In the fall spray again at 10 and 80% leaf fall. For brown rot blossom blight, apply full cover spray at delayed dormant (bud swell), popcorn, full bloom and petal fall stages. Repeat application every 7-10 days if needed.
	Anthrachnose leaf and fruit spot, Coryneum blight, Peach leaf curl	335 fl. oz.	Apply as a dormant spray in late fall to before bud break. Repeat application every 7-10 days if needed.

Crop	Diseases Controlled	Maximum Annual Rate (fl. oz. product/ 1000 sq. ft.)	Application Notes
Strawberry	Angular leaf spot, Leaf scorch, Mycosphaerella leaf spot, Phomopsis leaf blight, Powdery mildew, Septoria leaf spots	152 fl. oz.	Spray 1 month after planting (or before flowering on established plants). Repeat application every 7-10 days if needed.
	Anthrachnose fruit rot, Gray mold (Botrytis)	152 fl. oz.	Apply at the start of flowering and repeat application every 7-10 days until harvest.
Tobacco	Blue Mold (Downy Mildew)	149 fl. oz.	Repeat application every 10 days if needed.
Tomato, Potato, Eggplant, Pepper	Anthrachnose leaf and fruit spot, Bacterial speck, Bacterial spot, Cercospora leaf spot, Early blight, Gray mold, Late blight, Leaf mold, Septoria leaf spot	Tomato: 149 fl. oz. Potato: 466 fl. oz. Eggplant: 147 fl. oz. Pepper: 221 fl. oz.	Repeat application every 7-10 days if needed. Use the 2 fl. oz. per gallon of water dilution rate for late blight.
Walnut	Blight	596 fl. oz.	Repeat application every 7-10 days if needed.
Avocado	Anthrachnose, blotch, Scab	353 fl. oz.	Apply when blossom buds open. Repeat application every 14-30 days if needed.

*Not registered for use in California.

NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law, buyer assumes all risk of any such use. Seller makes no other warranties, either expressed or implied.



Sold under a license of W.
Neudorff GmbH KG, Germany

U.S. Patent No.: 5,246,716

Distributed by Gardens Alive!®, Inc.
706 W. Cumberland St. Lewisburg, OH 45338
Phone: 513-354-1483 www.GardensAlive.com