

Professional The Constant Professional Andersons Turf™ Products

Weed & Grass Preventer with 5% Treflan* Herbicide (Dispersible Granules)

Net Wt. 40 Lbs. (18.14 kg)

Not For Turf Use

A selective, preemergence herbicide for the control of certain annual grasses and broadleaf weeds in Lendscape Ornamentals, Ground Covers, Establehed Flowers, Ornamental Bulbs, Non-bearing Fruit and Nut Trees and Non-bearing Vineyards

ACTIVE INGREDIENT: triffurakn: alpha, alpha, alpha-trifluoro-2,6-dinitro-N,N-dipropyl-

EPA Reg. No. 9198-129 EPA Est. No. 9198-OH-1<u>M</u>, 9198-OH-2<u>B</u>, 9198-AL-001<u>A</u> Underlined letter is first letter used in run code on bag.

KEEP OUT OF REACH OF CHILDREN CAUTION **Precautionary Statements**

- FRET AID

 If never Hold siye open and rinse slawly and gently with water for 15-20 minutes. Remove contacts, if present, after the linst 5 minutes, then continue minute. If present, after the linst 5 minutes, then continue minute.

 If walllowed: Call a poten control center or doctor for headment article. Have person sip a glass of water in able to swallow. Do not induce vomiting unless told to by a poten content or doctor. Do not give amything by multit to an unconscious person. It also different and doctors. I have of contaminated doctors. related contaminated doctors. I have also minutes. I have a first minutes.

Call a poison control center or doctor for theatment advice.

Move person to fresh ar.

If initialed:

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if

Possible.

- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for irealment.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

AUTION: Casses moderate eye intation, Audit contact with eyes or docting rolonged or frequently repeated skin contact may cause Allergic reaction in ome individuals.

Personal Protective Equipment (PPE)
Applicators and other handlers must wear
Long-sleeved shirt and long pants
Socks and shoes

User Safety Requirements: Follow manufacturer's instructions for cleaning/ mainfaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gurn, using lobacco, or using the loilel. Users should remove clothing immediately it pesidiot gets inside. Then wash thoroughly and put on clean dollving. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean dollning.

ENVIRONMENTAL HAZARDS
This pestioide is extremely loate to freshwater marine, and estuarine fish and equatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to interfided areas below the mean high water mark. Do not apply in a menner which will directly expose canals, lakes, steams, poorls, marrises or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Do not contaminate water when disposing of equipment washwaters.

II is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying. Not for use on plants being grown for sale or commercial use, or for commercial seed production, or for research purposes. For use on plants inlended for assisting purposes or limitation modification and being grown in ornamental gardens or parks, or on got courses or lawns.

STORAGE AND DISPOSAL.

Do not contaminate water, food, or feed by giorage or disposal. Peaticle Storage: Effect in display contamination city, it cases or lawns in contamination of the contamination city, it cases or lawns in contamination of the contamination city, it cases to all as or at an approved waste disposal backty. Container Disposal: Non-Hillable container. Once it causes or reliff the container. Other for excitation of the container. Other container is the container of the container. Other for excitation of the container. Other for excitations of the container.

or by inchestion or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product is a preemergence herbicide for control of certain annual grasses and broaded reviewed in containing rown ornamentals, landscape commentals, namery slock, ground covers, established flowers, contained bottos, non-bearing trial and ruit frees and non-bearing visiguants. Apply this product before or after plaining but prior to germination of target weeds, or immediately after cutivation. Length of weed control will visit yeally weed to produce the product before and produce of control will visit yeally will weed opposition, portion period or set of conditions, temperature, watering regime, and other factors. Fellowing application, set should recoller and observed with weed control over time to determine when additional applications, may be needed. Repeat applications should not be made abone than 60 days allor a previous application of his product. Do not apply over 240

pounds per acre total of this product within a 12 month period.

This product does of with pocular winning at a transmission. This product does not control established weeds. Existing weeds should be cointrolled by collivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into sol prior to inclaiment, 308 should be in good condition and free of clodes at the time of application. A single raintail or sprinider irrigation of 0.5 inches or more, or flood irrigation, is required to activate this product, of priorimum weed control is obtained when this product is activated within 3 days of application and tillage is possible, this product may be activated using cultivation equipment capital or uniformly mixing line herbicide into the upper 1-2 inches of sol. Failure to activate this product within 3 days of application may result in errato weed control. Do not apply when wind conditions favor drift of this product's granules from the target area.

Oplimum weed control will be obtained when followed by overhead irrigation or rainfall within a few hours after surface application, Weeds controlled by this product when applied at 80 lbs./acre (1,8 lbs./1000 sq. ft.):

Annual Graéses:
Barfey, Hare
Barnyardgrass (Watergrass)
Bluognass, Annual
Brome
Unglefro
Cheal
Crabgrass
Oat, Wild
Cupgrass, Southwestern
Pancum, Fall Signalgrass Spranglelop, Mexican Stink Grass Witchgrass

Landscape Ornamentals, Ground Covers, Established Flowers, Ornamental Bulbs, Non-bearing Fruit and Nut Trees and Non-bearing

Vineyards:
This product is recommended as a preemergence treatment for control of certain annual grasses and broadlest weeds in landscape ormanentals, ground covers, established flowers, omanental busy, non-bearing fruit and nut treas and non-bearing vineyards. Apply this product before or after plateing but prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information", "General Precaution and Restrictions" and "Application Directions" sections of this label for general use precautions and information on application, application rates and weeds controlled.

GENERAL PRECAUTIONS AND RESTRICTIONS
Entry Restrictions: Do not enter or allow others to enter the treated area
until dusts have settled. If soil incorporation is required after the application,
do not enter or allow others to enter the breated area (except those persons
involved in the incorporation) until the incorporation is complete. If the
incorporation is exemplished by watering-in, do not enter or allow others to
enter the treated area until the surface is dry after the watering-in.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

To avoid possible injury, do not apply this product to:

- cuttings that have been planted in post for the first time
- ground covers until they are established and well rooted

Do not apply this product to newly transplanted ornamentals, ground covers, flowers and non-bearing futil and nut trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury

Do not make preplent applications of this product to areas where gladiolicorns less than one inch in diameter will be planted or injury may occur.

Do not apply this product in greenhouses or other enclosed structures.

Users who wish to use this product on plant species not recommended on this label may determine suitability for such uses by making field application of the product at a recommended rate to small number of plants. Prior to using this product on a larger number of plants, the treated plants should be observed for signs of herbicidal lefury during 30 to 60 days of normal growing condition to determine if the treatment is corriniptious to the larget plant species. The user assumes responsibility for any plant damage or other liability resulting from the use of this product on plant species not recommended on this label.

application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

APPLICATION TECHNIQUES FOR APPLYING THIS PRODUCT When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

A chain lastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

For Irealing smaller areas or rows of nursery stock or ornamental beds, a hand-held or push-type rotary applicator such as whirtybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed,

A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, hard to treat areas. Carefully measure the amount of product needed to avoid over-application.

This product may be used on the following established plant species

TREES:
Balsam Fr
White Fir
Novey Marte
Floar
Nov

Copper Leal Temple Fire
Dwarf Japanese Japanese Boxwood
Maple Korean Boxwood
California Gold

Asparagus Fern Crown Vetch Coreopsis Blue Fescue

ESTABLISHED FLOWERS:

NON-BEARING FRUIT AND NUT TREES AND NON-BEARING IVINEYARDS."

For the control of VINEYARIOS:

föxi Orenge Plum
Lemon Peach Raspbenry
Macadarnia Mut Pear Walhut, Black

*Non-bearing fruit and ruit tiese and non-bearing vineyards are del
plants which will not bear fruit for at least one year after treatment.

IMPORTANT: READ BEFORE USE:

INFORTANT: HEAD BEFORE USE:
Read the entire Directions for Use and the Warranty Disclaimer and
Limitation of Liability before using this product. It farms are not acceptable,
return the unopened product container at once, By using this product, user
or Buyer accepts the following Warranty Disclaimer and Limitation of

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrens that this product conforms to the chemical description on the
jude land is reasonably file for the purposes stated on the jude where used in stoff
accordance with the derections. Manufacturer makes No OTHER EXPRESS OR
IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR
PARTICULAR PURPOSE. To the extent consistent with applicable law, it is
Manufacturer's intent to LIMIT ANY LIABILITY FOR SECILAL, CONSECUENTIAL.
OR INCIDENTIAL ECONOMIC DAMAGES to related of products at Buyer's choice, to the extent constant with applicable law, its
DISCLAMS ANY LIABILITY OF COMPORATORY OR OTHER DAMAGES ANISING
OUT OF ANY USE CONTRANTY OF LABEL DIRECTIONS. Use contrary to label
directions is not permitted.

LEGAL HIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.





Professional Turf™ Products



Professional Weed & Grass Preventer

Turf Products with 5% Treflan* Herbicide (Dispersible Granules)



Visit our website at www.Andersonsplantnutrient.com

©2016 The Andersons, Inc. The Andersons logo is a registered trademark of The Andersons, Inc.,
Professional Turf is a trademark of The Andersons, Inc., Treflan is a registered trademark of Dow AgroSciences LLC

> APTDG5TF4.4 APTDG5TF4.4B