

ENFORCER® Mosquito & Flying Insect Killer II DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Use Restrictions

Do not allow adults, children or pets to enter the treated area until vapors, mists i and aerosols have dispersed and the treated area has been thoroughly ventilated. Do not spray animals. Do not water the treated area to the point of run-off. Do not make applications

during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1) Treatment to soil or vegetation around structures:

Applications to lawns, turf, and other vegetation:

3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches, and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewers or drains, or to any area like a gutter where i drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Not for use in outdoor residential misting systems.

į	This product kills	troublesome outdo	or insects which	n destroy lawns,	roses, flowers, trees ar	nd s
	Ants Aphids	Bagworms Beetles Chinab Bugo	Cicadas Cutworms	Inchworms Leafrollers	Sod Webworms Spider Mites	
	Armyworms	Chinch Bugs	Fleas	Moths	Whiteflies	

Also kills

Brown Dog Ticks Fall Cankerworms Mealybugs Budworms Grasshoppers Crickets Japanese Beetles Mole Crickets Earwigs Leafminers Mosquitoes Elm Spanworms Lone Star Ticks Sowbugs (pillbugs) HOSE-END SPRAYER:

Easy-to-use. For application to lawns, outdoor ornamentals, outside surfaces of buildings and other outdoor residential areas.

32 OZ. TREATS UP TO 5.000 SQUARE FEET.

Snap off tab on top of sprayer before first use.

1. Make sure knob control lever is in the **OFF** position. Shake well and attach to hose.

2. Turn water on at faucet, aim nozzle toward lawn and turn knob

control lever to **ON** to begin spraying.

3. When finished turn control lever to **OFF** position. Turn faucet to OFF position. Discharge residual water pressure by turning ON and OFF again. Store out of reach of children.

LBLPFl32BF • Copy: 7-13 • EPA Reg. No. 28293-271-40849 EPA Est. No. 40849-GA-1(E), or EPA Est. No. 59144-GA-1(A), or EPA Est. No. 44616-MO-1(H), or EPA Est. No. 1270-GA-1(1) Postscript is first letter or number of batch code on container. **Product Code: PFI32**

Sold By: Zep Commercial Sales & Service, a unit of Zep Inc.

1310 Seaboard Ind. Blvd., Atlanta, GA 30318

Questions? Call 1-888-805-4357, Mon - Fri, 9am to 5pm EST.



Tent Caterpillars

Ticks, including

and many others.

Deer Tick (which may

carry Lymè disease)

PEEL OPEN for

precautionary

statements

250

4.

0.500" 3.125" **ADHESIVE PRINT** AREA AREA

BACK LABEL Back of Top Layer

3.625"

NUISANCE PESTS IN OUTDOOR AREAS

OUTSIDE SURFACES OF BUILDINGS

PESTS: Mosquitoes. Ants (including foraging fire ants), Chiquer, Clover Mites, Crickets. Earwigs, Millipedes and Sowbugs (pillbugs).

HOW TO USE: To help prevent infestation of buildings, treat the building foundation to a

height of 3 feet, where pests are active and may find entrance. Also, apply as a residual spray i to outside surfaces of building, including garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness. OTHER OUTDOOR AREAS

PESTS: Ants (including foraging fire ants), Chiggers, Crickets, Earwigs, Fleas, Lone Star Ticks and Ticks, including Deer Tick (which may carry Lyme disease)

HOW TO USE: For treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For Ants, thoroughly wet hills and runways. Repeat application as reinfestation occurs. PESTS: Tent Caterpillars

HOW TO USE: Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have it returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.

HOME LAWNS

BENT, BERMUDA, BLUEGRASS, DICHONDRA, FESCUE, IRISHMOSS, MERION, ST. AUGUSTINE; PESTS: Armyworms, Brown Dog Ticks, Chiggers, Chinch Bugs, Cutworms, Fleas, Japanese; Beetle Grubs, Mole Crickets, Mosquitoes, Sod Webworms, Ticks, including Deer Tick (which i may carry Lyme disease), Ants (including foraging fire ants), Crickets, Grasshoppers **HOW TO USE:** Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed. For heavy infestations, repeat application after 2 weeks.

TREES & SHRUBS

ARIZONA CYPRESS. AZALEA. BIRCH. CHERRY. NON-BEARING CITRUS. CONIFERS. ELM. ENGLISH IVY, EUONYMUS, WEEPING FIG. FIR. HONEYSUCKLE, LILAC, MOCK-ORANGE, OAK. PALM. PHILODENDRON. PINE. POINSETTIA. POPLAR. RHODODENDRON. TAXUS. TULIP-PESTS: Aphids, Bagworms, Cicadas, Exposed Thrips, Fall Cankerworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider Mites, Whiteflies CHERRY, HONEYSUCKLE - PESTS: Fall Webworms

CONIFER - PESTS: Budworms

ELM - PESTS: Elm Leaf Beetles, Elm Spanworms, Fall Webworms

DOUGLAS and GRAND FIRS - PESTS: Tussock Moths.

NON-BEARING CITRUS - PESTS: Citrus Black Flies

OAK - PESTS: Oakworms

PINE - PESTS: Pine Beetles, Pine Moths, Pine Needleminers, Needle Scales

HOW TO USE: Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4 – 8 days. Application can be i made up to the day of collection.

3.625"

BACK LABEL Bottom Layer

ROSES & FLOWERS

AGERATUM, ASTER, AZALEA, BEGONIA, COLEUS, COMMON NINEBARK and SNOWBERRY, EXACUM, GLADIOLUS, GOLD BELLS, HYPOESTES, IVY, LILAC. MARIGOLD. MOCK-ORANGE. NANNYBERRY. ORCHID. PANSY. PEA SHRUB. PETUNIA, POINSETTIA, RHODODENDRON, ROSE, SNAPDRAGON, ZINNIA PESTS: Aphids, Bagworms, Cicadas, Exposed Thrips, Fall Cankerworms, Fall Webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider Mites, Whiteflies

HOW TO USE: Apply as a thorough spray, wetting the plants to the dripping point. Try to in hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray, at the first sign of insects. Repeat as necessary; use intervals of 4 - 8 days. Application can be made up to the day of collection.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area away from heat or open flame. **PESTICIDE DISPOSAL:** If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. CONTAINER DISPOSAL: If empty - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eves or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

0.500" 2.625" 0.500" **ADHESIVE** PRINT **ADHESIVE** AREA AREA AREA

0.500" **ADHESIVE** AREA

250'

2.625" PRINT AREA

0.500" **ADHESIVE** AREA