



# Turf Supreme®

## 16-6-8 with Trimec®

Kills Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds.

Contains TRIMEC® Herbicide to control the toughest weeds

**1 Bag Covers 15,000 sq ft**

<b>ACTIVE INGREDIENTS:</b>	
2,4-dichlorophenoxyacetic acid.....	0.55%
(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid.....	0.12%
Dicamba: 3,6-dichloro-o-anisic acid.....	0.05%
<b>OTHER INGREDIENTS:</b> .....	<b>99.28%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>
THIS PRODUCT CONTAINS:	
11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.	
2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.	
1.0 lb. 3,6-dichloro-o-anisic acid equivalent per ton.	
Isomer Specific by AOAC Methods.	
EPA REG. NO. 2217-643-7001	EPA EST. NO. 93243-TX-1
	EPA EST. NO. 089135-AZ-001

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material.

FIRST AID	
<b>IF ON SKIN OR ON CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center, or doctor, or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

<b>16-6-8 GUARANTEED ANALYSIS</b>	
TOTAL NITROGEN (N).....	16.00%
16.00% Ammoniacal Nitrogen	
AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> ).....	6.00%
SOLUBLE POTASH (K <sub>2</sub> O).....	8.00%
Sulfur (S).....	16.00%
16.00% Combined Sulfur	
Iron (Fe).....	2.00%
0.04% Water Soluble Iron	
Derived from ammonium sulfate, monoammonium phosphate, muriate of potash and iron sulfate.	

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

### INSTRUCTIONS:

Turf Supreme® 16-6-8 with Trimec® controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns as listed. In addition, a 16-6-8 fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

The maximum broadcast application rate is 5.0 lb of product per 1,000 sq ft per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 lb of product per 1,000 sq ft.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Turf Supreme® 16-6-8 with Trimec® effectively controls the following weeds when used as directed.

Broadleaf Weeds Controlled				
Bindweed, field	Burdock	Dichondra	Pearlwort	Scarlet pimpernel
Bird vetch	Burning nettle	Dollarweed	Poison ivy	Scotch thistle
Bitter wintercress	Buttonweed, common	Fiddleneck	Poison oak	Shepherd's purse
Bittercress, hairy	Canada thistle	Goldenrod	Prickly lettuce	Smooth dock
Black medic	Carpetweed	Ground ivy	Prostrate knotweed	Smooth pigweed
Black mustard	Chickweed, common	Henbit	Prostrate pigweed	Sowthistle, annual
Blackseed plantain	Chickweed, mouseear	Hoary cress	Prostrate spurge	Speedwell, common
Blessed thistle	Chicory	Hoary plantain	Puncturevine	Spotted spurge
Bloodflower milkweed	Clover, crimson	Knotweed, prostrate	Purslane, common	Tall Nettle
Bracted plantain	Clover, white	Kochia	Red sorrel	Tumble Pigweed
Bristly oxtongue	Cocklebur	Lambsquarters	Redroot pigweed	Venice mallow
Broadleaf dock	Common mullein	Mallow, common	Redstem filaree	Wayleaf thistle
Broadleaf plantain	Creeping buttercup	Morningglory, tall	Rough cinquefoil	White mustard
Buckhorn plantain	Curly dock	Mustard, tumble	Rough fleabane	Wild carrot
Bulbous buttercup	Daisy, English	Narrowleaf plantain	Russian pigweed	Woolly Plantain
Bull thistle	Dandelion, common	Narrowleaf vetch	Russian thistle	Yarrow, common
Burclover				

Bag Size lb	Calibration Rate lb per sq ft	Application Rate lb per 1,000 sq ft	Treated area for this bag size (sq ft)	
			Cool Season Grass	St. Augustinegrass
50 lb	1 lb per 300 sq ft	3.33 lb	15,000 sq ft	—

### SPREADER SETTINGS

Apply 50 lb (for 15,000 sq ft coverage per bag)

	SGN 150
BEST® PRO.....	4B
Spyker / BEST® Models 34, 44, 64.....	3.7
Earthway.....	12
Lesco.....	13
Scotts R-8A (Cone 8).....	J½
Lely 1250 / Vicon.....	6

As a guide, set spreader at the setting indicated by the above chart before applying. Remember the settings provided are only a guide and depend on the age, condition and speed of operation of your spreader. Check spreader calibration often and make adjustment as needed. Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### LIMITED WARRANTY AND DISCLAIMER

#### FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

© 2024 J.R. Simplot Company. All rights reserved. SIMPLOT®, BEST® and Turf Supreme® are trademarks of the J.R. Simplot Company registered in the United States and other countries. TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**NOTICE:** This product can expose you to chemicals including cadmium which are known to the State of California to cause cancer, birth defects or other reproductive harm. For more information, go to [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapco.org/metals.html>

**Net Weight: 50 lb (22.68 kg)**



Distributed by:  
J.R. SIMPLOT COMPANY  
P.O. Box 70013, Boise, ID 83707  
(208) 336-2110

24305D  
GHS24190  
**ITEM # 9603150**  
Product Labels, SDS and other information on BEST® products are available at [www.BestFertilizer.com](http://www.BestFertilizer.com)