3.75"

plusTrimec

Kills Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds; including those listed on the back panel.

Contains \*TRIMEC® Herbicide to Control the Toughest Weeds 1 Bag Covers 15,000 Sq. Ft.

THIS PRODUCT CONTAINS: **ACTIVE INGREDIENTS:** 2,4-dichlorophenoxyacetic acid ..... 11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton. 2.4 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy) propionc acid equivalent per ton. (+)-(R)-2-(2 methyl-4-chlorophenoxy) propionic acid .............. 0.12% Dicamba: 3,6-dichloro-o-anisic acid...... .... 0.05% 1.0 lb. 3,6-dichloro-o-anisic acid equivalent per ton. Isomer Specific by AOAC Methods. ...100.00% EPA Reg. No. 2217-643-7001 EPA Est. No. 61842-CA-1

# **KEEP OUT OF REACH OF CHILDREN** CAUTION

See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal. \*TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Net Weight: 50 LBS. (22.68 Kg)

# STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY. PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals **CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material.

lf on skin or on clothing:	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If Inhaled:	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
f swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have a person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product co	ontainer or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street
back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.
GUADANTEED ANALYSIS: 16-6-8 SPREADED SETTINGS

<u> </u>	<u> </u>			
GUARANTEED ANALYSIS: 16-6-8	SPREADER SETTINGS			
OTAL NITROGEN (N) 16.0%	Apply 50 lbs. (for 15,000 sq. ft. coverage pe	er b		
40.00/ A	BEST® Pro	4B		
$VAILABLE$ PHOSPHATE ( $P_2O_5$ )	Spyker / BEST® Models 34, 44, 64	3.7		
OLUBLE POTASH (K <sub>2</sub> O)	Earthway	12		
ULFUR (S)	Lesco			
RON (Fe)	Scotts R-8A (Cone 8)	J1/:		
ANGANESE (Mn) 0.2%	Lely 1250 / Vicon	visi		

OIN (	-e)		<b>3</b> 72
ANG	NÉSE (Mn)	Lely 1250 / Vicon	visit www.bestfertilizer.com
	I from Monoammonium Phosphate, Ammonium Sulfate, Muriate of Potash, ysulfate and Manganese Oxysulfate.	As a guide, set spreader at the setting indicated by the settings provided are only a guide and depend	
orma ernet	tion regarding the contents and levels of metals in this product is available on the at http://www.aapfco.org/metals.htm	of your spreader. Check spreader calibration often product off sidewalks, patios, driveways, etc. This fe	ertilizer contains iron which will stain concrete
<b>A</b>	NOTICE: This product can expose you to Cadmium (Ca), which is known	and painted surfaces. If it comes in contact with corflushing with water. If concrete is wet when contact of	

30 days between applications. The maximum seasonal rate is 6.66 pounds of product per 1,000 sq.ft.

4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

to the State of California to cause cancer, birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov.

Have the state of California to cause cancer, birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov.

## It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

• Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. • Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floratam' St. Augustinegrass in Florida. • Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application. • Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result. • Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Turf Supreme® Plus Trimec controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns as listed. In addition, a 16-6-8 fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. The maximum broadcast application rate is 3.33 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of

 Mow lawn to normal height 1 to 2 days before application. 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4). 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.

WEEDS CONTROLLED								
Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy	Broadleaf plantain Buckhorn plantain Bulbous buttercup Bull thistle	Chicory Clover, crimson Clover, white Cocklebur	Goldenrod Ground ivy Henbit Hoary cress	Narrowleaf vetch Pearlwort Poison ivy Poison oak	Redstem filaree Rough cinquefoil Rough fleabane Russian pigweed	Spotted spurge Tall Nettle Tumble Pigweed Venice mallow		
Black medic Black mustard Blackseed plantain	Burdock Burning nettle Burclover	Common mullein Creeping buttercup Curly dock	Hoary plantain Kochia Knotweed, prostrate	Prickly lettuce Prostrate pigweed Prostrate knotweed	Russian thistle Scarlet pimpernel Scotch thistle	Wavyleaf, thistle White mustard		
Blessed thistle Bloodflower milkweed	Buttonweed, common Canada thistle	Daisy, English Dandelion, common	Lambsquarters Mallow, common	Prostrate knotweed  Prostrate spurge  Puncturevine	Shephard's purse Smooth dock	Wild carrot Wolly Plantain Yarrow, common		
Bracted plantain Bristly extendue	Carpetweed common	Dichondra Dollarweed	Morningglory, tall Mustard tumble	Purslane, common	Smooth pigweed			

## Do not contaminate water, food, or feed by storage and disposal.

FOR USE ONLY AS DIRECTED.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. Distributed By: J.R. SIMPLOT COMPANY, P.O. BOX 198, LATHROP, CA 95330







