

IDENTEY THE PROBLEM—Browsing by deer or elk can usually be determined by the uneven tears on leaves and stems, pulling plents from the ground and stripping of stems; tracks and elongoted dropping may be evident. Hares and rabbits clip stems at an angle within about 2 feet above ground or higher on anow; tracks and round droppings may be evident without careful examination of the site. Some rabbits and some rodents will also peel bark from the stems. Moles leave rounded piles of soll and ridges in soil.

PREVENTIVITE PROBLEMS—Deer and elk usually damage conferous trees in winter soon after planting, or, in apring after new growth begins. Deer damage landscape plants, gardens, and orchards in spring, summer and winter. Here and rabbit damage is usually most severe on conifers and shrubs in winter, and on garden and landscape plants in summer. Moles also burrow around plants and bause roots to dry out.

WHEN TO APPLY SEADIST WIRDLIFF CONTROLLAND*—Prior to winter damage, apply the granular dust to those plants and plant parts expected to be damaged. During the growing sesson apply efter 1 or 2 inches of new foliage has developed. If some damage has already occurred, apply the granular dusts to both damaged and undamaged parts. For moles, aprinkle Seadust in new planting holes. For established plants, place Seadust in shovel silices around plant to intercept burrows.

HOW TO APPLY SARIST WRDLER CONTROLLANT "—To protect plants for a few days or longer during dry weather, apray follage and stems with water and dust with Seadust Wildlife Controllant". This is particularly effective for garden plants against robbit damage. Additional applications may be needed as new follage develops. To protect plants for periods of 2 months or more, including periods of rain or irrigation, spray dry follage and stems with Bond Spreader Sticker (CAS No. 36208-50005) at 3% to 5% solids in water and immediately apply Seadust Wildlife Controllant" onto the spreader sticker and let dry 2 hours or more. CAUTION: Adhesive sprays can harm plants if not epplied according to the label. Keep Wildlife Controllant and Adhesive Sprays out of the reach of children. For Moles: Use probe or drill to locate 3 burrows around fresh mounds and pour 1/4 to 1/2 cup of Soadust in each hole. Repeat if there are new mole mounds after 1 week. Typically apply about 11/2 pounds Seadust per 2500 square feet Do not exceed 180 libs. per acre.

STORAGE AND DEPOCAL...Keep dry in air tight gleas or heavy plastic containers in a dark location at room temperature. Dispose of according to state regulations.

SUPPLEMENTAL AMESINE LASE.—Application: Dust wet foliage or stems except during heavy rain or irrigation. Formulation: Water activated dry adhesive. (Dow Cellosize ER 4400 and feed grade urea). EPA exempted (25b)

This product has not been registered by the Linited States Environmental Protection Agazer, Wildlife Scrivices Company, Lnc.
represents that this product qualifies for earnysten from regists allon under the Foderst researchide,
fungionals, and Andersticion Act (FARA). This product is manufactured by:

Wildlife Services Company, Inc. 5227 Gifford Road SW, Olympia, WA 98512

> EMAIL: seadust@comcast.net Phone/Fax: 360-352-5150

MADE IN THE U.S.A.

US. PATENT NO. 6,652,870 B2