

Contains
Glyphosate

GORDON'S®

PRONTO®

FAST ACTING

**Weed & Grass
KILLER CONCENTRATE**

**Makes Up To
24 Gallons**

- Lawn Renovation
- Garden Preparation
- Fenceline Cleanup
and more!



Rain Fast In Hours

EVEN KILLS THE ROOTS!

ACTIVE INGREDIENT:
Isopropylamine salt of
glyphosate* 25.0%

OTHER INGREDIENTS: 75.0%

TOTAL 100.0%

*N-(phosphonomethyl)glycine

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS:
ONE U.S. QUART
(32 Fl. Oz.)
954/2-2010

CAUTION

*See attached booklet for complete
Precautionary Statements and Directions for Use
(including First Aid and Storage and Disposal).*

GORDON'S®

PRONTO®

FAST ACTING

**Weed & Grass
KILLER CONCENTRATE**

Contains Glyphosate


Makes Up To 24 Gallons

Even Kills The Roots!

***See inside pages for
Precautionary Statements
and Directions for Use
(including First Aid and
Storage and Disposal).***

954/8-2015 AP052115
EPA REG. NO. 2217-846
EPA EST. NO. 2217-KS-2

 **pbi / GORDON
CORPORATION**
An Employee-Owned Company



32 OUNCES

GORDON'S®

PRONTO®

FAST ACTING

Weed & Grass
KILLER CONCENTRATE

- ***Lawn Renovation***
- ***Garden Preparation***
- ***Fenceline Cleanup and more!***

ACTIVE INGREDIENT:

Isopropylamine salt of glyphosate* 25.0%

OTHER INGREDIENTS: 75.0%

TOTAL 100.0%

*N-(phosphonomethyl)glycine

KEEP OUT OF REACH OF CHILDREN

CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give

First Aid *(Continued)*

	<p>artificial respiration, preferably mouth-to-mouth if possible.</p> <ul style="list-style-type: none">• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

ENVIRONMENTAL HAZARDS:

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater and rinsate.

PHYSICAL OR CHEMICAL HAZARDS:

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets out of treated areas until sprays have dried.

HOW IT WORKS:

Facts and Tips

- Pronto[®] Fast Acting Weed & Grass Killer Concentrate is a multipurpose herbicide designed to control a broad spectrum of emerged annual and perennial weed species. This product will control unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees, shrubs, fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like — they may die too. Not recommended for spot weed control in lawns since this product kills lawn grasses.
- Plants that have not emerged at the time of application will not be controlled.
- Time to symptoms — Best results will be obtained when the weed species are actively growing. Visible effects are gradual — wilting and yellowing, advancing to complete browning and root destruction. Weeds usually yellow and wilt within 2 to 4 days and plant death occurs in 1 to 2 weeks.

Product Description and Spray Coverage: This product is a nonselective herbicide that will kill all vegetation contacted. This herbicide kills unwanted plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots. Avoid direct spray or drift onto desirable plants, lawn grasses,

shrubs and trees. Protect desirable plants and vegetation with a piece of plastic or cardboard. If desirable plants are accidentally sprayed, wash plants with water immediately. Any glyphosate not absorbed by the plant is broken down into natural materials. This product has no soil residual activity and thus will not affect nearby untreated plants.

WHERE TO USE:

For General Weed Control and Spot Spraying

- Use to prepare sites for large plantings, for planting flower beds, for planting fruit trees, ornamentals, vegetables, garden sites and to renovate lawns.
- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.
- Along ornamental lawns and turfgrass. Edging along/around lawns/yards and landscape plantings.
- In areas for lawn replacement or lawn renovation/replacement.
- Around or near ornamental/shade trees, fruit and nut trees, grapevines, shrubs, groundcovers, flower beds, mulched landscape beds.
- For garden plot preparation, vegetable plot preparation, and small fruits and berries plot preparation.
- For flower garden preparation.
- Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.

- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fence-lines, around shelter belts and other areas where nonselective weed control is desired.

TOUGH WEED CONTROL

(ANNUALS, PERENNIALS, BRUSH)

CONTROLS WEEDS AND GRASSES INCLUDING LAWN AND TURFGRASS SPECIES: This product will provide full or partial control (suppression) of these lawn and turfgrass species:

Bahiagrass

Bentgrass, colonial, creeping and redtop

Bermudagrass, common and hybrid

Bluegrass, annual, rough, and Kentucky

Bromegrass, downy and smooth

Buffalograss

Canarygrass, reed

Carpetgrass

Centipedegrass

Fescues, chewings, hard, red, tall, sheep

Foxtail, green, yellow, creeping, meadow, giant

Johnsongrass

Kikuyugrass
Orchardgrass
Paspalum *spp.*
Ryegrass, annual, Italian, perennial, and hardy
St. Augustinegrass
Zoysiagrass or Japanese lawngrass

WOODY SPECIES:

This product will provide full or partial control (suppression) of these woody species:

Alder	Mock orange*
Blackberry	Manzanita
Barberry	Pine
Caenothus <i>spp.</i>	Poison ivy
Cherry	Poison oak
Coyote brush	Poison hemlock
Dewberry	Oak
Elm	Quaking aspen
Elderberry	Raspberry
Honeysuckle	Sumac
Hickory	Trumpet creeper
Kudzu	Wild blackberry
Lantana	Western red cedar*
Maple	Willow
Multiflora rose	*Not for use in California

ANNUAL AND PERENNIAL WEED SPECIES:

This product will provide full or partial control (suppression) of these annual and perennial weed species:

Artichoke, Jerusalem

Barnyardgrass

Beggarweed

Bittercress

Black medic

Broadleaf plantain

Buckhorn plantain

Bull thistle

Burclover

Canada thistle

Chickweed

Clover

Cocklebur

Common groundsel

Common mullien

Common ragweed

Coral-bead

Crabgrass

Creeping beggarweed

Creeping Charlie

Curly dock

Dallisgrass

Dandelion

Evening primrose

Fall panicum

False dandelion

Fennel

Fiddleneck

Field bindweed

Field pennycress

Field sandbur

Filaree

Florida pusley

Garden spurge

Goosegrass

Green foxtail

Guineagrass

Hemp sesbania

Henbit

Horsenettle

Horseradish

Horseweed/marestail

Iceplant

Ironweed

Knapweed

Knawel

Knotweed

Lambsquarters

London rocket

Lovegrass

Mallow

Mayweed

Milkweed

Mouseear chickweed

Nimblewill

(cont. on next page)

ANNUAL AND PERENNIAL WEED SPECIES: (cont.)

Oleander*	Toadflax
Oxalis	Torpedograss
Pampas grass	Trailing lantana
Pennsylvania smartweed	Tumble mustard
Pigweed	Vaseygrass
Plantain	Velvetleaf
Prickly lettuce	Virginia-creeper
Primrose	White clover
Prostrate spurge	Whitetop
Puncturevine	Wild barley
Purple cudweed	Wild carrot
Purple nutsedge	Wild geranium
Purslane	Wild morningglory
Quackgrass	Wild mustard
Ragweed	Wild oats
Red clover	Wild sweet potato
Red sorrel*	Witchgrass
Redroot pigweed	Yarrow
Shepherd's purse	Yellow nutsedge
Smartweed	Yellow oxalis
Smooth catsear	Yellow starthistle
Smooth pigweed	
Sowthistle	
Spotted spurge	
Spurge	
Sunflower	
Tansy ragwort	
Tansy mustard	
Texas panicum	
Thistle	
Timothy	

*Not for use in California

WHEN TO USE:

Apply to actively growing weeds when the vegetation is green. Use when the wind is calm to avoid drift. Make applications on warm sunny days when the day-time temperature is above 60°F. Weed control is accelerated by warm sunny weather. Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks. Rain or watering 2 hours after application will not wash away effectiveness. Hard-to-control weeds and thick lawn grasses (like bermudagrass) may require a second application.

HOW TO USE:

This product is formulated for spot treatment of individual weeds and grasses or for treatment of large areas as instructed on this label.

To apply this product use any of the following equipment:

- Plastic, aluminum or stainless steel tank sprayer
- Ortho Dial 'N Spray®
- Ortho Spray-Ette®
- A hose end sprayer recommended for herbicides.
- Hand-trigger pump sprayer.
- **Do not use or store in galvanized or unlined steel sprayer, or through any irrigation system or sprinkling can.**
- Apply to evenly and thoroughly wet weeds and grasses.
- Use a tank type sprayer with a coarse spray nozzle for spot treatments.

- Use the Ortho Dial 'N Spray® or other hose end sprayer to treat large areas such as for lawn renovation or to prepare sites for planting flower beds or vegetable gardens.
- Treated weeds begin to wilt in about 2 to 4 days with complete kill of the weed including the root in 2 to 4 weeks.
- This product does not move or build up in the soil to effect untreated plants growing nearby.
- Ornamental plantings can be made 3 days after treatment. Vegetable gardens can be seeded or planted 3 days after treatment.
- For lawn replacement, lawns may be replanted 1 week after spraying.
- Measure the correct amount of this product and water to add to sprayer from the directions on next page.

TABLE 1. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> • Use 2.25 fl. oz. of this product in 1 gallon of water. 	<ul style="list-style-type: none"> • Use 4.5 fl. oz. of this product in 1 gallon of water. 	300 sq. ft. (30 ft x 10 ft)
Ortho Dial 'N Spray [®] Hose End Sprayer	<ul style="list-style-type: none"> • Add 2.25 fl. oz. of this product to sprayer jar. • DO NOT ADD WATER. • Set dial to 3 oz. and apply. 	<ul style="list-style-type: none"> • Add 4.5 fl. oz. of this product to sprayer jar. • DO NOT ADD WATER. • Set dial to 5 oz. and apply. 	300 sq. ft.

(Continued on next page)

TABLE 1. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS (Cont.)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Ortho Spray-Ette® Hose end Sprayer	<ul style="list-style-type: none"> • Add 8 fl. oz. of this product to sprayer jar. • Fill to 4 gallon mark. 	<ul style="list-style-type: none"> • Add 16 fl. oz. of this product to sprayer jar. • Fill to 4 gallon mark. 	1200 sq. ft. (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> • Use 1 tbs. (0.5 fl. oz.) of this product in 24 fl.oz. of water. 	<ul style="list-style-type: none"> • Use 2 tbs. (1.0 fl. oz.) of this product in 24 fl.oz. of water. 	Use for spot treatments or single weeds

EQUIVALENT MEASUREMENTS
1 teaspoon (tsp.) 1 tablespoon (tbs.) = 3 teaspoons (tsp.) 1 fluid ounce (fl.oz.) = 2 tablespoons (tbs.) 2 fluid ounces (fl.oz.) = 4 tablespoons (tbs.)

This product in these container sizes will treat these approximate areas:

CONTAINER SIZE	ANNUAL WEEDS (Treated Area, Sq. Ft.)	PERENNIAL WEEDS (Treated Area, Sq. Ft.)
1 pint	2,400 sq. ft.	1,200 sq. ft.
1 quart	4,800 sq. ft.	2,400 sq. ft.

CLEANING APPLICATION EQUIPMENT:

To clean sprayer after use, rinse spray equipment thoroughly and flush all sprayer components with water 3 times after use. Spray rinsate on bare soil or gravel surface. Do not apply rinsate to treated weeds. Only after a very thorough cleaning can the sprayer or spray equipment be used for application of other products.

WHERE TO USE

For General Weed Control: Use near fences, in paths, patios, sidewalks, curbs and driveways, around ornamental trees, shrubs, flower beds and plantings, around buildings. Control weeds in brick and gravel

walks. Use for trimming and edging landscapes. NOT recommended for spot weed treatment in lawns, since glyphosate kills all green plants, including lawn grasses.

Spot Treatments: Spray directly on actively growing weeds in and around flower beds, ornamental trees, fruit and nut trees, grapevines, shrubs, fences, driveways and walkways. Do not spray fruit and nut trees or grapevines directly.

For Landscaping:

- Prepare areas for planting of ornamentals, trees, shrubs, desert landscapes, rock gardens, flower beds or similar plantings.
- Delay mowing, tilling, or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results. However, if the treated vegetation is not disturbed, planting or transplanting of trees, flowers, shrubs, bedding plants may be done at one (1) day after application.

For Lawn Renovation and Lawn Replacement:

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock. This product may be applied before planting, seeding, sprigging, plugging, or sodding.
- Use this product to remove the existing lawn and kill weeds and grasses. Uniform and complete spray coverage of all existing vegetation is required.
- Bermudagrass, nutsedge, torpedograss and deep rooted perennials may require a second treatment in 5 to 7 days.

- For best results, apply in spring or fall when daytime temperatures are at least 60°F.
- Best results are obtained when one mowing is skipped before and after the application of this product.
- If the lawn is dry, water lawn before applying this product. Water again 2 to 3 days after application.
- Hose end sprayers and tank sprayers are appropriate for localized spot treatments and broadcast (entire) applications on small turfgrass areas, e.g. ~10,000 sq. ft. Refer to Table 1 for the spray preparation using hose end sprayers and tank sprayers.
- Conventional sprayers fitted with spray booms and spray guns are appropriate for broadcast (entire) applications on large turfgrass areas, e.g. ~20,000 sq. ft. Refer to Table 2 for the spray preparation using power spray equipment suitable for large turfgrass areas.
- Seeding, sprigging, plugging or sodding may be done at seven (7) days after application of this product. Prepare the soil for planting by raking, roto-tilling, vertical cutting, or other tillage method.
- Rake up and remove loosened thatch or debris.
- Mix in starter fertilizer and soil amendments if needed. Establish the lawn according to local cultural practices.

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
Lawn & turf seedbeds (prior to establishment):	Many annual and perennial grasses and broadleaf weeds, including annual bluegrass, crabgrass, lambsquarters and ragweed.	Annual weeds: 1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
		Perennial weeds: 5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bahiagrass	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bentgrass, colonial creeping	2.7 - 3.6	≤ 25	2.0 - 2.6	< 1

(Continued on next page)

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
	Bermudagrass, common and hybrid	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bluegrass, annual rough and Kentucky	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Bromegrass, downy and smooth	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Orchardgrass	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Canarygrass, reed	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1

(Continued on next page)

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
	Fescue, tall	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1
	Fescues, chewings, hard, red, sheep	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Johnsongrass and ryegrass annual, perennial	1.8 - 5.5	≤ 25	1.3 - 4.0	< 1
	Kikuyugrass	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1
	Quackgrass and yellow nutsedge	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1

(Continued on next page)

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./1000 sq. ft.	Spray Volume, Gals./1000 sq. ft.
	Other perennial grasses listed on this label	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1

PRECAUTIONS, RESTRICTIONS & COMMENTS FOR LAWN RENOVATION AND LAWN REPLACEMENT

SITE PREPARATION (preplant, preemergence, renovation): This product may be applied before planting or renovation of turf. Applications must be made prior to the emergence of the turf to avoid turf injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm season grasses, such as bermudagrass, summer or fall applications provide best control.

Do not disturb soil or underground plant parts before treatment.

Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

This product does not provide residual control.

FARMSTEADS AND FENCEROWS: This product will control kochia, multiflora rose, thistle in areas where total vegetation control is desired. Specific instructions for farmsteads are presented in Table 3.

TABLE 3. SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelter belts and other area where nonselective weed control is desired.	Annual and perennial weeds, including crabgrass, foxtail, Johnsongrass, kochia, morningglory, multiflora rose, pigweed, trumpet creeper, thistle, woody brush.	Annual weeds: 1.8 - 3.6	3 - 40	1.3 - 2.6	≤ 1.0
		Perennial weeds: 5.5 - 9.5	3 - 40	4.0 - 7.0	≤ 1.0
		Woody brush & trees: 5.5 - 9.5	3 - 40	4.0 - 7.0	≤ 1.0

See next page for Table 3 – Precautions, Restrictions & Comments

TABLE 3. SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT (Cont.)

PRECAUTIONS, RESTRICTIONS & COMMENTS:

SITE PREPARATION (preplant, preemergence, renovation):

BARE GROUND: Repeated applications of this product may be used on emerged weeds to maintain bare ground.

NONSELECTIVE POSTEMERGENCE:

Annuals and Perennials: Apply to foliage of emerged weeds. Apply at early growth stages of annual weeds and when perennial weeds are approaching maturity.

Woody brush and trees: Apply this product after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when applications are made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Reduced performance may result if fall treatments are made following a frost.

GARDENS (VEGETABLES, HERBS, SMALL FRUITS AND BERRIES):

This product may be applied before planting or transplanting of vegetables (garden crops), herbs and small fruits and berries. This product may be used as a broad spectrum postemergent herbicide prior to planting to control annual and perennial weeds including field bindweed, Johnsongrass, or dallisgrass.

Spray the undesirable vegetation on a spray-to-wet basis. Uniform and complete spray coverage of all existing vegetation is required.

RESEEDING AND REPLANTING IN SPRAYED AREAS

ORNAMENTALS AND FLOWERS: Reseed or replant one (1) day after application.

VEGETABLES: Wait three (3) days before planting or transplanting the vegetables listed below:

Beans	Endive
Beet greens	Garlic
Broccoli	Gourds
Brussels sprout	Honeydew melon
Cabbage	Honeyball melon
Cantaloupe	Horseradish
Carrot	Jerusalem artichoke
Cauliflower	Kale
Casaba melon	Kohlrabi
Chard (Swiss)	Leek
Chicory	Lentils
Collards	Lettuce
Corn	Mango melon
Crenshaw melon	Melons
Cucumber	Muskmelon
Eggplant	Mustard greens

Okra
Onion
Rape greens
Parsley
Parsnips
Peas
Pepper
Persian melon
Potato
Pumpkin
Radish
Rhubarb

Rutabaga
Spinach
Summer squash
Sweet potato
Tomatillo
Tomato*
Turnip
Watercress
Watermelon
Winter squash
Yams

*Tomatoes planted from seed only. Wait thirty (30) days for transplants. For vegetables not listed or for tomato transplants, wait thirty (30) days before planting.

SMALL FRUITS AND BERRIES: Blackberry, black raspberry, blueberry, boysenberry, currant, dewberry, elderberry, gooseberry, huckleberry, loganberry, olallieberry, and red raspberry may be planted by seed, rooted cutting, or transplant 7 days after spraying this product. Wait thirty (30) days after application before planting of all other unlisted small fruits and berries.

FRUIT TREES, NUT TREES, CANEBERRIES OR GRAPEVINES: If this product is used to control weeds around fruit or nut trees, caneberreries or grapevines, wait twenty-one (21) days before harvesting and eating the fruits or nuts.

Use the following table to determine the amount of this product to mix with water for use in gardens (vegetables, herbs, small fruits and berries). Close cap tightly after use.

TABLE 4. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> • Use 2.25 fl. oz. of this product in 1 gallon of water. 	<ul style="list-style-type: none"> • Use 4.5 fl. oz. of this product in 1 gallon of water. 	300 sq. ft. (30 ft x 10 ft)
Ortho Dial 'N Spray® Hose End Sprayer	<ul style="list-style-type: none"> • Add 2.25 fl. oz. of this product to sprayer jar. • DO NOT ADD WATER. • Set dial to 3 oz. and apply. 	<ul style="list-style-type: none"> • Add 4.5 fl. oz. of this product to sprayer jar. • DO NOT ADD WATER. • Set dial to 5 oz. and apply. 	300 sq. ft.

(Continued on next page)

TABLE 4. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES) (Cont.)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Ortho Spray-Ette® Hose end Sprayer	<ul style="list-style-type: none"> • Add 8 fl. oz. of this product to sprayer jar. • Fill to 4 gallon mark. 	<ul style="list-style-type: none"> • Add 16 fl. oz. of this product to sprayer jar. • Fill to 4 gallon mark. 	1200 sq. ft. (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> • Use 0.5 fl. oz. (1 tbs.) of this product in 24 fl. oz. of water. 	<ul style="list-style-type: none"> • Use 1 fl. oz. (2 tbs.) of this product in 24 fl. oz. of water. 	Use for spot treatments or single weeds

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Protect container from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED

WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Pronto® is a registered trademark of PBI/GORDON Corporation

Ortho®, Ortho Dial 'N Spray®, and Ortho Spray-Ette® are registered trademarks of Monsanto Company.

954/8-2015 AP052115
EPA REG. NO. 2217-846
EPA EST. NO. 2217-KS-2

**MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com**

THIS PAGE IS BLANK INTENTIONALLY.

THIS PAGE IS BLANK INTENTIONALLY.

THIS PAGE IS BLANK INTENTIONALLY.

**STOP! READ THE ENTIRE LABEL
FIRST. OBSERVE ALL
PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PHYSICAL OR CHEMICAL HAZARDS:

Combustible. Do not use or store near heat or open flame.

954/8-2015 AP052115
EPA REG. NO. 2217-846
EPA EST. NO. 2217-KS-2

**MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com**

PEEL HERE TO OPEN ▲