



CONTROLS:



Fleas



Cockroaches



Spiders



Flies



Bed Bugs

# VIPER RTU

## READY-TO-USE INSECTICIDE

- With Country Fresh Fragrance
- Provides 60-Day Control of Fleas, Cockroaches and Stored Product Pests
- For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs
- For Control of Pests on Non-Commercial Livestock

### ACTIVE INGREDIENT:

Permethrin (CAS #52645-53-1).....0.25%

**OTHER INGREDIENTS:**.....99.75%

**TOTAL:**.....100.00%

*Cis/trans* ratio: min. 35% (+/-) *cis* and max. 65% (+/-) *trans*.

EPA Reg. No. 4-393-53883

EPA Est. No. 53883-TX-002


**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Inside Booklet for Additional Precautions.

A **CONTROL SOLUTIONS INC.** PRODUCT

**NET CONTENTS: 1 GALLON**

LIFT  
HERE



**READ ENTIRE LABEL BEFORE USE.****PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter rain or storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:** Except when applying to pets, companion animals or non-commercial livestock, do not enter or allow others to enter until sprays have dried. Except when applying to pets, companion animals or non-commercial livestock, do not apply this product in a way that will contact adults, children or pets, either directly or through drift. Do not use on outdoor residential misting systems. Do not use as a broadcast treatment on outdoor surfaces at residential sites, including nurseries, day care centers, schools, hospitals and nursing homes. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Do not water the treated area to the point of runoff. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation.

**STORAGE AND DISPOSAL**

**Pesticide Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.  
**Pesticide Disposal & Container Handling:** If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. If empty - Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

**SHAKE WELL BEFORE USING****INDOOR APPLICATION**

For spot application only. Do not use as a space spray indoors.

**SURFACE SPRAYING:** FOR CATS: DO NOT USE IN CONTROL OF: COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDS, CENTIPEDS, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, FLOWER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOOR BEETLES, RUST RED FLEA BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

**DO NOT CONTROL CARPET BEETLES:** Apply as a spot treatment to edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

**DO NOT CONTROL BEDBUGS:** Spray lightly around tufts and seams of mattresses. Do not spray entire mattress. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**DO NOT CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE):** Apply as a spot treatment to infested areas, dog beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding must be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas. Do not spray areas inhabited by cats.

**DO NOT CONTROL CLOTHES MOTHS:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

**DO NOT KILL LICE AND LOUSE EGGS:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 12 inches. Treat only those garments and parts of bedding and furniture that cannot be either laundered or dry cleaned. For mattresses, spray lightly around tufts and seams. Do not spray entire mattress. Allow all treated articles to dry thoroughly before use.

**ON COMPANION ANIMALS AND NON-COMMERCIAL LIVESTOCK:** Protects cattle (beef and dairy), hogs (swine) and horses from Stable Flies, Horn Flies, Face Flies, Horse Flies, Deer Flies, Blood-Sucking Lice, Mosquitoes and Gnats.

**CATTLE SPOT APPLICATION TO BACK, FACE, LEGS, TAIL & EARS:** Apply 2 fl. oz. of product per animal to infested areas. For blood-sucking lice, use a spray stiff brush to get the spray to the base of the hair. Reapply as needed.  
**HOGS BODY SPRAY:** Mix 2 fl. oz. of product in 1 gallon (½ fl. oz. in 1 quart) of water. Apply as a light mist to the hair thoroughly, using 1-2 quarts of dilute solution per animal. For face flies, spray face and head, but do not spray into ears. Repeat treatment in 10 days, if needed.

**HOGS (SWINE): SPOT APPLICATION TO BACK, FACE, LEGS, TAIL & EARS:** Apply 2 fl. oz. of product per animal to infested areas. For blood-sucking lice, use a spray stiff brush to get the spray to the base of the hair. Reapply as needed.

**WHOLE BODY SPRAY OR DIP:** Mix 2 fl. oz. of product in 1 gallon of water (½ fl. oz. in 1 quart). Apply as a light mist to wet the hair thoroughly, using 1-2 quarts of dilute solution per animal. For face flies, spray face and head, but do not spray into ears. Repeat treatment in 14 days, if needed.

**HORSES: SPOT APPLICATION TO BACK, FACE, LEGS, TAIL & EARS:** Apply 2 fl. oz. of product per animal to infested areas. For face flies, spray face and head, but do not spray into eyes. Reapply as needed. **WHOLE BODY WASH OR DIP:** Mix 2 fl. oz. of product in 1 gallon of water. Apply to wet the hair thoroughly. Reapply as needed.

**POLYTRY:** To Control Poultury Lice, Bedbugs and Mites on poultry and in non-commercial poultry houses. **POLYTRY:** apply as a fine mist, using approximately 1 gallon per 75 birds (1 quart per 20 birds) (1 pint per 125 sq. ft.) (7 fl. oz. per 500 sq. ft.) using wetting rooster, wails and cages. For bedbugs and mites, spray crevices of roost poles, cracks in walls and cracks in nests where these insects hide. Do not treat poultry litter. Reapply in 1 to 2 weeks, if required.

**DOGS: USE ONLY ON DOGS. DO NOT USE ON CATS OR KITTENS.** Do not use on puppies less than twelve weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals.  
**TO CONTROL FLEAS, TICKS, AND LICE:** Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Do not spray animal's genitalia. Do not reapply product for 7 days.  
In case of an emergency, call 1-800-548-2423 (Animal Poison Control Center).

**OUTDOOR APPLICATION**

| Plant:  | Pests:  | How To Use:  |
|---|---|--|
| Ageratum, Aster, Ardisia, Azalea, Baby's Breath, Begonia, Birds-nest Fern, Bleeding Heart, Camellia, Carnation, Chrysanthemum, Coleus, Common Ninkerbak and Snowberry, Crown of Thorns, Cockscomb, Crassula, Cyclamen, Dahlias, Delphiniums, Dogwood, Dracaena, Dumbcane, English Ivy, Euconymus, Excaxum, Fuchsia, Geranium, Gladiolus, Grape Ivy, Grape Vines, Gold Bells, Hypoestes, Ivy, Juniper, Kentia palms, Laurel, Lilac, Marigold, Mock-Orchid, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Pinsetta, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rhododendron, Rose, Rubber plants, Schellera, Snapdragon, Snake Plant, Statice, Stocks, Velvet Plant, Verbena, Violets, Wandering Jew, Zinnia | Ashids, Armyworms, Bagworms, Cabbage Worms, Cicadas, Clover Mites, Corn earworms, Cutworms, Exposed Trifles, Fall Cankerworms, Fall Webworms, Fungus Gnat, Inchworms, Japanese Beetles, Lace Bugs, Leafminers, Leafhoppers, Mealybugs, Pavement Ants, Scale Crawlers, Spring Cankerworms, Spider Mites, Whiteflies, Rose Chafers, Psyllids, Scale Insects | Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of spotting. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection. |

**TO HELP CONTROL TICKS** in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**FOR CONTROL OF GARDEN INSECTS:** When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur. To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. Buyer's Guarantee Limited to Label Claims. For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents) call National Pesticide Information Center at 1-800-858-7378.