



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container. Keep in an area inaccessible to children and pets.

PESTICIDE DISPOSAL: If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly full: Call your local solid waste agency for disposal instructions. Never place unused product down any sewer or intend drain.

GENERAL INFORMATION

H-YIELD® TURF RANGER INSECT CONTROL GRANULES CONTAINING DELTAMETHRIN is a broad spectrum residual insecticide for the control of certain insects in Home Lawns. Treat as directed - water in granules with 1 inch of water. This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply on saturated soil. Do not allow runoff to occur. Do not contaminate ornamental fish ponds.

In New York State, this product may not be applied to lawns that directly border coastal marshes. For control of Turf Insects other than Fleas and Ticks in New York State, a second application may be made after two weeks if target insect activity persists.

Not for use in commercial greenhouses or nurseries.

Not for use on sod lawns, grass grown for seed or on golf course turf.

Food plants such as spuds and measuring crops must not be used for food purposes after use in measuring pesticides. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

HOME LAWN RECOMMENDATIONS: Use **H-YIELD® TURF RANGER INSECT CONTROL GRANULES CONTAINING DELTAMETHRIN** to control specified Pests of Lawns. Apply **H-YIELD® TURF RANGER INSECT CONTROL GRANULES CONTAINING DELTAMETHRIN** to give complete and uniform coverage of Pest Infested Areas. Applications should be made using the rates shown below.

VEGET	DOSEAGE OF PRODUCT	APPLICATION PROCEDURE
Ants, Armyworms, Bermudagrass Miles, Billbugs (Adult), Boxelder Bugs, Caterpillars, Chiggers, Chinch Bugs, Clover Miles, Cockroaches, Grasshoppers, Japanese Beetle, Japanese Beetle Larvae, Japanese Beetle Pupa, Japanese Beetle Weevils, Eastern Skipper, Field Crickets, Flea Snapper, Fire Ants, Flea Beetles, Flea, Grasshoppers, Hyperisks, Japanese Beetles, Lawn Moths, Leafhoppers, Mealy Bugs (Adult), Milkweed Bugs (Adult), Onion Thrip, Onion Thrip Larvae, Onion Thrip Nymph, Onion Thrip Pupa, Snoutbeetles, Spiders, Saltmarshes, Sprinklers, Ticks (including Lyme disease vector), Vegetable Weevils	2 lbs to 3 pounds per 1,000 sq. ft.	Apply this product uniformly to Home Lawns, under shrubsbery and other ornamental/ornamental grassy areas, and areas adjacent to any body of water. Immediately after application, water the lawn thoroughly with half an inch of water.
European Crane Fly Larvae (Ladache Jackants)	2 to 3 pounds per 1,000 sq. ft.	Apply in early April when insects are young.
Chinch Bugs	2 to 3 pounds per 1,000 sq. ft.	Start treatment when temperatures reach 60°F or above. A second application may be necessary in 6 weeks if insect activity continues.

APPLICATION EQUIPMENT AND INSTRUCTIONS: This product can be applied with a drop or broadcast (starry) spreader. For best results, apply this product evenly and uniformly over the target (lawn) area avoiding streaking, skips or overlaps.

Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 3 lbs. per 1000 sq. ft., and your area is 2000 sq. ft., then place 6 lbs. of product in the hopper and set the spreader setting at 3.1/4 (per table below) if using an AgriSource drop spreader.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the target area and then applying the remaining one-half of this product at a right angle to the previous application direction.

NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will effect the rate of application.

Spreader	Spreader Settings		
	Lbs. of product per 1,000 sq. ft.	2 lbs.	3 lbs.
Scotta/Republic AgriSource (Drop)	2 1/2	3 1/4	
Scotta/Republic Speedy Green (Broadcast)	2 3/4	3 1/2	
Farmer's Edge Earthway EV-V-Spread (Broadcast)	15	14	

FLEA AND TICK CONTROL (WITBOMB): For control of Fleas and Ticks (including, but not limited to, Boxer Tick, Black-legged Tick, Brown Dog Tick, Deer Tick, Lone Star Tick, Western Black-legged Tick, Head Tick), a regular treatment program should be followed. Treat areas monthly as directed above when Pests first appear in late Spring (May-June) and continue until late Summer (September-October).

SOIL WORMS: If Lawn Moths (and Webworms/Adults) are noticed at large numbers of ducks, or at first sign of damage to lawn, treat entire lawn every

PERIMETER TREATMENT: To help prevent infestation of buildings by Ants, Boxelder Bugs, Brown Dog Ticks, Centipedes, Cockroaches, Clover Miles, Crickets, Field Cricket, Fire Ants, Fleas, Scorpions, Spiders, Spittlebug, Springtails (Collembola) and Ticks, treat a band 5 to 10 feet wide around and adjacent to the building where pests are active and may find entrance. Treat at a rate of 1 to 1 1/2 pounds per 500 sq. ft. or 2 to 3 pounds per 1,000 sq. ft.

SPOT TREATMENT: To control Ants, including Acrobat, Argentine, Big-headed, Camponotus, Harvester, Leafcutter, Little Black, Pavement and Pheidole Ants. Product should be applied to thoroughly cover the nest (1 to 2 tablespoons) and surrounded with sufficient water to dissolve granules. In addition to treating Ant nests or mounds, entry points of buildings and the soil surface along the perimeter of the building should be treated.

DEGGER TREATS: Apply 1 to 2 tablespoons of granules in and around each individual ground nest opening. Make the application at dusk when Wasps are not active.

SIMPSON FIRE ANT: Apply two tablespoons over each 8" to 10" mound and up to 2 to 3 inches to prevent escape of the queen. Immediately after application, water the area with the soil partly with a light stream of water, then cover with soil again. Do not apply in cool weather during the Spring and Fall (55-65°F) or in early morning or late evening for best results. Treat new mounds as they appear. If possible, make applications after a recent rainfall.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

In the event of a medical or chemical emergency contact:

North America 1-800-252-3824 or Worldwide Int'l. + 61-819-248-0545



ChemTel

1000-252-3824

Worldwide Int'l. + 61-819-248-0545

ChemTel

1000-252-3824

Worldwide Int'l. + 61-819-248-0545</p