

Available only at authorized

TREATS AND FEEDS **5,000** SQUARE FEET




Lawn Pro

Dealers

Weed and Feed

Weed Control Plus Lawn Fertilizer



Controls more than 50 common lawn weeds* such as dandelion, clover and chickweed
*as listed

Builds thick green lawns

ACTIVE INGREDIENTS:	
2,4-D†	1.18%
Mecoprop-p††	0.59%
OTHER INGREDIENTS:	98.23%
Total 100.00%	

*0.0118 lbs 2,4-D acid equivalent per pound of product
**0.0059 lbs MCP-p acid equivalent per pound of product


†CAS#475-7
††CAS#16484-77-8
EPA Reg. No. 538-270

EPA Est. No. 538-OH-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back for additional precautionary statements

NET WEIGHT 14.63 lb (6.63 kg) **26-0-3**

TREATS AND FEEDS **5,000** SQUARE FEET



Weed and Feed
Weed Control Plus Lawn Fertilizer



51105
3

51105
3

EPA Reg No. 538-270 (5M)
Scotts Weed & Feed Weed Control Plus Lawn Fertilizer



Weed and Feed
Weed Control Plus Lawn Fertilizer



Weed and Feed

Weed Control Plus Lawn Fertilizer

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds of 2,4-D per acre per application.

APPLICATION

- Apply when weeds are actively growing (when daytime temperatures are consistently above 60° F).
- Apply on a calm day.
- Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying this product.
- On bahia and centipede lawns, wait until after the second mowing of the year but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine and dichondra lawns. On Bermudagrass, bahia, centipede grass, zoysia lawns, under certain conditions, some temporary discoloration may result.
- Kills listed weeds

RESTRICTIONS

Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

The maximum number of broadcast applications per treatment site is two per year. A minimum of 30 days between applications is required.

WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product.

This product should not be applied if rain is expected within the next 24 hours.

IMPORTANT

WHAT TO EXPECT

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control. This product will not prevent weed seeds from producing new plants.

EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees F.

SEEDING/SODDING/SPRIGGING

Wait until the new grass has been mowed at least four times before applying Weed and Feed Weed Control Plus Lawn Fertilizer.

If you are planning to seed, sod or sprig, you should wait at least one month after you have applied Weed and Feed Weed Control Plus Lawn Fertilizer.

WEEDS CONTROLLED

aster	clover (red)	knotweed	ragweed
bindweed	clover (white)	(seedling)	red sorrel (sheep sorrel)
bittercress	cudweed	lambquarter	scarlet pimpernel
black medic	curly dock	lespedeza	shepherd's purse
buckhorn	dandelion	(seedling)	smartweed
bull thistle	dichondra	lippia (matchweed)	sow thistle
burclover	dollarweed	mallow	spurweed
buttonweed	English daisy	moneywort	vetch
Canada thistle	eveningprimrose	morningglory	wild carrot
carpetweed	filaree	mustard	yarrow
catsear (coast dandelion)	fleabane	pepperweed	yellow rocket
chickweed	ground ivy	plantain	
chicory	hawkweed	purslane	
cinquefoil	healall	pusley (Richardia spp.)	
clover (hop)	henbit		
	horseweed		

Spreader Settings This package applies 2.9 lbs of product per 1000 sq. ft.

	Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader.	Covers 6,000 sq. ft.
	Scott's® Broadcast/Rotary Spreaders	
	Each pass should be within 5 feet of each other to ensure an overlap.	3½
	Scott's® Drop Spreaders	
	Makes sure spreader wheel completely overlaps previous wheel marks.	5½
	For additional spreader settings, please call 1-800-543 TURF (8873)	

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical-resistant gloves made of any waterproof material.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Environmental Hazards: This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.

PESTICIDE DISPOSAL and CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Scott's® Weed and Feed Weed Control Plus Lawn Fertilizer 26-0-3 F 643

Guaranteed Analysis

Total nitrogen (N)	26%
7.7% ammoniacal nitrogen	
10.1% urea nitrogen	
7.5% other water soluble nitrogen*	
0.7% water insoluble nitrogen	
Soluble potash (K ₂ O)	3%
Sulfur (S)	10%
10.0% combined sulfur (S)	

Derived from: urea, methyleneurea, potassium sulfate and ammonium sulfate.

*Contains 5.6% slowly available nitrogen from methyleneureas.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Scott's No-Quibble Guarantee®


If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase, and we will mail you a refund check promptly.

Questions / Information

Consumer Helpline:
1-800-543-TURF (8873)

Visit our website:
www.scotts.com

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with directions.



©2012 The Scott's Company LLC.
World rights reserved.

Sold and Guaranteed by:
The Scott's Company
14111 Scottslawn Rd.
Marysville, OH 43041

Available only at authorized

TREATS AND FEEDS **5,000** SQUARE FEET



**Lawn
Pro**[®]

Dealers

Weed and Feed

Weed Control Plus Lawn Fertilizer

EPA Reg No. 538-270 (5M)
Scotts Weed & Feed Weed
Control Plus Lawn Fertilizer

Controls more than 50 common lawn weeds[‡] such as dandelion, clover and chickweed

[‡]as listed

Builds thick green lawns

ACTIVE INGREDIENTS:	
2,4-D†	1.18%
Mecoprop-p††	0.59%
OTHER INGREDIENTS:	98.23%
Total	100.00%

*0.0118 lbs 2,4-D acid equivalent per pound of product

**0.0059 lbs MCPP-p acid equivalent per pound of product

†CAS#94-75-7

††CAS#16484-77-8

EPA Reg. No. 538-270

EPA Est. No. 538-OH-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back for additional precautionary statements

NET WEIGHT 14.63 lb (6.63 kg)

26-0-3

DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM, DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS



Weed and Feed Weed Control Plus Lawn Fertilizer

EPA Reg No. 538-270 (5M)

Scotts Weed & Feed Weed Control Plus Lawn Fertilizer

1105
3



Weed and Feed

Weed Control Plus Lawn Fertilizer

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds of 2,4-D per acre per application.

APPLICATION

- Apply when weeds are actively growing (when daytime temperatures are consistently above 60°F).
- Apply on a calm day.
- Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying this product.
- On bahia and centipede lawns, wait until after the second mowing of the year but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine and dichondra lawns. On Bermudagrass, bahia, centipedegrass, zoysia lawns, under certain conditions, some temporary discoloration may result.
- Kills listed weeds

RESTRICTIONS

Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

The maximum number of broadcast applications per treatment site is two per year.

A minimum of 30 days between applications is required.

WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product.

This product should not be applied if rain is expected within the next 24 hours.

IMPORTANT

WHAT TO EXPECT

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control. This product will not prevent weed seeds from producing new plants.



EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees F.

7/30/2012 SSL

Page 5 of 6

Spreader Settings This package applies 2.9 lbs of product per 1000 sq. ft.

	Apply header stripes. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader	Covers 5,000 sq. ft.
	Scott's® Broadcast/Rotary Spreaders Each pass should be within 5 feet of each other to ensure an overlap.	3½
	Scott's® Drop Spreaders Make sure spreader wheel completely overlaps previous wheel marks.	5½
	For additional spreader settings, please call 1-800-543 TURF (8873)	

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical-resistant gloves made of any waterproof material.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Environmental Hazards: This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in EPA Reg No. 538-270 (5M)

STORAGE AND DISPOSITION: Scott's Weed and Feed Weed Control Plus Lawn Fertilizer

SEEDING/SODDING/SPRIGGING

Wait until the new grass has been mowed at least four times before applying Weed and Feed Weed Control Plus Lawn Fertilizer.

If you are planning to seed, sod or sprig, you should wait at least one month after you have applied Weed and Feed Weed Control Plus Lawn Fertilizer.

WEEDS CONTROLLED

aster	clover (red)	knotweed	ragweed
bindweed	clover (white)	(seedling)	red sorrel (sheep
bittercress	cudweed	lambsquarter	sorrel)
black medic	curly dock	lespedeza	scarlet pimpernel
buckhorn	dandelion	(seedling)	shepherd's purse
bull thistle	dichondra	lippia (matchweed)	smartweed
burclover	dollarweed	mallow	sow thistle
buttonweed	English daisy	moneywort	spurweed
Canada thistle	eveningprimrose	morningglory	vetch
carpetweed	filaree	mustard	wild carrot
catsear (coast	fleabane	pepperweed	yarrow
dandelion)	ground ivy	pigweed	yellow rocket
chickweed	hawkweed	plantain	
chicory	healall	purslane	
cinquefoil	henbit	pusley (Richardia	
clover (hop)	horseweed	spp.)	

cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.
PESTICIDE DISPOSAL and CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Scotts® Weed and Feed 26-0-3 F 643
 Weed Control Plus Lawn Fertilizer

Guaranteed Analysis

Total nitrogen (N)	26%
7.7% ammoniacal nitrogen	
10.1% urea nitrogen	
7.5% other water soluble nitrogen*	
0.7% water insoluble nitrogen	
Soluble potash (K ₂ O)	3%
Sulfur (S)	10%
10.0% combined sulfur (S)	

Derived from: urea, methyleneurea, potassium sulfate and ammonium sulfate.

*Contains 5.6% slowly available nitrogen from methyleneureas.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase, and we will mail you a refund check promptly.

Questions / Information



Consumer Helpline:
 1-800-543-TURF (8873)



Visit our website:
www.scotts.com

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with directions.



©2012 The Scotts Company LLC.
 World rights reserved.

Sold and Guaranteed by:
 The Scotts Company
 14111 Scottslawn Rd.
 Marysville, OH 43041

EPA Reg No. 538-270 (5M)
 Scotts Weed & Feed Weed Control Plus Lawn Fertilizer