


84401

Chlorine-free

BAQUACIL® EASY START KIT

For Plaster Pools
and Attached Water
Features and Spas

- ✓ BAQUACIL® Swimming Pool Sanitizer and Algistat (2-64oz. bottles)
- ✓ BAQUACIL® Performance Algicide (1-32oz. bottle)
- ✓ BAQUACIL® Test Strips (25 strips) with Pool Care Guide
- ✓ BAQUACIL® Chlorine Neutralizer (2-5 oz. bottles)
- ✓ BAQUACIL® Filter Cleaner (1-32 oz. bottle)
- ✓ BAQUACIL® Metal Control (2-1.25 lb. bottles)
- ✓ BAQUACIL® Water Sample Bottle
- ✓ Free Pool Care CD-ROM Inside



AS EASY AS 1-2-3

1. Balance Pool Water
2. Remove and Neutralize Chlorine
3. Add BAQUACIL Products

Swim chlorine-free in less than 24 hours.

For all pools up to 20,000 gallons



84401

Chlorine-free

BAQUACIL[®]

EASY START KIT

For Plaster Pools
and Attached Water
Features and Spas



AS EASY AS 1-2-3
1. Balance Pool Water
2. Remove and Neutralize Chlorine
3. Add BAQUACIL Products

Swim chlorine-free in less than 24 hours.

For all pools up to 20,000 gallons

**Chlorine-free
BAQUACIL[®]
SWIMMING POOL
SANITIZER AND ALGISTAT**

ACTIVE INGREDIENT:	
POLY (Oximinodimethylamino)carbamoylamine	
Nonaquidocarbonylmethoxyhexamethylene	
HYDROCHLORIDE	20%
INERT INGREDIENTS	80%
TOTAL	100%

**KEEP OUT OF REACH OF CHILDREN
WARNING**

**PRECAUTIONARY STATEMENTS. HAZARDS
TO HUMANS AND DOMESTIC ANIMALS.
WARNING**

**CAUSES SUBSTANTIAL BUT TEMPORARY EYE
INJURY. HARMFUL IF SWALLOWED.**

Do not get concentrate in eyes or on clothing. Avoid contact with skin. Avoid breathing vapor or mist. Wash thoroughly with soap and water after handling. Wear goggles or face shield when handling concentrate. Keep container closed. May cause allergic skin reactions.

FIRST AID

IF IN EYES

- Hold eyes open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**MEDICAL EMERGENCY INFORMATION CENTER
1-800-654-6811**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner other than for which it was intended. BAQUACIL[®] can be used in both residential and public swimming pools and spas.

TREATING POOL

POOLS PREVIOUSLY TREATED WITH CHLORINATING CHEMICALS
Important - BAQUACIL SWIMMING POOL SANITIZER AND ALGISTAT (BAQUACIL SANITIZER) is incompatible with chlorinating chemical products. Consult your Authorized Dealer of BAQUACIL for information on balancing your pool water. Note: If the pool is a plaster pool, add 1 1/4 lbs. of BAQUACIL METAL CONTROL per 10,000 gallons after the chlorine has been neutralized, and allow pump/filter to operate for 8-12 hours. Check with on-time test

kit for free and combined chlorine prior to the beginning of neutralization. To destroy chlorine, add BAQUACIL CHLORINE NEUTRALIZER (per label directions) (add 1 gallon of BAQUACIL CHLORINE NEUTRALIZER (27% hydrogen peroxide) per 10,000 gallons of pool water. Filter overnight. Check with on-time test kit. If free chlorine is detected, repeat neutralizer treatment.

**FRESHLY FILLED AND DECHLORINATED POOLS
START UP WITH BAQUACIL**

Note: If the pool water contains algae or is hazy, consult the problem-solving section of "The Pool Care Guide for BAQUACIL," available from your Authorized Dealer of BAQUACIL.

- Rockwash or clean pool filter. Check pool pH; adjust to 7.2-7.8 if necessary.
- Add BAQUACIL SANITIZER to give a content of 50 ppm (1/2 oz. gallon product/10,000 gallons of pool water).
- After 24 hours, check pH and BAQUACIL SANITIZER concentration with BAQUACIL TEST STRIPS and adjust if necessary.

NORMAL POOL ROUTINE

- Every week, check BAQUACIL SANITIZER concentration and pH.
- Maintain BAQUACIL SANITIZER concentration at about 50 ppm. If concentration falls 10 ppm or more, add a top-up dose. The level should not be allowed to drop below 30 ppm. For example, to raise the BAQUACIL SANITIZER level 10 ppm, add 1 pint per 10,000 gallons of water. Consult the booklet, "The Pool Care Guide for BAQUACIL," for calculation instructions.

NOTE: BAQUACIL SANITIZER AND CHLORINE ARE INCOMPATIBLE. Refer to "The Pool Care Guide for BAQUACIL" for information on how to deal with accidental chlorination.

PROBLEM-SOLVING AND SUPPLEMENTAL POOL INFORMATION

Do not use in freestanding spa/hot tubs. Do not use in unpainted plaster pools previously exposed to metals which may cause staining. (See current "Pool Care Guide" for specific information.) BAQUACIL is not compatible with many detergents and some supplemental algaecides (such as those based on copper).

If algae development, cloudiness, or pool problems occur or additional information on water quality is desired, consult "The Pool Care Guide for BAQUACIL," available from your Authorized Dealer of BAQUACIL.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contain late water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL
Keep container closed when not in use. Store in original container in an area inaccessible to children.
PROTECT FROM FREEZING. In case of a leak or spill, soak up with absorbent such as sand, earth, or sawdust, and shovel into waste container. Remove waste to chemical waste area, and dispose of in accordance with local pollution regulations. To dispose of excess product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL

Do not reuse container. To dispose of empty container, rinse thoroughly before discarding in trash.

**FOR INFORMATION ON SPILLS,
CALL 1-800-654-6811.**

EPA Reg. No. 1258-1263
Superficial use in Lot. No.
EPA EST. NO. 1258-TN-01(R)
EPA EST. NO. 68996-MQ-001(M)
EPA EST. NO. 38209-1J-01(M)

**Chlorine-free
BAQUACIL[®]
PERFORMANCE ALGICIDE**

ACTIVE INGREDIENTS:	
POLYOXYETHYLENE DIMETHYLIMIDIO	
ETHYLENE DIMETHYLIMIDIO ETHYLENE	
DICHLORIDE	60%
INERT INGREDIENTS	40%
TOTAL	100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS. HAZARDS
TO HUMANS AND DOMESTIC ANIMALS.
CAUTION**

Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**MEDICAL EMERGENCY INFORMATION CENTER
1-800-654-6811**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. BAQUACIL[®] Performance Algicide is used to control the growth of algae in swimming pools. For maximum effectiveness, pools containing heavy growth of algae should be cleaned prior to using BAQUACIL Performance Algicide.

For pools having just visible algae growth, add an initial dose of 11 to 17 fluid ounces of BAQUACIL Performance Algicide per 10,000 gallons of water and remove settled sludge. For treatment of freshly cleaned and filled pools, add 11 fluid ounces of BAQUACIL Performance Algicide per 10,000 gallons of water. Subsequent additions of 2 to 4 fluid ounces of BAQUACIL Performance Algicide per 10,000 gallons of water should be made every 5 to 7 days after initial treatment for maintenance.

BAQUACIL Performance Algicide will prevent growth of most algae throughout the season winter months when pools are not in use. For pools which have

been properly treated during the swimming season and are free of visible algae, add 8 to 10 fluid ounces of BAQUACIL Performance Algicide per 10,000 gallons of water. At the end of swimming season, pools have some visible algae growth, add 12 to 17 fluid ounces per 10,000 gallons. It is necessary that a uniform distribution of algicide throughout the water and the pool be made for maximum effectiveness. If pool is not covered, it may be necessary to repeat this treatment one or more times during the season.

BAQUACIL Performance Algicide is compatible with those chemicals normally used to treat pools and is effective at both acid and alkaline pH. BAQUACIL Performance Algicide can be used in pools treated with chlorine chemicals and may reduce the amount of those chemicals normally required. However, do not mix BAQUACIL Performance Algicide with concentrated dry or liquid chlorine products.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not contaminate water by the cleaning of equipment or disposal of waste. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the sewer treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Do not stack more than four drums high. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

PLASTIC
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

METAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**FOR INFORMATION ON SPILLS,
CALL 1-800-654-6811.**

EPA Reg. No. 1258-1272
EPA EST. NO. 4991-TN-3

BAQUACIL[®] is a registered trademark of Arch UK Bioceutics Ltd. Arch Chemicals, Inc., Smyrna, GA 30080
For more information visit our website at www.baquacil.com

CONSUMER COMMODITY

