



# FIFRA Section 24(c) Special Local Need (SLN) Label FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

## Proline® 480 SC Fungicide

EPA REG No. 264-825

EPA SLN No. WA-260007

For Suppression of Black Leg (*Leptosphaeria maculans*) in Listed Brassica Crops Grown For Seed

THIS LABEL FOR PROLINE 480 SC FUNGICIDE EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2030.

### ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of prothioconazole and degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Degradates of prothioconazole are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

**Brassica Crops Grown for Seed: Broccoli; broccoli, Chinese; broccoli raab; Brussels sprouts; cabbage; cabbage, Chinese, bok choy; cabbage, Chinese, napa; cauliflower; collards; kale; kohlrabi; mustard greens; mustard spinach; and rape greens.**

PEST SUPPRESSION	RATE PER APPLICATION fluid ounces/acre
Black Leg ( <i>Leptosphaeria maculans</i> )	5.0 – 5.7 (0.156 – 0.178 lbs ai/A)

**Apply by foliar broadcast only; ground or aerial application.**

Note: Use in rotation in resistance management programs where additional products are available.

### RECOMMENDED APPLICATION TIMINGS FOR BIENNIAL BRASSICA SEED CROPS:

Up to 2 applications in the fall may be followed with up to 3 applications in the spring/summer when using the 5.0 fl oz/acre (0.156 lbs ai/A) rate.

- **Fall Applications:** Apply Proline 480 SC in the fall just prior to the onset of cool and wet conditions that favor spread of black leg, particularly if the disease has been detected in the crop.
- **Spring Applications:** Apply Proline 480 SC in the spring when conditions become warm enough for the plants to start growing again and are moist and cool (particularly west of the Cascade Mountains) – 1 or 2 applications may be needed depending on disease pressure and environmental conditions.
- **Summer Applications:** Apply Proline 480 SC in the summer when pods begin to form and conditions are cool and moist, particularly if black leg has been detected in the crop earlier in the season and there is a risk of infection occurring on seed developing in the pods.

Maximum Proline 480 SC applications must not exceed 5 for the crop production cycle and **DO NOT** exceed the total maximum amount of 25.0 fl oz (0.781 lbs prothioconazole) of Proline 480 SC Fungicide per acre per crop production cycle.

#### **RECOMMENDED APPLICATION TIMINGS FOR ANNUAL BRASSICA SEED CROPS:**

Up to 3 applications in the spring/summer.

- **Spring Applications:** Apply Proline 480 SC after plants have emerged and conditions are moist and cool (particularly west of the Cascade Mountains) – 1 or 2 applications may be needed depending on disease pressure and environmental conditions.
- **Summer Applications:** Apply Proline 480 SC in the summer when pods begin to form and conditions are cool and moist, particularly if black leg has been detected in the crop earlier in the season and there is a risk of infection occurring on seed developing in the pods.

Maximum Proline 480 SC applications must not exceed 3 for the crop production cycle and **DO NOT** exceed the total maximum amount of 17.1 fl oz (0.534 lbs prothioconazole) of Proline 480 SC Fungicide per acre per crop year.

#### **RESTRICTIONS / PRECAUTIONS**

- **DO NOT** apply within 14 days of harvest.
- **DO NOT** windrow within 14 days of last application.
- **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- For use in accordance with this SLN label, **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply prothioconazole with mechanically pressurized handgun equipment.
- **Crops Grown for Seed Production:**
  1. All brassica seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, compostor or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
  2. No portion of the brassica seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
  3. Brassica seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
  4. Brassica seed may not be distributed for human consumption or animal feed.

**WSDA Container Disposal Guidance:** Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

#### **IMPORTANT—READ BEFORE USE**

PROLINE 480 SC FUNGICIDE, WHEN USED ON BRASSICA GROWN FOR SEED UNDER THIS SECTION 24(C) LABEL HAS NOT BEEN TESTED UNDER ALL CONDITIONS AND IN ALL TANKMIX SCENARIOS. BAYER RECOMMENDS THAT THE USER AND/OR GROWER TEST PROLINE 480 SC FUNGICIDE IN ORDER TO DETERMINE ITS COMPATIBILITY WITH THE INTENDED USE. **RISK OF CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS ARE BEYOND THE CONTROL OF BAYER, AND BAYER INTENDS THAT ALL SUCH USE IS AT USER'S AND/OR GROWER'S RISK, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW.**

THIS SPECIAL CONDITIONS AND DISCLAIMER STATEMENT IS REQUIRED BY BAYER AND IS NOT REQUIRED OR ENFORCED BY THE USEPA OR THE WSDA.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LP, 800 N. Lindbergh Blvd., St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)**

Proline is a registered trademark

3/13/2026

Page 2 of 2