

3.00" Customer Seal Area

- Use anytime spring through summer.
- For residential use on lawns & landscaped areas around trees.



Kills white grubs & mole crickets early, before they damage your lawn!

FOR OUTDOOR RESIDENTIAL USE ONLY Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, and Quee Countles of New York.

ACTIVE INGREDIENT:

Inidaclopid, 1-(I6-Chloro-3-pyridryl)methyl(N-nitro-2-inidazolidnimine, OTHER INGREDIENTS:

STOP - Read the label before use. Keep out of reach of children KEEP OUT OF REACH OF CHILDREN

CAUTION
SEE BACK PANEL FOR FIRST AID AND ADDITIONAL
PRECAUTIONARY STATEMENTS

EPA Reg. No. 42750-154-9198

EPA Est. No. 9198-OH-1M, 9198-OH (Underlined letter is first lietter used



Provides up to 5,200 sq ft of white grub controll

NET WT 15 LBS (6.82 kg)

ortify GRUB CONTROL

	FIRST AID
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice, Have person sp a glass of water if able to swallow. Co not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Pinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	Hold eye open and more slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye, Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh ar, If person is not breathing, call 911 or an ambulance, then give artificial respiration. Call a poison control center or doctor for further treatment advice.
	ncy call CHEMTREC toll free at 1-809-424-9300. Have a product container or label- ing a poison control center or doctor, or going for treatment.
Note To Physician	No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin, Causes eye irritation, Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates, Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

WHAT FORTIEY SEASON LONG GRUR CONTROL DOES

White grubs can cause extensive damage to furf by feeding on grass roots. Affected turf turns yellow, then brown, and is easily pulled up in pieces with your hands. Mole crickets can seriously damage lawns by their tunneling activity. Some species also feed on the grass roots, as well, This product prevents turl damage by killting grubs and mole crickets when they are newly hatched and before they can cause turf damage.

This product works best when applied before grubs and mole crickets hatch. The active ingredient will remain in the root zone of the turf until the eggs of grubs or mole crickets hatch. These newly hatched insects are highly susceptible to this product, and are killed before they become large enough to cause turf damage. This product also kills grubs in landscaped ornamental areas containing flowers, ground covers, shrubs and trees,

If you have questions regarding the best time to make applications in your area, you can consult your local County University Extension Office. Be sure that you describe the product you are using when seeking such advice.

HOW TO APPLY FORTIFY SEASON LONG GRUB CONTROL DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For residential use on lawns and landscaped areas.

Do not apply this product by any application method, to Linden, Basswood or other Titia species in

READ ALL DIRECTIONS FOR USE AND CONDITIONS OF SALE BEFORE USING APPLICATION TO LAWNS:

Pesis Controlled:

White grubs (including the larvae of Japanese beetle, Oriental beetle, Chaters, Asiatic garden beetle, May or June beetles, and black turigrass alaenius) and mole crickets

Using either a drop-type or rolary-type spreader, apply this product uniformly over the area being treated. Avoid the use of spreaders which may apply the product in narrow, concentrated or uneven bands, Calibrate your spreader according to the spreader manufacturer's directions to ensure the proper amount of this product is applied. Check frequently to be sure spreader is working property and spreading granules evenly,

This product will work best if the lawn is watered thoroughly or if rainfall occurs within 24 hours after application to move the active ingredient through the thatch and down into the soil. Avoid mowing lawn until watering or rainfall has occurred so that the uniformity of the granules is not disturbed.

When to Apply: This product works best when applied before grubs and mole crickets hatch. For best results, apply as follows:

White grubs: Apply once, anytime from April through mid-August.

Mole crickets: In Florida, apply once from early April through early May. Elsewhere, apply once from mid-May through mid-June.

Do not apply this product when turf areas are waterlogged. The product may not work effectively

For control of white grubs, this product should be applied at the rate of 2,87 pounds per 1,000 square feet, For mole cricket control, apply 3,8 pounds per 1000 square feet. The table below can be used as a guide for applying the proper rate of this product. However, the application rate can vary depending on the speed of application and the condition of the spreader. For best results, use a normal, steady pace when applying to obtain an accurate, even application,

Settings are approximate, The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's

Tin: $1.000 \text{ so ft} = 20 \text{ ft} \times 50 \text{ ft}$

Do not use a spreader that applies granules in narrow, concentrated, or uneven bands. Be sure the spreader works properly and spreads granules evenly

SPREADER SETTING TABLE			
SPREADERS	SETTINGS Lbs. of Product/1000 Sq. Ft		
Lb			
	2.87 Lbs.	3,80 Lbs.	
BROADCAST SPREADERS:			
Agri-Fab Model 100	3 1/4	4	
Andersons Model 4300	4	5	
Andersons Model 4500	4 1/2	5 1/4	
Ear thway Model 2000A	13	14	
Precision Model S850	6 3/4	7 3/4	
Republic EZ 540/75450	3	4	
Scatts Model SG3000	3 1/4	4 1/4	
Scotts Edge Guard	3	4	
Scotts Essygreen EG3	25	26	
Andersons Accupro 2000, SR-2000 (Cone6)	K 1/2	M	
Width 11 ft.			
DROP SPREADERS:			
Earthway Model 7350	13	14	
Precision Model FH22	9	10	
Republic Model EZ530,75350	5 3/4	6 3/4	
Scotts Model AG3000	4 1/2	5 1/4	
Scotts Model PF4	5 3/4	6 3/4	

APPLICATION TO LANDSCAPE PLANTINGS (Trees, Shrubs, Evergreens, Flowers, Ground covers)

Pests Controlled:

While grubs (including larvae of Japanese beelle, Oriental beelle, chalers, Asiatic garden beetle, May and June beetle, and black turfgrass ataenius),

Apply as a broadcast treatment to soil, mulch or other substrate. Prior to application, remove black plastic from area to be treated to allow water to penetrate into the soil, Permeable weed barriers need not be removed prior to application.

This product will work best if material is worked into the soil around plants or if a thorough watering or rainfall occurs within 24 hours after application to move the product down through

The best control of grubs will be obtained if this product is applied before grubs begin to hatch. For best results, application should occur from April through mid-August,

DO NOT apply to waterlogged areas, This product may not work effectively under these

How Much to Apply:

This product should be applied at the rate of 2.87 pounds per 1.000 square feet. This 15 pound bag will treat 5,400 square feet of landscaped area. The table above can be used as a guide for applying the proper rate of this product. However, the application rate can vary depending on the speed of application and the condition of the spreader.

NOTE: DO NOT apply this product to vegetable gardens.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal STORAGE CONDITIONS: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

CONTAINER DISPOSAL: Non-refilable container. Do not reuse or refill this container, Paper and plastic bags: Completely empty bag into application equipment. Then dispose of in sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container al once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Distributed by: The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537

FRT2GC15 FRT2GC15B



