

161
MAY 11/1.10

Produced For:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87
BONHAM, TEXAS 75418



Visit Us At: www.hi-yield.com
EPA Reg. No. 19713-235-7401 • EPA Est. No. 7401-TX-1

FIRST AID	
If In Eyes:	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person.
If Inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
If On Skin Or Clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product including health concerns, medical emergencies or pesticide incidents, call the National Pesticide Information Center at 1-800-858-7378.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

WARNING: Causes substantial but temporary eye damage. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), long-sleeved shirt, long pants, shoes, and socks. Prolonged or frequent repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

The following PPE are recommended to be worn when handling this product: Goggles or face shield, chemical-resistant gloves when using indoors or outdoors, long pants and long-sleeved shirt. Remove contaminated clothing and launder separately before re-use. Wash non-disposable clothes thoroughly with soap and water before removing.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using to let. 2) Remove clothing PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow the product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

COMPATIBILITY AND PLANT SAFETY

This product can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and Bordeaux mixture will reduce the fungicidal activity of this product. Do not apply captan in combination with, immediately before or closely following oil sprays. Do not allow oil sprays on adjacent areas to drift onto plants which have been or will shortly be treated with captan. The time factor governing the safe interval between captan and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and advisors to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, captan may cause necrotic spotting of tender immature leaves of certain varieties of Apples, Cherries, Peaches and Plums. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use captan and other spray materials at lowest recommended rates and avoid drenching trees.

APPLICATION INFORMATION

Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitator during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read "COMPATIBILITY AND PLANT SAFETY" section above.)

GENERAL USE PRECAUTIONS

Do not apply this product to seeds or seed products. Apply to flowers so as volume to provide thorough, even coverage. Do not apply this product through any type of irrigation system. Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control, but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor diseases. Apply the low rate and/or spray at longer intervals when climatic conditions least favor diseases. If you are unaware of the climatic conditions favorable for diseases claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions. IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the Eastern or Western U.S. Follow those directions for your growing area where specified.

ENTRY RESTRICTIONS

Do not allow people or pets to enter treated area until sprays have dried.

Hi-Yield Captan 50W Fungicide



• A Fungicide For Plant Disease Control.

ACTIVE INGREDIENTS:
Captan 43.0%
Related derivatives 1.1%

OTHER INGREDIENTS:
100.0%

TOTAL:
146.00%

14-Tetrahydro-4-methyl-2H-pyridin-4-one, 4-cyano-1,2-dithiobenzoate

**KEEP OUT OF REACH OF CHILDREN
WARNING / AVISO**
See Side Panel for First Aid

**Si usted no entiende la etiqueta,
busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**

WARRANTY - CONDITIONS OF SALE

OUR REPRESENTATIONS AND WARRANTIES FOR THIS PRODUCT ARE LIMITED TO THE FACTS SET FORTH ON THIS LABEL. THERE IS NO WARRANTY OR GUARANTEE OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO, THE MERCHANTABILITY OF THIS PRODUCT OR THE FITNESS OF THIS PRODUCT FOR ANY PARTICULAR PURPOSE. THE USER ASSUMES ALL LIABILITY FOR ANY AND ALL DAMAGE, LOSS OF PROFITS, OR OTHER CONSEQUENCES, INCLUDING BUT NOT LIMITED TO, THE COST OF REPLACEMENT OR REPAIR OF ANYTHING DAMAGED BY THIS PRODUCT, AND THE COST OF ANY LITIGATION, ATTORNEY'S FEES, AND COURT COSTS, ARISING OUT OF THE USE OF THIS PRODUCT. THIS WARRANTY DOES NOT APPLY TO PRODUCTS WHICH ARE NOT IDENTIFIED AS SUCH ON THIS LABEL. THE USER ASSUMES ALL LIABILITY FOR ANY AND ALL DAMAGE, LOSS OF PROFITS, OR OTHER CONSEQUENCES, INCLUDING BUT NOT LIMITED TO, THE COST OF REPLACEMENT OR REPAIR OF ANYTHING DAMAGED BY THIS PRODUCT, AND THE COST OF ANY LITIGATION, ATTORNEY'S FEES, AND COURT COSTS, ARISING OUT OF THE USE OF THIS PRODUCT.

NET CONTENT 12 OZS.

FRUIT AND NUT CROPS		
Crop	Diseases†	Rate Level (Tablespoons) per 1 Gallon of Water
Apples (Eastern U.S.)	Black rot, Alternaria, Scab, blossom end rot, Fire-blast, rust	1 to 5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small-wood trees, 1 to 2 quarts for medium-wood trees and 1 to 3 gallons for large-wood trees. Apply at 7- to 14-day intervals as needed to maintain control in orchards. Bloom, petal fall and first cover sprays. For Fire-blast, first spray also apply during the anthesis and bloom periods to a 40- to 50-day schedule. Better control of Black rot, Scab, and Alternaria may be obtained by using 1 to 1.5	1 to 1.5
	SPECIFIC DIRECTIONS: Apply at 7- to 14-day intervals in seed and fruit cover sprays.	
Note: May be applied up to day of harvest. Do not use this product in combination with or closely following or in alternation with volatile sulfur products or sulfur compounds. Varieties of Apples such as Golden, King, Red Delicious, Sturman, etc., as severe injury and defoliation may occur.		
Cherries (Western U.S.)	Berry rot, Botrytis gray mold, Blight, rust	1 to 2
	SPECIFIC DIRECTIONS: Begin at mid-blossom stage at 7- to 14-day intervals and continue to day of harvest.	
Cherries (Eastern U.S.)	Botrytis rot, Blight, Leaf spot	1 to 5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small-wood trees, 1 to 2 quarts for medium-wood trees and 1 to 3 gallons for large-wood trees. Apply in orchards. Bloom, petal fall, shoot, cover and harvest sprays. Applications of 7- to 14-day intervals may be necessary during bloom to control blossom blight. Repeat application at 7- to 14-day intervals as needed to maintain control at start of harvest. Postharvest application of 1 to 1.5 level (tablespoons per 1 gallon) of water. Apply immediately after harvest and repeat application in 7- to 14 days. May be applied up to day of harvest.	
Grades (Except CA)	Brown rot, Leaf spot, Phomopsis, Gum suppression of Black rot	1 to 1.5
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage when shoots are one-half to 1.5 inches long, when shoots are 2 to 3 inches long and when shoots are 2 to 12 inches long. Repeat petal fall, bloom, immediately after bloom and continue at 7- to 14-day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. May be applied up to day of harvest.	
Peaches	Brown rot, Scab	1 to 1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small-wood trees, 1 to 2 quarts for medium-wood trees and 1 to 3 gallons for large-wood trees. Apply in full bloom, bloom, petal fall, shoot stages and in cover and harvest sprays. When conditions are favorable, make applications at 7- to 14-day intervals during bloom to control blossom blight. Repeat application at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor disease development. Post-harvest application of 1 to 1.5 level (tablespoons per 1 gallon) of water. Apply immediately after harvest and repeat application in 7- to 14 days. May be applied up to day of harvest.	
Plums (Fresh Prunes (Eastern U.S.))	Brown rot	1 to 5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small-wood trees, 1 to 2 quarts for medium-wood trees and 1 to 3 gallons for large-wood trees. Apply in full bloom, bloom, petal fall and cover sprays. Repeat applications at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor disease development. May be applied up to day of harvest.	
Stamborras	Botrytis gray mold, Leaf spot	1 to 8
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage. Begin applications when new growth starts in the Spring and before fruit starts to form. Repeat at 7- to 14-day intervals. Under conditions favorable to Fruit rot, continue applications through harvest. Repeat immediately after each harvest. May be applied up to day of harvest.	

ORNAMENTALS

USE PRECAUTIONS: Do not apply to Lawns. Do not apply to Ornamental plants below the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of a volume of mixture per square foot area.

Crop	Diseases†	Rate per Gallon of Water
Lawns	Fusarium-off of seedlings	2 cups
	SPECIFIC DIRECTIONS: Dip seedlings before planting.	
	SPECIFIC DIRECTIONS: Apply at 7- to 14-day intervals through bloom.	
Camellias	Petal blight	1 lb/gal
	SPECIFIC DIRECTIONS: Apply to drench soil around plants beginning when flowers start to form. Repeat at 7- to 14-day intervals through bloom.	
Chrysanthemums	Botrytis flower blight, Septoria leaf spot	2 lbs/gal
	SPECIFIC DIRECTIONS: Apply at first sign of disease. Repeat at 7- to 14-day intervals.	
	Fusarium-off of seedlings	2 lbs/gal
	SPECIFIC DIRECTIONS: Dip cuttings before planting.	
Woods	Black rot, Botrytis blossom blight	2 lbs/gal
	SPECIFIC DIRECTIONS: Begin at first growth on first signs of disease. Repeat at 7- to 14-day intervals and make applications at time of harvest and heavy dews.	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal and cleaning of equipment.

PESTICIDE STORAGE: Keep pesticides in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in cool, dry place.


PESTICIDE DISPOSAL: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Reuse/Recycle container. Do not reuse or refill this container. If empty, clean to trash or offer for recycling if available. If partially filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Produced For:
VOLUNTARY PURCHASING GROUPS, INC.
 230 FM 87
 BONHAM, TEXAS 75418

Visit Us At: www.hi-yield.com

EPA Reg. No. 19713-235-7401 • EPA Est. No. 7401-TX-1



15M-10-59

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
If On Skin Or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye damage. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), long-sleeved shirt, long pants, shoes, and socks. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)


The following PPE are recommended to be worn when handling this product: Goggles or face shield, chemical-resistant gloves when using indoors or outdoors, long pants and long-sleeved shirt. Remove contaminated clothing and launder separately before re-use. Wash non-disposable gloves thoroughly with soap and water before removing.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.



Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow this product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

COMPATIBILITY AND PLANT SAFETY

This product can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of this product. Do not apply captan in combination with, immediately before or closely following oil sprays. **Do not allow oil sprays on adjacent areas to drift onto plants which have been or will shortly be treated with captan.** The time factor governing the safe interval between captan and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, captan may cause necrotic spotting of tender, immature leaves of certain varieties of Apples, Cherries, Peaches and Plums. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use captan and other spray materials at lowest recommended rates and avoid drenching trees.

APPLICATION INFORMATION

Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read "COMPATIBILITY AND PLANT SAFETY" section above.)

Do not apply this product through any type of irrigation system.

Note: There are 5 level tablespoons or 15 level teaspoons per ounce of this product.

GENERAL USE PRECAUTIONS

Do not apply this product to seeds or seed products. Apply sufficient spray volume to provide thorough, even coverage. Do not apply this product through any type of irrigation system. Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control, but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at longer intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions. **IMPORTANT:** Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the Eastern or Western U.S. Follow those directions for your growing area where specified.

ENTRY RESTRICTIONS

Do not allow people or pets to enter treated area until sprays have dried.




• A
Di

ACTIVE ING
Captan*
Related deriv
OTHER ING
TOTAL:
*N-Trichlorom
2-dicarboxim

bus
(If you do no

OUR RECOMMEN
and method of app
influencing factors
Buyer assumes all
extent consistent w
damages resulting
given herewith. Th



Hi-Yield®

Captan 50W Fungicide



in sip a glass
control center

1, preferably

th you when
ct (including
on Center at

ctive eyewear
or frequently

ical-resistant
clothing and
removing.

ove clothing/
ne outside of

of equipment

Apple
(East)

Note:
alterr
Stayr
Bluet
(Wes

Cherr
(East

Grapi
(Exce
SPEC
to 1.5
bloor
the lo
devel
disea
Peacl

Plum
Fresh
(East

of equipment

with this product
to avoid drift. Keep

on this label.

used fungicides
may lime, lime-
combination with,
plants which
captan and oil
and authorities
combinations with
should not be used
on young or tender
plants, especially
likely to occur
such as leaf spotting
on anchoring trees.

to obtain agitation
in all liquids or
on plants. (Read

even coverage;
use before or at
interval as specified,
when climatic
conditions least
favorable. For
specific use
IMPORTANT: Read
label to either the



• A Fungicide For Plant Disease Control.

ACTIVE INGREDIENTS:

Captan*	48.9%
Related derivatives	1.1%

OTHER INGREDIENTS:	50.0%
---------------------------------	-------

TOTAL:	100.0%
---------------------	--------

*N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

**KEEP OUT OF REACH
OF CHILDREN**

WARNING / AVISO
See Side Panel for First Aid

**Si usted no entiende la etiqueta,
busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**

WARRANTY - CONDITIONS OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable laws, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable laws, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

NET CONTENT 12 OZS.

Plum
Fresh
(East

Straw

USE P
the lea
the do

Azale

Came

Chrys

Rose

Do n
PES'
from
PES'
dow
CON
for ri
unus

D[®]

in

cide



FRUIT AND NUT CROPS		
Crop	Disease(s)	Rate: Level Tablespoon(s) per 1 Gallon of Water
Apples (Eastern U.S.)	Black rot (Frogeye), Botrytis blossom end rot, Primary scab	1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small/dwarf trees, 1 to 2 quarts for medium-size trees and 1 to 3 gallons for large/mature trees. Apply at 10- to 14-day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays. For Frogeye leaf spot also apply during the prebloom and bloom periods on a 10- to 14-day schedule.	
	Bitter rot, Black pox, Black rot, Botryosphaeria rot, Brooks fruit spot, Fly speck, Secondary scab, Sooty blotch	1 to 1.5
SPECIFIC DIRECTIONS: Apply at 10- to 14-day intervals in second and later cover sprays.		
Note: May be applied up to day of harvest. Do not use this product in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of Apples such as Baldwin, King, Red Delicious, Staymen, etc., as severe injury and defoliation may occur.		
Blueberries (Western U.S.)	Berry rot, Botrytis gray mold, Mummy berry	1 to 2
	SPECIFIC DIRECTIONS: Begin at mid-bloom, repeat at 7- to 10-day intervals until maturity. May be applied up to day of harvest.	
Cherries (Eastern U.S.)	Botrytis rot, Brown rot, Leaf spot	1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small/dwarf trees, 1 to 2 quarts for medium-size trees and 1 to 3 gallons for large/mature trees. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3- to 4-day intervals may be necessary during bloom to control Blossom blight. Repeat application at 7- to 20-day intervals as needed to maintain control up to start of harvest. Postharvest sprays: Leaf spot - Use 1.5 level tablespoons per 1 gallon of water. Apply immediately after harvest and repeat application in 10 to 14 days. May be applied up to day of harvest.	
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage when shoots are one-half to 1.5 inches long; when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom and continue at 10- to 14-day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible Grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible Grape varieties and during periods of weather highly favorable for disease development. May be applied up to day of harvest.	
Grapes (Except CA)	Downy mildew, Leaf spot, Phomopsis, Cane suppression of Black rot	1 to 1.5
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage when shoots are one-half to 1.5 inches long; when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom and continue at 10- to 14-day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible Grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible Grape varieties and during periods of weather highly favorable for disease development. May be applied up to day of harvest.	
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage when shoots are one-half to 1.5 inches long; when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom and continue at 10- to 14-day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible Grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible Grape varieties and during periods of weather highly favorable for disease development. May be applied up to day of harvest.	
Peaches	Brown rot, Scab	1 to 1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small/dwarf trees, 1 to 2 quarts for medium-size trees and 1 to 3 gallons for large/mature trees. Apply in full pink, bloom, petal fall, shuck stages, and in cover and preharvest sprays. When conditions are favorable, make applications at 3- to 4-day intervals during bloom to control Blossom blight. Repeat application at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown rot.	
	Coryneum blight (Peach blight, Shothole)	1.5
SPECIFIC DIRECTIONS: Apply in pink bud, full bloom, petal fall, and cover sprays as necessary and as a postharvest spray (before leaves drop). Preharvest spray may be applied up to day of harvest.		
Plums, Fresh Prunes (Eastern U.S.)	Brown rot	1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small/dwarf trees, 1 to 2 quarts for medium-size trees and 1 to 3 gallons for large/mature trees. Apply in full pink, bloom, petal fall sprays. Repeat applications at 7- to 14-day intervals as needed to maintain control. Continue applications through	



Plant

OUT OF REACH
CHILDREN
/ AVISO
Panel for First Aid

en detalle.
in it to you in detail.)

Follow directions carefully. Timing
specifically recommended and other
consistent with applicable laws.
directions given herewith. To the
consequential, special or indirect
strict accordance with directions
user.

	needed to maintain control. Continue applications through harvest if conditions favor Brown rot.	
	Coryneum blight (Peach blight, Shothole)	1.5
	SPECIFIC DIRECTIONS: Apply in pink bud, full bloom, petal fall, and cover sprays as necessary and as a postharvest spray (before leaves drop). Preharvest spray may be applied up to day of harvest.	
Plums, Fresh Prunes (Eastern U.S.)	Brown rot	1.5
	SPECIFIC DIRECTIONS: Spray 1 to 2 pints for small/dwarf trees, 1 to 2 quarts for medium-size trees and 1 to 3 gallons for large/mature trees. Apply in full pink, bloom, petal fall sprays. Repeat applications at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown rot. May be applied up to day of harvest.	
Strawberries	Botrytis (Gray mold), Leaf spot	4 to 8
	SPECIFIC DIRECTIONS: Apply sufficient spray volume to provide thorough, even coverage. Begin applications when new growth starts in the Spring and before fruit starts to form. Repeat at 7- to 14-day intervals. Under conditions favorable to Fruit rot, continue applications through harvest period, treating immediately after each picking. May be applied up to day of harvest.	

ORNAMENTALS

USE PRECAUTIONS: Do not apply to Lawns. Do not apply to Ornamental plants below/beyond the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot area.

Crop	Disease(s)	Rate per Gallon of Water
Azaleas	Damping-off of cuttings	3 tbsps.
	SPECIFIC DIRECTIONS: Dip cuttings before bedding.	
	Petal blight	1.5 tbsps.
	SPECIFIC DIRECTIONS: Apply to soil area around plants and spray flowers just before bloom. Repeat at 7- to 14-day intervals through bloom.	
Camellias	Petal blight	1 tbsps.
	SPECIFIC DIRECTIONS: Apply to drench soil around plants beginning when flowers start to open. Repeat at 7- to 10-day intervals through bloom.	
Chrysanthemums	Botrytis flower blight, Septoria leaf spot	2 tbsps.
	SPECIFIC DIRECTIONS: Apply at first sign of disease. Repeat at 7- to 10-day intervals.	
	Damping-off of cuttings	3 tbsps.
	SPECIFIC DIRECTIONS: Dip cuttings before bedding.	
Roses	Black spot, Botrytis blossom blight	2 tbsps.
	SPECIFIC DIRECTIONS: Begin at first growth or first sign of disease. Repeat at 7- to 14-day intervals and more frequently during frequent rains and heavy dews.	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal and cleaning of equipment.

PESTICIDE STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling-if available. If partially filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.