

2.5"
1.0"
31.5"
3"
Web Weight
48.50g

15

1.5"
SEAL AREA
CONCRETE, 11-1600 23054 High Yield Grub Free Zone II 15lb Bag
1.5"
Black
Black
4.5"
Arizona
Turf
Turf

33054

Grub Free Zone II

Hi-Yield®

Grub Free Zone II



Contains Imidacloprid

One Application Controls Grubs All Season Long

- Apply Before Insects Hatch To Prevent Turf Damage.
- Newly hatched insects are highly susceptible to Imidacloprid.
- Treats Up To 5,000 Sq. Ft.

FOR OUTDOOR RESIDENTIAL USE ONLY

Not for sale, sale info, distribution and use in Alaska, Hawaii, Kings and Queens Counties of New York State.

ACTIVE INGREDIENT:	
Imidacloprid, 1-(1E)-2-chloro-N-[(2E)-2-methylprop-1-en-1-yl]pyridin-4-amine	9.2%
OTHER INGREDIENTS:	90.8%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For First Aid, Precautionary Statements And Directions For Use

NET WEIGHT 15 LBS. (6.8 KG)



Hi-Yield
Grub Free Zone II
33054

SEAL AREA

1.5" SEAL AREA

Hi-Yield®

Grub Free Zone II

STOP - Read the label before use.

FIRST AID	
If Swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If On Skin or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If In Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. Call a poison control center or doctor for further treatment advice.

In the event of a chemical or physical emergency, contact 1-800-368-3000.
 North America 1-800-368-3000 or Worldwide toll-free 81-814-84-8088. Have the product container with you when calling a poison control center or doctor or going for treatment.
Read The Precaution: The specific articles is available. Treat the patient symptomatically.

APPLICATION EQUIPMENT AND INSTRUCTIONS

This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply the product evenly and uniformly over the target area (lawn) with a spreading rate of 2.87 lbs. per 1,000 sq. ft. before each application. Determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the total area to apply 2.87 lbs. per 1,000 sq. ft. and your area to treat is 6,000 sq. ft., then place 16 lbs. of product in the hopper and set the spreader setting at 3 (per table below) if using an AccuSpin™ type spreader.

A more uniform application can be made by spreading over-half of the recommended amount of this product over the target area and then applying the remaining one-half of the product at a right angle to the previous application direction.

NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working conditions of the spreader will affect the rate of application.

Spreader	Spreader Settings	
	Lbs. of product per 1,000 sq. ft.	Sp. No.
AccuSpin™ AccuSpin™ (Drop)	2.87 lbs.	3.3 lbs.
Scotts® Turf Builder® (Broadcast)	2.54	3.34
Permatron® Turf Builder® (Broadcast)	1.4	1.6

Tip: 1,000 sq. ft. = 20 ft. x 50 ft.

Do not use a spreader that applies granules in narrow concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

WHAT HI-YIELD® GRUB FREE ZONE II DOES

White grubs can cause extensive damage to lawns by feeding on grass roots. Affected lawns turn yellow, thin, brown, and finally pull up in clumps with your hands. White grubs can seriously damage lawns by their tunnelling activity. Some species also feed on the grass roots, as well. **HI-YIELD® GRUB FREE ZONE II** prevents grub damage by killing grubs and male cicadas when they are newly hatched and before they can cause further damage.

HI-YIELD® GRUB FREE ZONE II works best when applied before grubs and male cicadas hatch. The active ingredient will remain in the root zone of the lawn until the eggs of grubs or male cicadas hatch. These newly hatched insects are highly susceptible to **IMIDACLOPRID**. **DO NOT** use **HI-YIELD® GRUB FREE ZONE II** on lawns that have been damaged by grubs or male cicadas. **HI-YIELD® GRUB FREE ZONE II** also kills grubs in landscaped ornamental areas containing flowers, ground covers, shrubs and trees.

HOW TO APPLY HI-YIELD® GRUB FREE ZONE II DIRECTIONS FOR USE

If you have questions regarding the best time to make applications in your area, you can consult your local County University Extension Office. Be sure that you describe the product you are using when calling such advice.

HOW TO APPLY HI-YIELD® GRUB FREE ZONE II DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For residential use on lawns and landscaped areas.

READ ALL DIRECTIONS FOR USE AND CONDITIONS OF SALE BEFORE USING

APPLICATION TO LAWNS

Precaution: White grubs (including larvae of Japanese beetle, Oriental beetle, Chafers, Asiatic garden beetle, May or June beetle, and black turfgrass stemmors) and male cicadas.

How to Apply: Using either a drop-type or rotary-type spreader, apply **HI-YIELD® GRUB FREE ZONE II** uniformly over the area being treated. Avoid the use of sidewalks, which may apply the product in narrow, concentrated or uneven bands. Calibrate your spreader according to the spreader manufacturer's directions to assure the proper amount of **HI-YIELD® GRUB FREE ZONE II** is applied. Check frequently to be sure **HI-YIELD® GRUB FREE ZONE II** is working properly and spreading granules evenly.

After Applying: **HI-YIELD® GRUB FREE ZONE II** will work best if the lawn is watered thoroughly or if rainfall occurs within 24 hours after application to move the active ingredient through the thatch and down into the soil. Avoid mowing lawn until watering or rainfall has occurred so that the uniformity of the granules is not disturbed.

When to Apply: **HI-YIELD® GRUB FREE ZONE II** works best when applied before grubs and male cicadas hatch. For best results, apply as follows:
 White grubs: Apply area anytime from April through mid-August.
 Male cicadas: In Florida, apply area from early April through early May. Elsewhere, apply area from mid-May through mid-June.

Do not Apply HI-YIELD® GRUB FREE ZONE II when turf areas are waterlogged. The product may not work effectively under these conditions.

How Much to Apply: For control of white grubs, **HI-YIELD® GRUB FREE ZONE II** should be applied at the rate of 2.87 pounds per 1,000 square feet. For male cicada control, apply 3.0 pounds per 1,000 square feet. The table below can be used as a guide for applying the proper rate of **HI-YIELD® GRUB FREE ZONE II**. However, the application rate can vary depending on the speed of application and the condition of the spreader. For best results, use a normal, steady pace when applying to obtain an accurate even application.

APPLICATION TO LANDSCAPE PLANTINGS
 (Trees, Shrubs, Evergreens, Flowers, Ground covers)

Precaution: White grubs (including larvae of Japanese beetle, Oriental beetle, Chafers, Asiatic garden beetle, European Crane fly, May and June beetle, and black turfgrass stemmors).

How to Apply: Apply as a broadcast treatment to soil, mulch or other substrates. Prior to application, remove black plastic from area to be treated to allow water to penetrate into the soil. Permeable weed barriers need not be removed prior to application.

After Applying: **HI-YIELD® GRUB FREE ZONE II** will work best if material is worked into the soil around plants or if a thorough watering of rainfall occurs within 24 hours after application to move the product down through mulch and into the soil.

When to Apply: The best control of grubs will be obtained if **HI-YIELD® GRUB FREE ZONE II** is applied before grubs begin to hatch. For best results, application should occur from April through mid-August.

DO NOT apply to waterlogged areas. **IMIDACLOPRID** may not work effectively under these conditions.

How Much to Apply: **HI-YIELD® GRUB FREE ZONE II** should be applied at the rate of 2.87 pounds per 1,000 square feet. This 15 square foot will treat 5,000 square feet of landscaped area. The table below can be used as a guide for applying the proper rate of **HI-YIELD® GRUB FREE ZONE II**. However, the application rate can vary depending on the speed of application and the condition of the spreader.

NOTE: DO NOT apply **HI-YIELD® GRUB FREE ZONE II** to susceptible plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE CONDITIONS: Store in original container in a cool, dry place, out of the reach of children, preferably in locked storage cabinet.
CONTAINER DISPOSAL: Non-refillable containers. Do not reuse or refill this container. Paper and plastic bags: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Seller's control, it is impossible for Seller to eliminate all risks associated with the use of this product. As a result, full theory or effectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SELLER MAKES NO OTHER REPRESENTATION, EXPRESS OR IMPLIED, OF MERCHANTABILITY OF OR FITNESS FOR A PARTICULAR PURPOSE, OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Voluntary Purchasing Group, Inc. is authorized to make any representation beyond those contained herein or to modify the warranties contained herein. Seller disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, DAMAGES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER BY CONTRACT, WARRANTY, TORT NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, ON AT SELLER'S ELECTION, THE REPLACEMENT OF PRODUCT.



Manufactured by:
VPG
 Voluntary
 Purchasing
 Group, Inc.
 230 FM 87 • BONHAM, TEXAS
 Visit Us At: www.h-yield.com
 EPA Reg. No. 42750-154-7401 • EPA Est. No. 7401-TX-1
 (RV090910)

