



Grub-No-More

GRANULES



- ONE APPLICATION STOPS GRUBS ALL SEASON LONG
- USE SPRING THROUGH LATE SUMMER



ACTIVE INGREDIENT:
Imidacloprid 0.2%
OTHER INGREDIENTS: 99.8%
TOTAL 100.00%

Active Ingredient: 0.2% imidacloprid,
CAS Registry No. 138264-41-3

FOR OUTDOOR RESIDENTIAL USE ONLY
Not for sale, sale into, distribution and/or
use in Nassau, Suffolk, Kings, and Queens
counties of New York state.

See back panel for complete Precautionary
Statements and Directions for Use (including
First Aid and Storage and Disposal).

KEEP OUT OF
REACH OF CHILDREN
CAUTION

526/4-2025

NET WEIGHT: 30 POUNDS



Grub-No-More

GRANULES



Grub-No-More

GRANULES

Grub-No-More

GRANULES



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency call toll-free at 877-800-5556.
Active Ingredient: 0.2% imidacloprid, CAS Registry No. 138261-41-3.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions For Use and Conditions of Sale before using. For residential use on lawns and landscaped areas. Do not apply this product by any application method, to linden, basswood or other Tilia species in the state of Oregon.

APPLICATION TO LAWNS:

What Grub-No-More® Granules does:


White grubs can cause extensive damage to turf by feeding on grass roots. Affected turf turns yellow, then brown, and is easily pulled up in pieces with your hands. Mole crickets can seriously damage lawns by their tunneling activity. Some species also feed on the grass roots, as well. This product prevents turf damage by killing grubs and mole crickets when they are newly hatched and before they can cause turf damage. This product works best when applied before grubs and mole crickets hatch. The active ingredient will remain in the root zone of the turf until the eggs of grubs or mole crickets hatch. These newly hatched insects are highly susceptible to imidacloprid, and are killed before they become large enough to cause turf damage. This product also kills grubs in landscaped ornamental areas containing flowers, groundcovers, shrubs and trees. If you have questions regarding the best time to make applications in your area, you can consult your local County University Extension Office. Be sure that you describe the product you are using when seeking such advice.

Pests Controlled:

White grubs (including the larvae of Japanese beetle, Oriental beetle, chafers, Asiatic garden beetle, May or June beetles, and black turfglass ateneus) and mole crickets.

How to Apply:

Using either a drop-type or rotary-type spreader, apply this product uniformly over the area being treated. Avoid the use of spreaders which may apply the product in narrow, concentrated or uneven bands. Calibrate your spreader according to the spreader manufacturer's directions to ensure the proper amount of this product is applied. Check frequently to be sure spreader is working properly and spreading granules evenly.



After Applying:

This product will work best if the lawn is watered thoroughly or if rainfall occurs within 24 hours after application to move the active ingredient through the thatch and down into the soil. Avoid moving lawn until watering or rainfall has occurred so that the uniformity of the granules is not disturbed.

When to Apply:

This product works best when applied before grubs and mole crickets hatch.

For best results, apply as follows:

White grubs: Apply once, anytime from April through mid-August.
Mole crickets: In Florida, apply once from early April through early May. Elsewhere, apply once from mid-May through mid-June.

Do not apply this product when turf areas are waterlogged. The product may not work effectively under these conditions.

How Much to Apply:

For control of white grubs, this product should be applied at the rate of 2.87 pounds of product per 1,000 square feet. For mole cricket control, apply 3.8 pounds of product per 1,000 square feet. Table 1 can be used as a guide for applying the proper rate of this product. However, the application rate can vary depending on the speed of application and the condition of the spreader. For best results, use a normal, steady pace when applying to obtain an accurate, even application.

Table 1. SPREADER SETTING TABLE AND APPROXIMATE TREATED AREAS

Spreader Model	Application Rates	
	Setting to deliver 2.87 pounds of product per 1,000 sq. ft.	Setting to deliver 3.8 pounds of product per 1,000 sq. ft.
EARTHWAY ROTARY	13	15
PRECISION DROP	12	14
PRECISION ROTARY	4.7	5.9
RED DEVIL ROTARY	7	8
SEARS ROTARY	3	3.5
SCOTT'S ACCUGREEN	5.25	5.75
SCOTT'S EDGEGUARD	3.25	4
SCOTT'S PRECISIONGREEN	4	4.5
SCOTT'S SPEEDYGREEN	3.25	4
SCOTT'S TRACTOR	3	3.5
SPYKER ROTARY	4.75	5.25

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.

Tip: 1,000 sq. ft. = 20 ft. x 50 ft.

Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

APPLICATION TO LANDSCAPE PLANTINGS

(Trees, Shrubs, Evergreens, Flowers, Groundcovers)

Pests Controlled: White grubs (including larvae of Japanese beetle, Oriental beetle, chafers, Asiatic garden beetle, May and June beetle, and black turfglass ateneus).

How to Apply:

Apply as a broadcast treatment to soil, mulch or other substrate. Prior to application, remove black plastic from area to be treated to allow water to penetrate into the soil. Permeable weed barriers need not be removed prior to application.

After Applying:

This product will work best if material is worked into the soil around plants or if a thorough watering or rainfall occurs within 24 hours after application to move the product down through mulch and into the soil.

When to Apply:

The best control of grubs will be obtained if this product is applied before grubs begin to hatch. For best results, application should occur from April through mid-August.

DO NOT apply to waterlogged areas. Imidacloprid may not work effectively under these conditions.

How Much to Apply:

This product should be applied at the rate of 2.87 pounds of product per 1,000 square feet. **This 30 pound bag will treat 10,000 square feet of landscaped plantings.** However, the application rate can vary depending on the speed of application and the condition of the spreader.

NOTE: DO NOT apply this product to vegetable gardens.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE CONDITIONS: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN THIS PRODUCT UNOPENED IMMEDIATELY TO THE POINT OF PURCHASE, AND THE PURCHASE PRICE WILL BE REFUNDED IN FULL. THE TERMS OF THIS LIMITED WARRANTY AND DISCLAIMER CANNOT BE VARIED BY ANY WRITTEN OR VERBAL STATEMENTS OR AGREEMENTS AT THE POINT OF SALE OR ELSEWHERE.

MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350, SHAWNEE, KANSAS 66286
www.GordonsUSA.com

For questions or comments
call toll-free 800.884.3179
Mon.-Fri. 8:00 a.m. - 4:30 p.m. CST

or visit: GordonsUSA.com



Learn More in
English or Spanish



Obtener más información
en inglés o español

526/4-2025 APO60625
EPA REG. NO. 2217-909
EPA EST. NO. 32802-MO-1



526

#526570