

3 in

4 in



**Botanical Insecticide, Miticide, and Nematicide
REPELLANT, ANTIFEEDANT, AND INSECT GROWTH REGULATOR (IGR)**

ACTIVE INGREDIENT	% by Wt.
Azadirachtin	1.2%
OTHER INGREDIENTS	98.8%
TOTAL	100.0%

Contains 0.35 grams azadirachtin per fluid ounce.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE
WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS,
AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Botanical product for control of insects on indoor and outdoor ornamental flowers, hemp, trees, shrubs, vegetables, fruit and nut trees, and plants, including plants grown in containers, residential recirculatory, aeorponic, and hydroponic systems, interiorscapes, home and garden use.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Net Contents: EPA Reg. No. 71908-1-81268
One Pint (16 fl. oz.) EPA Est. No 72806-OK-1^c
Superscript is first letter of lot number.




PARRY AMERICA INC.
1521 N. Cooper St., Ste 350
Arlington, TX 76011



Customer Approval _____ Date: _____



**Botanical Insecticide, Miticide, and Nematicide
REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)**

- BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR ORNAMENTAL FLOWERS, HEMP, TREES, SHRUBS, VEGETABLES, FRUIT AND NUT TREES, AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, RESIDENTIAL RECIRCULATORY, AEROPONIC, AND HYDROPONIC SYSTEMS, INTERIORSAPES, HOME AND GARDEN USE.



FOR ORGANIC GARDENING

See the Directions for Use for a Complete List of Insects Controlled.

ACTIVE INGREDIENT:	% By Wt.
Azadirachtin	1.2%
OTHER INGREDIENTS	<u>98.8%</u>
TOTAL	100.00%

Contains 0.35 grams azadirachtin per fluid ounce.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PARRY AMERICA INC.
1521 N. Cooper St. Suite 350
Arlington, TX 76011

EPA Reg. No. 71908-1-81268

USA Phone Number: 972-325-1227

JOB 165549

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.
CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY
AT 1-888-478-0798.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS -

CAUTION: Harmful if absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

MODE OF ACTION:

AzaMax® controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

AzaMax® will provide control results comparable to the synthetic insecticide standards. **AzaMax®** provides broad spectrum control with very low environmental impact.

AzaMax® provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in **AzaMax®** - Azadirachtin – is a unique botanical insecticide, miticide and nematocide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin.

Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
Anti-ovipository	Insect do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
Repellent	Insects do not prefer treated plants.

PESTS CONTROLLED OR SUPPRESSED

Use **AzaMax®** against the following pests.

TARGET PEST SPECIES OF AzaMax®

HEMIPTERA AND HOMOPTERA

including but not limited to: **true bugs** including boxelder bugs, chinch bugs, lygus bugs and stink bug; **lacebugs**; **leafhoppers** including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; **mealy bugs** including apple mealy bugs, citrus mealy bugs, grape mealy bugs; **whiteflies** including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; **aphids** including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; **psyllids** including pear psyllids and **scales** including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.

(continued on next page)

TARGET PEST SPECIES OF AzaMax® (continued)

LEPIDOPTERA

including but not limited to: **moths** including European pine shoot moth, pine tip moth and Tussock moth; **leafrollers** including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; **cutworms** including black cutworm and citrus cutworm; **caterpillars and loopers** including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; **armyworms** including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; **webworms** and **leaf perforators**.

COLEOPTERA

including but not limited to: **beetles, grubs and weevils** including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girdlers.

(continued on next page)

TARGET PEST SPECIES OF AzaMax® (continued)

DIPTERA

including but not limited to: * **flies** including Caribbean fruit fly, cherry maggots, crane fly, **fungus gnat**, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; **leafminers** including citrus leafminers and serpentine leafminers.

*Not intended for use on public health pests

THYSANOPTERA

including but not limited to: **thrips** including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.

ACARINA

including but not limited to: * **mites including**, red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.

*Not intended for use on public health pests

ORTHOPTERA

including but not limited to: **crickets; grasshoppers; locusts**

HYMENOPTERA

including but not limited to: * **sawflies** including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.

*Not intended for use on public health pests

NEMATODA

Nematodes (suppression)

FOR USE ON ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers including but not limited to:

Actinopterus, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hедера, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia

*** Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.**

(continued on next page)

**FOR USE ON ORNAMENTALS AND
LANDSCAPE PLANTINGS** *(continued)*

Ornamental Trees and Shrubs including but not limited to:

Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an **AzaMax®** application.

Applications of **AzaMax®** will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

**FOR USE ON GARDEN VEGETABLES, HERBS,
SPICES, FRUITS AND BERRIES**

Leafy Vegetables including but not limited to:	Broccoli , Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce , Spinach
Root Vegetables, including but not limited to:	Beet, Carrot , Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables including but not limited to:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables including but not limited to:	Cucumber , Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables including but not limited to:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables including but not limited to:	Garlic, Onion , Shallot
Berries including but not limited to:	Blackberry, Blueberry, Raspberry, Strawberry , others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry <i>(continued on next page)</i>

**FOR USE ON GARDEN VEGETABLES, HERBS,
SPICES, FRUITS AND BERRIES** *(continued)*

Herbs and Spices including but not limited to:	Chive, Dill, Fennel, Mustard , Sage, Sweet bay, others include: Anise, Balm, Basil , Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop , Marjoram, Marigold , Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme , Wintergreen, Woodruff, Wormwood
Nut Trees including but not limited to:	Almond , Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Pome Fruits including but not limited to:	Apple , Quince, or Pear <i>(Comice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)</i>
Stone Fruits including but not limited to:	Apricot, Cherry , Nectarine, Peach, Plum
Citrus Fruits including but not limited to:	Grapefruit , Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine , Satsuma (orange mandarin), Tangerine
Other Crops	Hemp

AzaMax® has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying **AzaMax®** up to the time of harvest.

SPRAY PREPARATION

AzaMax® is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion and which separates upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

AzaMax® is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

A jar test can quickly determine physical compatibility. The process of conducting jar test is given below:

STEP 1: Add one pint of water to a glass jar with a lid. (Use the same water source that will go in the tank.).

STEP 2: Check spray water pH and adjust if necessary. Often, the pesticide label will give the optimal pH range for best results.

STEP 3: Add the pesticides to the jar you plan to use one at a time, and shake vigorously after each addition.

STEP 4: After all products have been added, shake again, let the solution stand for 15 minutes and then shake one last time and observe the results.

Results: Jar is cool to the touch, and mixture is smooth. Then it is compatible mixture.

If a broader spectrum of control is required tank-mix **AzaMax®** with insecticides or miticides. If a rapid knock-down of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with **AzaMax®**.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. *Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.*

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. **AzaMax®** is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use **AzaMax®** with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Apply **AzaMax®** with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers, hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

APPLICATION SCHEDULE

For the most effective control, apply the product when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. For best results, spray in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with your state agricultural experiment station or local extension horticulturist/arborist for information on tactics and windows of application.

APPLICATION METHODS

- For the most effective control, spray the product as soon as possible after pests appear and are in immature stages.
- Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels use higher label rates and increase the spray frequency.
- For best results, spray in the morning or evening hours.
- Repeat spraying if rain occurs within two to three hours of spraying.

Apply **AzaMax®** as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl. oz. per 1,000 sq. ft. per application.

Apply **AzaMax®** alone to food/garden crops on the day of harvest.

Dilute **AzaMax®** with water at a rate of 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

FOLIAR APPLICATION

USE	SPRAY CONCENTRATION%	Amounts of AzaMax®	
		Fluid Ounces (Tbs.) Per Quart	Fluid Ounces (Tbs.) Per Gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals & garden plants, hemp, and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.08 - 0.25 fl. oz. (1/6 - 1/2 Tbs.)	0.32 - 1.0 fl. oz. (2/3 - 2.0 Tbs.)
	Medium rate ranges of 0.75 - 1.25% vol/vol:	0.25 - 0.40 fl. oz. (1/2 - 5/6 Tbs.)	1.0 - 1.6 fl. oz. (2.0 Tbs. - 3 1/5 Tbs.)
	Upper rate ranges of 1.25 - 1.70% vol/vol:	0.40 - 0.50 fl. oz. (5/6 - 1.0 Tbs.)	1.6 - 2.0 fl. oz. (3 1/5 - 4 Tbs.)

DRENCH APPLICATION

Use **AzaMax®** as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of **AzaMax®**. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute **AzaMax®** with water for concentrations of 0.4 to 0.8% vol/vol. See use rate table below. Add the required amount of **AzaMax®** to a clean bucket with at least one-half of the water to be drenched. Agitate the mixture thoroughly and then fill with the remaining water and continue agitation until the product is thoroughly dispersed. Drench the soil in the pot with one (1) pint of finished product dilution per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% vol/vol spray concentration. For leafminers and other difficult to control pests, use the 0.8% vol/vol spray concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure make applications every 5 to 6 days. Additional applications of **AzaMax®** may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of AzaMax®			Application Interval
	0.4%	0.6%	0.8%	
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 - 14 days
1 gallon	0.5 fl.oz.	0.8 fl.oz.	1.0 fl.oz.	10 - 14 days
5 gallons	2.7 fl.oz.	4.0 fl.oz.	5.0 fl.oz.	10 - 14 days
10 gallons	5.4 fl.oz.	8.0 fl.oz.	10.0 fl.oz.	10 - 14 days

**RESIDENTIAL RECIRCULATORY, AEROPONIC,
AND HYDROPONIC APPLICATION**

Use **AzaMax®** in recirculatory, aeroponic, or hydroponic systems for the control of foliar pests, soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or soil borne thrips for interiorscapes, hydroponic, aeroponic and container plants.

Dilute **AzaMax®** with water for concentrations of 0.1% to 0.8% volume/volume in a recirculatory or in a hydroponic liquid system. See use rate table below. Agitate the mixture thoroughly until the product is thoroughly dispersed.

For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure applications make applications every 5 to 7 days. Additional applications of **AzaMax®** may be required with increased and prolonged pest infestation.

**DILUTION TABLE FOR RECIRCULATORY, AEROPONIC,
AND HYDROPONIC APPLICATIONS**

Gallons of Water	Amount of AzaMax®				Application Interval
	0.1%	0.2%	0.4%	0.6%	
1 gallon	¼ Tbs.	½ Tbs.	1 Tbs.	1.5 Tbs.	2.0 Tbs.
1 gallon	0.14 fl. oz.	0.25 fl. oz.	0.5 fl. oz.	0.8 fl. oz.	1.0 fl. oz.
5 gallons	0.7 fl. oz.	1.3 fl. oz.	2.5 fl. oz.	4.0 fl. oz.	5.0 fl. oz.
10 gallons	1.4 fl. oz.	2.6 fl. oz.	5.0 fl. oz.	8.0 fl. oz.	10.0 fl. oz.

19

Preventive applications as a recirculatory system application may be warranted for certain pests.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers in a dry, cool, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container $\frac{1}{4}$ full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or dispose of in trash or in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

NOTICE ON CONDITIONS OF SALE

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of PARRY. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

AzaMax® is a registered trademark of Parry America Inc

▲ Peel Here To Open

FIRST AID

If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT
1-888-478-0798.**



PARRY AMERICA INC.

1521 N. Cooper St. Suite 350

Arlington, TX 76011

USA Phone Number: 972-325-1227

EPA Reg. No. 71908-1-81268
Rev. 5/30/18

PF-71121-4

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.




Dieline does not print.

 **Cimarron Label**

Experts in Extended Text Labeling

4201 North Westport Ave. • Sioux Falls, SD 57107

Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
3/13/20	165549	Parry America
LABEL SIZE	BOOKLET SIZE	
4" x 3"	3.75" x 2.25"	
LABEL COLORS	BOOKLET OUTSIDE COLORS	BOOKLET INSIDE COLORS
 BLK	 BLK	 BLK
PATTERN VARNISH: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

Form: CS 006B - 3/29/2017

ARTWORK IS APPROVED

REVISED PROOF NEEDED

**WE CANNOT PROCESS
THIS ORDER WITHOUT AN
AUTHORIZED SIGNATURE**

Signed _____ Date _____