TRUSTED SINCE 1926 STUMP-OUT® Stump & Vine Killer

Kill stumps and prevent sprouting

Controls woody plants and vines

ACTIVE INGREDIENT:

FPA Fst No. 4-NY-1

Triclopyr: (3,5, 6-tricloro-2-pyridinyloxyacetic acid), as the triethylamine salt 8.8% INERT INGREDIENTS:

TOTAL

Contains 6.36% triclopyr acid-0.6 lb./gal. EPA Reg No. 53883-216-4

For outdoor residential use only.

Si usted no entiende la etiqueta, busque a alquien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Keep Out Of Reach Of Children CAUTION See back panel for additional Net Contents 8 FL.OZ. (.23 liter) 51040416



STUMP-OUT STUMP & VINE KILLER

FOR THE CONTROL OF WOODY PLANTS, VINES, AND BROADLEAF WEEDS AROUND HOMES, CABINS, FENCES, WALKWAYS, AND OTHER NON-CROP AREAS.

FOR OUTDOOR RESIDENTIAL USE ONLY.

Store and transport in an upright position.

EPA Est. No. 4-NY-1 EPA Reg. No. 53883-216-4

Buyers Guarantee Limited to Label Claims.

©Bonide Products, Inc. All Rights Reserved

Distributed by Bonide Products, Inc. 6301 Sutliff Road Oriskany, NY 13424





DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

This product provides control of undesirable woody vines, plants, and broadleaf weeds around homes, cabins, fences, walkways, and other non-crop areas.

APPLICATION INSTRUCTIONS

To apply as a Stump Treatment: Apply this product undiluted to kill stumps and prevent sprouting. Using the applicator top, completely cover the freshly cut surface.

To use for Vine Control: For best results when treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, cut the vine first and treat as directed above for stump treatment. Do not let product contact desirable plants.

To apply as a Foliar Spray: In a hose-end or tank sprayer, mix 8 to 16 tablespoons (4 to 8 fl. oz.) in enough water to make one gallon of spray mixture. Apply at the rate of one gallon spray mixture per 500 sq. ft. Coverage will depend on the amount of foliage to be treated. Apply as a thorough foliage spray to wet all leaves using a coarse spray pattern with low pressure.

Use the higher rate for mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine, or winged elm), or during drought conditions. Using low rates may result in resprouting of weeds in a year following treatment.

Make applications when brush and weeds are actively growing with full leaf coverage. In 1 to 6 weeks, symptoms such as wilting will occur. Make applications when the air is calm. Effectiveness may be reduced if rainfall occurs with 24 hours of application.

USE RESTRICTIONS

- Do not use on or around edible food or food crops.
- Avoid contact with eyes or clothing during and after application.
- Keep people and pets out of the area during application.
- Measuring utensils and spray equipment should not be reused for purposes other than berbicide application.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- . Do not enter or allow others to enter the treated area until sprays have dried.

SPRAY DRIFT AND NON-TARGET PLANT PRECAUTIONS

- Do not spray when the wind is blowing toward desirable plants. Do not apply this product directly to, or allow to drift to.
- grapes, vegetable crops, flowers, ornamentals, or desirable foliage. Applications should be made when wind is low and there is little chance of spray drift.
- Do not apply with nozzles that produce a fine-droplet spray.

WOODY PLANTS, VINES AND BROADLEAF WEEDS CONTROLLED

These include:

Arkansas rose

Alder

WOODY PLANTS AND VINES California rose

Flm

Hazel

Arrowwood	Ceanothus	Honeysuckle
Ash	Cherry	Hornbeam
Aspen	Chinquapin	Kudzu
Beech	Choke cherry	Locust
Birch	Cottonwood	Madrone
Blackberry	Crataegus (hawthorn)	Maples
Blackgum	Douglas fir	Mesquite
Box elder	Dogwood	Mimosa
Brazilian pepper	Elderberry	Mulberry

Cascara

WOODY PLANTS AND VINES Cont.

Oaks Sassafras Trumpet creeper
Persimmon Scotch broom Tulip poplar
Pine Sumac Virginia creeper
Poison ivy Sweetbay magnolia
Poison oak Sweetdum Willow

Polson Oak Sweegum Willow
Poplar Sycamore Winged elm
Raspberry Tanoak Wild grape
Salmonberry Thimbleberry Wild rose

PERENNIAL BROADLEAF WEEDS CONTROLLED INCLUDE:

Rindweed Field hindweed Smartweed Burdock Field horsetail rush Spurge Canada thistle Henhit Tansy ragwort Chicory Lambsquarter Vetch Oxalis Wild lettuce Clover Curly dock Plantain Wild violet

Dandelion Ragweed

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep out of the reach of children. Store in a dry, cool area. Avoid contamination of food and foodstuffs.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain.

gently with water for 15-20 minutes.
Remove contact lenses, if present,
after the first 5 minutes, then
continue rinsing eye.
Call a poison control center or doctor
for treatment advice.

Call a poison control center or

. Hold eve open and rinse slowly and

FIRST AID

IF IN FYES:

	physician immediately for treatment advice.
	 Have person sip a glass of water if able to swallow.
	 Do not induce vomiting unless told
	to do so by the poison control center or physician.
	Do not give anything by mouth to an
	unconscious person.

or physician.

• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

Note to Physician - Probable mucosal damage may contraindicate the use of qastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if swallowed, Causes moderate eve

irritation. Avoid contact with eves or clothing. Wash thoroughly with soan and water after handling, and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, chemical resistant gloves made of any waterproof material, and shoes and socks

when applying this product. User Safety Recommendations Users should:

 Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately after handling this

product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm

weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or

wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable.

particularly where the water table is shallow, may result in groundwater contamination.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Bonide Products, Inc. or the Seller. To the extent allowed by applicable law all such risks shall be assumed by the Buyer.

Bonile Products, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

BONIDE PRODUCTS. INC. NEITHER MAKES NOR

INTENDS. NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH INCONSISTENT WITH THE LABEL DIRECTIONS. WARNING, OR CAUTIONS. TO THE EXTENT ALLOWED BY APPLICABLE LAW. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT

MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT ALLOWED BY APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BONIDE PRODUCTS, INC. AND THE SELLER OFFER THIS PRODUCT, AND

THE BUYER AND USER ACCEPT IT, SUBJECT TO THE FOREGOING CONDITIONS OF SALE AND WARRANTY.

OR OTHERWISE, SHALL BE LIMITED, AT THE