


A-Vert Plus Lawn Food 10912

ferti·lome™



A-Vert Plus Lawn Food

A-Vert Plus Lawn Food

A-Vert Plus Lawn Food

18-0-12

• Covers Up To 2,000 Sq. Ft.

• A Fertilizer Containing Gallery® And Team™, Pre-emergence Herbicides For Control Of Annual Grasses And Broadleaf Weeds In Established Home Lawns.

ACTIVE INGREDIENTS:	
Isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
Trifluralin: α,α,α-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine	0.27%
Benfluralin: N-butyl-N-ethyl-α,α,α-trifluoro-2,6-dinitro-p-toluidine	0.53%
OTHER INGREDIENTS:	98.91%
TOTAL:	100.00%
Contains 0.1308 pounds of active ingredient per 12 pound bag.	
*TRADEMARK of Dow AgroSciences LLC	

KEEP OUT OF REACH OF CHILDREN

CAUTION

Refer To Back Panel For Precautionary Statements And Directions For Use Including Storage And Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened.

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs or Clothing.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

10912-0814-TP



Covers Up To 2,000 Sq. Ft.

Net Weight 12 LBS. (5.44 KG)

A-Vert Plus Lawn Food

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing dust.

USE PRECAUTIONS

Wash hands before eating, drinking, chewing gum, using tobacco, or using a toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If Inhaled: • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Move person to a place of fresh air if possible. • Do not give anything by mouth to an unconscious person.

If Swallowed: • Call a poison control center or doctor for treatment advice. • Call a poison control center or doctor for further treatment advice.

If on Skin: • If on face, use plenty of water to flush. • If on hands, use plenty of water to wash.

Have the product container or label with you when calling a poison control center or doctor, or going to hospital, in the event of an accidental or chemical emergency contact. 1-800-368-5768 North America 1-800-252-3024 or Worldwide toll-free 1-877-834-0033

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater aquatic life and aquatic invertebrates including shrimp and daphnia. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or drift.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Pesticide storage bins in unopened containers. Protect stored product from moisture. In case of spill or leak, contain material and dispose in waste container. Recycle Non-Hazardous container. Do not reuse or refill this container. Offer for recycling if available.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying **feri-lome® A-VERT PLUS LAWN FOOD**. Application Restrictions: Do not apply **feri-lome® A-VERT PLUS LAWN FOOD** in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Early Application: Do not allow people or pets to enter the treated area until dusts have settled. If watering is required after the application, do not enter or allow others to enter the treated area (except those involved in the watering) until the watering is complete and the surface is dry. Do not use as a top-dressing for lawn or other ornamental use, or for commercial use, production, or for research purposes.

HOW THIS PRODUCT WORKS

feri-lome® A-VERT PLUS LAWN FOOD is a granular herbicide containing **Gallip®** and **Team®**, pre-emergence herbicides for application to established lawns for control of certain broadleaf weeds and annual grasses. **feri-lome® A-VERT PLUS LAWN FOOD** also feeds the lawn with a blend of fertilizers to provide slow growing nutrient release to your lawn.

feri-lome® A-VERT PLUS LAWN FOOD will not control existing weeds. Existing weeds should be controlled with a herbicide applied post-emergence herbicide either before or after application of **feri-lome® A-VERT PLUS LAWN FOOD**.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

THIS PRODUCT IS RECOMMENDED FOR USE ON THE FOLLOWING ESTABLISHED TURFGRASS SPECIES:

- Do not apply to hydroponic or hydroponically-grown plants.
- Do not apply to lawns or hydroponically-grown plants of Bermuda grass, St. Augustine grass, Centipede grass or Zoysia grass until well established.
- Do not apply to lawns or hydroponically-grown plants of any other species unless specifically labeled for use on that species.
- Do not make more than 2 applications of **feri-lome® A-VERT PLUS LAWN FOOD** per year.
- Apply **feri-lome® A-VERT PLUS LAWN FOOD** only to established lawns. Apply to newly seeded lawns (including overseed lawns) only after seedlings are established and well established after 4 weeks.
- Repeat: Do not reseed treated areas for at least 8 weeks following application of **feri-lome® A-VERT PLUS LAWN FOOD**.
- Do not apply to non-lawn areas such as flower beds or gardens. Take a precaution to prevent inadvertent application to non-lawn areas.
- Do not apply to lawns in areas where **feri-lome® A-VERT PLUS LAWN FOOD** cannot be applied and may require additional applications.
- **feri-lome® A-VERT PLUS LAWN FOOD** may apply to lawns that are weakened due to severe winter weather or is not well established.
- Do not apply to lawns or hydroponically-grown plants of any other species unless specifically labeled for use on that species.
- Do not apply in the spring to Bermuda grasses that have been reseeded with winter annual grasses since timing may occur.
- Do not reseed or reseed during application in such a way as to apply to areas of labeled areas, but apply to any area.
- Apply **feri-lome® A-VERT PLUS LAWN FOOD** only to the grass species listed on this label. The user assumes responsibility for any plant damage or other liability resulting from use of **feri-lome® A-VERT PLUS LAWN FOOD** on any other species not recommended on this label.
- Maximum Single Application Rate: 7.2 lbs. of **feri-lome® A-VERT PLUS LAWN FOOD** per 1,000 sq. ft.
- Maximum Weekly Rate: 14.4 lbs. of **feri-lome® A-VERT PLUS LAWN FOOD** per 1,000 sq. ft.
- Minimum Re-treatment Interval: Cool season turf - 56 to 70 days, warm season turf - 70 to 84 days.

ESTABLISHED COOL SEASON GRASSES			ESTABLISHED WARM SEASON GRASSES		
Bermudagrass, Colonial Bermudagrass, Creeping Fescue, Kentucky Fescue, Chewings	Fescue, Creeping Red Fescue, Sheep Fescue, Tall Ryegrass, Perennial Ryegrass		Bermudagrass, Bermudagrass, Centipede grass, Fescue, Tall Fescue	St. Augustinegrass, Zoysia grass	

WEEDS CONTROLLED

WEEDS CONTROLLED BY A REPEAT APPLICATION 4 WEEKS AFTER THE FIRST APPLICATION WHEN APPLIED PRIOR TO WEED GERMINATION					
Annual Bluegrass, Barnyardgrass, Bureweed, Lawn Bittersweet, Carpetweed	Clover, Wild Chickweed, Common Chickweed, Mouseear (Large Crabgrass)	(Smooth Crabgrass), Field Sandspur (Field Sandspur), Footcandle, Furbals	(Bottlebrush), (Bottlebrush), (Bottlebrush), (Bottlebrush), (Bottlebrush)	(Bottlebrush), (Bottlebrush), (Bottlebrush), (Bottlebrush), (Bottlebrush)	Mallow, Little Panicle, Perennial Ryegrass, Pinpointweed, Speedwell, Thymelae

WEEDS CONTROLLED BY A REPEAT APPLICATION 8 WEEKS AFTER THE FIRST APPLICATION WHEN APPLIED PRIOR TO WEED GERMINATION					
Clover, egg, Clover, White	Dandelion, Groundsel, common	Medic, Black Plantain, Broadleaf	Pantain, Bunch, Pantain, slender		

HOW TO APPLY

Apply **feri-lome® A-VERT PLUS LAWN FOOD** using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid edge or overlap as poor weed control or turfgrass injury may occur. Many cultures application may be achieved by spreading half the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splitboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate **feri-lome® A-VERT PLUS LAWN FOOD**.

APPLICATION EQUIPMENT AND INSTRUCTIONS

SPREADER	Spreader Settings	
	Lbs. of product per 1,000 sq. ft.	Rate (lb./1000 sq. ft.)
Scotts®/Husqvarna Accugrip® (Drop)	5.75 LBS.	7.2 LBS.
Scotts®/Husqvarna Spray-on Green® (Broadcast)	5-1/2	7-1/2
feri-lome® A-VERT PLUS LAWN FOOD (Broadcast)	18	22

NOTE: The granule settings listed above are approximate. The spread of which you make and the working conditions of the spreader will affect the rate of application.

HOW MUCH TO APPLY

Apply **feri-lome® A-VERT PLUS LAWN FOOD** in cool season lawns at a rate of 5.75 lbs. per 1,000 sq. ft. and to warm season lawns at a rate of 7.2 lbs. per 1,000 sq. ft.

WHEN TO APPLY

feri-lome® A-VERT PLUS LAWN FOOD can be applied at any time of the spring or fall. The type of weeds controlled will vary depending upon time of application as other weed control products may be needed to provide a total annual weed control program.

Early spring (before Forsythia blooms): If applied in early spring, **feri-lome® A-VERT PLUS LAWN FOOD** will prevent Crabgrass and other annual grasses (including *Poa annua*) and broadleaf weeds, such as Dandelion, Clover, Plantain, and Spurge, that normally germinate in the spring and summer. When used in this manner, **feri-lome® A-VERT PLUS LAWN FOOD** may be substituted for a multi-step weed and feed program. For your area, consult the local extension service for optimum timing for control of crabgrass.

Mid-late spring (after Forsythia blooms, before Memorial Day): **feri-lome® A-VERT PLUS LAWN FOOD** will prevent annual grasses (including *Poa annua*) and broadleaf weeds, such as Dandelion, Clover, Plantain, and Spurge, that have yet to germinate in the spring and summer, ensuring season-long weed control.

Early fall (before average night temperatures fall below 60°F): **feri-lome® A-VERT PLUS LAWN FOOD** will prevent establishment of winter annual grasses (including *Poa annua*) and broadleaf weeds, such as Chickweed, Henbit, and Speedwell, that normally germinate in the fall and winter and broadleaf weeds that germinate the following spring season. A supplemental application of a Crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control. **Later fall:** **feri-lome® A-VERT PLUS LAWN FOOD** will prevent establishment of broadleaf weeds, such as Chickweed, Henbit, and Speedwell, that may germinate over winter and broadleaf weeds that germinate the following spring season. A supplemental application of a Crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

TERMS AND CONDITIONS OF USE

If none of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks of use.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from **feri-lome® A-VERT PLUS LAWN FOOD** (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of **feri-lome® A-VERT PLUS LAWN FOOD** unless Seller is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

18-0-12 GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
3-5% Ammoniacal Nitrogen	
14.7% Urea Nitrogen	
Soluble Phosph (P ₂ O ₅)	12%
Boron (B)	0.05%
Copper (Cu)	0.05%
Iron (Fe)	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn)	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc (Zn)	
Derived From Urea, Ammoniacal Nitrogen, Methyl Dithionyl Sulfate, Copper Sulfate, Ferric Sulfate, Manganese Sulfate, Zinc Sulfate	



Manufactured by: **VPG** Voluntary Purchasing Groups, Inc.
 230 FM 87 • BONHAM, TEXAS 75418
 EPA Reg. No. 62719-192-7401 • EPA Est. No. 7401-TX-1
 Visit Us At: www.fertilome.com
 Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at: <http://aapco.org/metals.htm>