

AMINOPYRALID	GROUP	4	HERBICIDE
METSULFURON-METHYL	GROUP	2	HERBICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label

For Distribution and Use Only in the State of Oregon

Opensight®

EPA Reg. No. 62719-597

EPA SLN No. OR-250013

For Control of Herbaceous Broadleaf Weeds and Woody Plants in Forest Plantations

This label is valid until December 31, 2030, or until otherwise amended, withdrawn, canceled, or suspended.

Active Ingredients:

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro-	62.13%
Metsulfuron methyl (Methyl 2-[[[(4-methoxy-6- methyl-1,3,5- triazin-2-yl)-amino]carbonyl] amino]sulfonyl]benzoate)	9.45%
Other Ingredients	28.42%
Total	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the label affixed to the main container of Opensight must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the main container of Opensight.
- Read this SLN label and the label affixed to the main container of Opensight before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Forest Plantations

Opensight may be applied to forest plantations as an aerial or ground broadcast application for site preparation, or as a spot application or directed spray application for conifer release (see Application Methods section) to control herbaceous broadleaf weeds and woody plants. Opensight may be applied alone or in tank-mix combinations with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated, and (2) mixing is not prohibited by the label of the registered tank mixed products, and (3) that the tank-mix combination is physically compatible (see Tank Mix Compatibility Testing on the label affixed to the main container of Opensight). When tank mixing, use only in accordance with the restrictions, precautions, and limitations on the respective product labels.

Unless otherwise directed, applications of Opensight must include either a crop oil concentrate or a nonionic surfactant. Refer to the label affixed to the main container of Opensight for mixing instructions and spray adjuvant information.

Application Methods

For best control from broadcast and directed spray applications of Opensight, use a spray volume that will provide thorough plant coverage. The recommended spray volumes are 10 to 30 gallons per acre by air or 10 to 100 gallons per acre by ground. When making an aerial broadcast application, do not apply less than 2 gallons per acre total spray volume.

Forest Site Preparation Directions (aerial or ground broadcast)

Use up to 3.3 oz per acre of Opensight and apply in a total spray volume of 10 to 30 gallons per acre by air or 10 to 100 gallons per acre by ground. Tank mixtures with other herbicides registered for forest use, such as Accord XRT II, may improve control of woody brush and herbaceous plants. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture must be followed; follow the most restrictive label directions including the longest recommended waiting period before planting. Refer to the label affixed to the main container of Opensight for additional directions, restrictions, and precautions regarding tank mixing.

Note: Conifers planted sooner than 2 months (west of the Cascade Mountains) and 4 months (east of the Cascade Mountains) after treatment with Opensight may be injured.

Directed Spray Applications for Conifer Release (including spot application)

NOTE: To prevent injury to or loss of desirable trees, including conifers, avoid spray contact with the foliage, especially small trees and their terminal buds, and avoid application in the root zone especially when making spot treatments.

Use Opensight to release conifers from competing invasive and broadleaf weeds such as thistles, mareetail, bracken fern, wild carrot, mustards, knapweeds, tansy ragwort, and sensitive woody plants such as blackberry, buckbrush, snowberry, locust, redbud, and wild roses. The spray mixture should be directed onto foliage of competitive plants using knapsack or backpack sprayers with flat fan nozzles or equivalent any time after plants have reached full leaf size, but before autumn coloration. The majority of treated hardwoods should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct spray away from contact with conifer foliage, particularly foliage of desirable conifers.

Spot Application for site preparation or conifer release: Spot applications may be made at rates equivalent to the broadcast-applied rates of Opensight at 2 to 3.3 oz. per acre per year depending on the target species. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage. Repeat treatments may be made, but the total amount of Opensight applied must not exceed 3.3 oz per acre per year. To prevent misapplication, spot treatments should be applied using calibrated application systems. Apply the spray mixture using a calibrated boom, boomless nozzle and hand-held or backpack sprayers.

Spot treatments may be applied at an equivalent broadcast rate of up to 6.6 oz of Opensight per acre per year; however, not more than 50% of an acre may be treated at that rate.

Restrictions

- For use under this SLN label, do not apply this product through any type of irrigation system.
- Read and follow all directions, restrictions, and precautions regarding composting and forage and manure management on the label affixed to the main container of Opensight.
- Do not apply more than a total of 3.3 oz of Opensight per acre per year as a result of broadcast, spot, and/or repeat applications.
- **BUFFER REQUIREMENTS**
 - This product must be applied in a manner consistent with Oregon Senate Bill (SB) 1602 of 2020 for helicopter pesticide applications and the Oregon Forest Practices Act for all methods of pesticide applications, including buffer requirements. SB 1602 requires a 300 foot buffer for inhabited dwellings or schools for helicopter pesticide applications. The landowner or school board may request the pesticide application forego all but the inner 60 feet of the SB 1602 300 foot buffer (ORS 527.672). For more information on buffer requirements under SB 1602 and the Oregon Forest Practices Act, visit <https://www.oregon.gov/odf/working/Documents/chemicals-and-other-petroleum-products.pdf>
 - In addition to the buffer requirements of SB 1602 of 2020 and the Oregon Forest Practices Act, the following label restrictions must be observed:
 - **Avoiding Injury to Non-Target Plants:** Do not aurally apply Opensight within 50 feet of a border downwind (in the direction of wind movement), or allow spray drift to come in contact with, any broadleaf crop or other desirable broadleaf plants. For more information, refer to the Avoiding Injury to Non-Target Plants section on the label affixed to the main container of Opensight.
 - If multiple buffers apply, use the greatest buffer distance

24(c) Registrant:

Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

®™ Trademarks of Corteva Agriscience and its affiliated companies

R372-037

Accepted: 09/05/2025

Replaces: R372-028