

OUR BEST  
ANNUAL  
PROGRAM

538-270\_Scotts Step 2 Weed Control Plus Lawn Fertilizer 28-0-6\_20180111\_121\_538\_.pdf

Scotts®

# 4STEP® PROGRAM



Improves Lawn's Ability to  
Absorb Water & Nutrients

Covers 15,000 SQ FT

KEEP OUT OF REACH OF CHILDREN  
CAUTION

See back for additional precautionary statements.

ACTIVE INGREDIENTS:

2, 4-DP ..... 1.10%

Mesoprop-py ..... 0.50%

OTHER INGREDIENTS: ..... 88.39%

Total 100.00%

\*0.010 lb 2,4-D acid equivalent per pound of product

\*0.005 lb MCPA acid equivalent per pound of product

28-0-6

NET WT. 43.92 lb (19.89 kg)

Available only at authorized Scotts

**LawnPro®**  
Dealers

# STEP®2

## Weed Control plus Lawn Fertilizer

DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM,  
DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS.



Apply in Late Spring:

APR | MAY | JUN



# Now it's Easy for You to Have a Better Lawn



### Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed.

### Simple and easy

No more guesswork. You simply apply the right product – one STEP® at a time.

### Guaranteed results

Each STEP® product is designed to the specific season to help give you a greener lawn.

### Florida Annual Fertilizer Applications

On Centipede lawns, do not exceed rate of 2 lbs of Nitrogen per 1,000 sq. ft. in the north and 3 lbs of Nitrogen per 1,000 sq. ft. in the central and south. On Bahia lawns, do not exceed rate of 3 lbs of Nitrogen per 1,000 sq. ft. in the north.

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

## SPREADER SETTINGS

**Spreading Tips:** Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Covers 15,000 sq. ft. This package applies 2.93 lbs of product per 1,000 sq. ft.

Scotts Broadcast / Rotary Spreaders	Spreader Settings	Directions	Drop Spreaders	Spreader Settings	Directions
Scotts Basic™, Standard™, Deluxe™ EdgeGuard®, Lawn Pro® or Speedy Green®	3 1/2	Each pass should be within 5 feet of each other to ensure overlap.	Scotts AccuGreen®	6 3/4	Make sure spreader wheel completely overlaps previous wheel marks.
Scotts Easy Green®	28	Each pass should be within 30 inches of each other to ensure overlap.			

For additional spreader settings please call 1-800-543-TURF (8873)

## GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline:  
**1-800-543-TURF**  
(8873)

Visit our website:  
**www.scotts.com**



### Well-fed lawns are more efficient at using available water.

Feeding with Scotts® fertilizer helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

## Kills most common lawn weeds\* DIRECTIONS FOR USE

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling.*  
This product applies 1.5 pounds of 2,4-D per acre per application.

### APPLICATION

Apply on a calm day. Make sure that the grass is wet from dew or sprinkling before applying. (This helps particles adhere to weed leaves to produce best results.) Set spreader to the setting indicated by the chart above before applying.

### WHAT TO AVOID

- Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.
- Do not apply more than twice per year. A minimum of 30 days between applications is required.
- Do not water or mow lawn for at least 24 hours after application. (Granules must remain on weed leaves for 24 hours to be effective.)
- Do not apply if rain is expected within 24 hours.
- Do not apply if day temperatures are regularly below 60° F or above 90° F.
- Apply only to grass.
- Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetables.

- Grass should not touch the bottom of a drop spreader. Tall grass should be mowed a day or two before application.
- Delay application on newly seeded, sodded or sprigged areas until new grass has been mowed four times.
- Do not seed, sod or plant grass sprigs until at least one month after using this product.
- Do not apply on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass.
- On bahia or centipede lawns, wait until after the second mowing of the year, but not before mid-March.
- In fall, DO NOT treat centipede grass after Oct. 1, nor bahia grass after Nov. 1.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical resistant gloves made of any waterproof material.

### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling poison control center or doctor or going for treatment.

### ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a cool, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE CONTAINER DISPOSAL:** If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

EPA Reg. No. 538-270 EPA Est. No. 538-OH-1  
†CAS #94-75-7 †† CAS #16484-77-8

STEP® 2 Weed Control Plus Lawn Fertilizer 28-0-6 F 643

### Guaranteed Analysis

Total nitrogen (N)	28%
4.9% ammoniacal nitrogen	
17.9% urea nitrogen	
5.7% other water soluble nitrogen*	
0.4% water insoluble nitrogen	
Soluble potash (K <sub>2</sub> O)	6%
Sulfur (S)	6%
6% combined sulfur (S)	

Derived from: Methyleneureas, potassium sulfate, ammonium sulfate, urea.

\*Contains 5.1% slowly available methylenediurea and dimethylenetriurea nitrogen.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

†Including dandelion, white clover, chickweed, plantain and ragweed.



The Scotts Company  
14111 Scottslawn Rd.  
Marysville, Ohio 43041

U.S. Patent No. 5,102,440

©2010 The Scotts Company LLC.

World rights reserved. Made in U.S.A.

### Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.



# Now it's Easy for You to Have a Better Lawn



## Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed.

## Simple and easy

No more guesswork. You simply apply the right product – one STEP® at a time.

## Guaranteed results

Each STEP® product is designed to help give you a greener lawn.

## Florida Annual Fertilizer Applications

On Centipede lawns, do not exceed rate of 2 lbs of Nitrogen per 1,000 sq. ft. in the north and 3 lbs of Nitrogen per 1,000 sq. ft. in the central and south. On Bahia lawns, do not exceed rate of 3 lbs of Nitrogen per 1,000 sq. ft. in the north.

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

## SPREADER SETTINGS

**Spreading Tips:** Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Covers 15,000 sq. ft. This package applies 2.93 lbs of product per 1,000 sq. ft.

Scotts Broadcast / Rotary Spreaders	Spreader Settings	Directions	Drop Spreaders	Spreader Settings	Directions
Scotts Basic™, Standard™, Deluxe™ EdgeGuard®, Lawn Pro® or Speedy Green®	3 1/2	Each pass should be within 5 feet of each other to ensure overlap.	Scotts AccuGreen®	6 3/4	Make sure spreader wheel completely overlaps previous wheel marks.
Scotts Easy Green®	28	Each pass should be within 30 inches of each other to ensure overlap.			



For additional spreader settings please call 1-800-543-TURF (8873)

## GET EXPERT ADVICE FOR YOUR LAWN NOW!



Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline:  
**1-800-543-TURF**  
(8873)



Visit our website:  
**www.scotts.com**



## Well-fed lawns are more efficient at using available water.

Feeding with Scotts® fertilizer helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

## Kills most common lawn weeds\* DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds of 2,4-D per acre per application.

### APPLICATION

Apply on a calm day. Make sure that the grass is wet from dew or sprinkling before applying. (This helps particles adhere to weed leaves to produce best results.) Set spreader to the setting indicated by the chart above before applying.

### WHAT TO AVOID

- Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.
- Do not apply more than twice per year. A minimum of 30 days between applications is required.
- Granules must remain on

- Grass should not touch the bottom of a drop spreader. Tall grass should be mowed a day or two before application.
- Delay application on newly seeded, sodded or sprigged areas until new grass has been mowed four times.
- Do not seed, sod or plant grass sprigs until at least one month after using this product.
- Do not apply on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass.
- On bahia or centipede/grass lawns, wait until after the second mowing of the year, but not before mid-March.
- In fall, DO NOT treat centipede/grass after Oct. 1, nor bahiagrass after Nov. 1.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

- Do not apply if rain is expected within 24 hours.
- Do not apply if day temperatures are regularly below 60° F or above 90° F.
- Apply only to grass.
- Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetables.

EPA Reg No. 538-270 (15M)

Step 2 Weed Control Plus Lawn Fertilizer

# KEEP OUT OF REACH OF CHILDREN

## CAUTION

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical resistant gloves made of any waterproof material.

#### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling poison control center or doctor or going for treatment.

#### ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a cool, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE CONTAINER DISPOSAL:** If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

EPA Reg. No. 538-270 EPA Est. No. 538-OH-1

+CAS #94-75-7 ++ CAS #16484-77-8

11/09/10 FPL

Page 4 of 4

STEP® 2 Weed Control Plus Lawn Fertilizer 28-0-6 F 643

#### Guaranteed Analysis

Total nitrogen (N) .....	28%
4.0% ammoniacal nitrogen	
17.9% urea nitrogen	
5.7% other water soluble nitrogen*	
0.4% water insoluble nitrogen	
Soluble potash (K <sub>2</sub> O) .....	6%
Sulfur (S) .....	6%
6% combined sulfur (S)	

Derived from: Methylenureas, potassium sulfate, ammonium sulfate, urea.

\*Contains 5.1% slowly available methylenediurea and dimethylenetriurea nitrogen.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

‡Including dandelion, white clover, chickweed, plantain and ragweed.



The Scotts Company  
14111 Scottslawn Rd.  
Marysville, Ohio 43041

U.S. Patent No. 5,102,440

©2010 The Scotts Company LLC.

World rights reserved. Made in U.S.A.

#### Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

STEP® 2 15M 3415-13



PROOF OF PURCHASE  
VID-BB  
P3415-13

EPA Reg No. 538-270 (15M)  
Step 2 Weed Control Plus Lawn Fertilizer