

WEED BEATER PLUS[®]

TRUSTED SINCE 1926

BONIDE[®]

Crabgrass & Broadleaf Weed Killer

Ready to Spray



Kills Crabgrass *PLUS* broadleaf weeds

Kills the weeds not the lawn

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 6.42%
Quinclorac..... 2.13%
Dicamba, dimethylamine salt 0.60%

OTHER INGREDIENTS: 90.85%

TOTAL 100.00%

See back panel for substatement of ingredient statement
EPA Est. No. 4-NY-1 EPA Reg. No. 4-466

Keep Out Of Reach Of Children

CAUTION (See inside booklet for additional
Precautionary Statements and First Aid)

Net Contents 32 FL. OZ. (946 ML.)

51000416



Kills to the root

WEED BEATER PLUS CRABGRASS & BROADLEAF WEED KILLER READY TO SPRAY

THIS PRODUCT CONTAINS:

0.456 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%

0.128 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%

0.043 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%

Isomer Specific by AOAC Method

- KILLS OVER 200 LAWN WEEDS PLUS CRABGRASS IN ONE EASY STEP
- ONE QUART COVERS UP TO 6,600 SQUARE FEET.

CONTROLS: CRABGRASS, DANDELION, BLACK MEDIC, CHICKWEED, CLOVER, CREEPING CHARLIE, GROUND IVY, LAMBSQUARTERS, MORNINGGLORY, OXALIS, POISON IVY, POISON OAK, SPEEDWELL, SPURGE, WILD CARROT, YARROW AND OTHER BROADLEAF WEEDS.

EPA Est. No. 4-NY-1

EPA Reg. No. 4-466

©Bonide Products, Inc. All Rights Reserved

Bonide Products, Inc.

6301 Sutliff Road

Oriskany, NY 13424

68510512 Quart



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING.

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grassy weeds including crabgrass, dandelion, foxtail, chickweed and clover. (see complete listing inside)

Where to Use: On Lawns: Kills the weeds, not the lawn when used as directed. Can be used on Kentucky bluegrass, Ryegrass (perennial and annual), Tall Fescue, Mixtures of cool season grasses, Bermudagrass* (common and hybrid), Zoysiagrass and Buffalograss.

When to Use: Apply when daytime temperatures are below 90°F.

Coverage: This container treats 6,600 sq. ft.

How it Works: This product contains three proven weed killers that target lawn weeds and crabgrass. It enters the lawn weeds through their leaves and moves throughout the plant to kill the weed, roots and all. Approved lawns listed on this label will not be harmed when used as directed.

*Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

HOW TO USE

- Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This 32 fl. oz. container will cover 6,600 square feet.
- Make sure sprayer nozzle is in "OFF" position then, connect sprayer to garden hose.
- Turn water on at faucet. Extend hose to the farthest area of lawn to be treated. To begin spraying, push lock tab down and turn knob control lever to "ON" position. The product mixes automatically with the water. Walk at a steady pace while spraying evenly and working back toward the faucet to avoid walking over wet, treated grass.
- To stop spraying, turn knob control lever to the "OFF" position. Turn off water at the faucet. To relieve pressure, turn knob control lever to the "ON" position and point the sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

Apply this product when weeds are small and actively growing.

Can be applied to the following turf grasses: Kentucky bluegrass, Ryegrass (perennial and annual), Tall Fescue, mixtures of cool season grasses, Bermudagrass* (common and hybrid), Zoysiagrass and Buffalograss.

*Do not apply to Bermudagrass lawns when daytime temperatures exceed 85°F. Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected.

The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

When to apply:

- When weeds are small and actively growing.
- When temperature is below 90°F.
- For Newly Seeded Lawns, do not apply until after the third mowing (approximately 6 weeks after seeding emergence)
- For Newly Sodded, Sprigged or Plugged Lawns, apply 3 to 4 weeks after planting.

Watering:

- Adequate soil moisture and favorable growing conditions enhance the performance of this product.
- Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule watering, before and after the application. For best results, delay the second watering until approximately 24 hours after the application.

Mowing:

- Delay mowing 2 days before and until 2 days after the application.
- Do not use clippings from the treated areas as mulch or compost, around flowers, ornamentals, trees, or in vegetable gardens.

Non-Target Plants:

- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers or other nontarget plants.

GRASS AND BROADLEAF WEEDS CONTROLLED

A partial list is presented below:

GRASS WEEDS

Crabgrass, large and smooth Foxtail, green, yellow and giant Signalgrass, broadleaf

BROADLEAF WEEDS

Annual yellow sweetclover	Buckhorn	Clover, white	Ground ivy
Aster	Buckhorn plantain	Cockle	Gumweed
Austrian fieldcress	Bulbous buttercup	Cocklebur	Hairy fleabane
Bedstraw	Bull thistle	Common mullein	Hawkweed
Beggarticks	Bullnettle	Creeping Jenny	Healall
Betony, Florida	Burclover	(Creeping Charlie)	Heartleaf drymary
Bindweed, field	Burdock	Cudweed	Heathaster
Bird vetch	Burning nettle	Curly dock	Hedge bindweed
Bitter wintercress	Burweed	Daisy, English	Hedge mustard
Bittercress, hairy	Buttercup	Daisy fleabane	Hemp
Bitterweed	Buttonweed	Daisy, oxeye	Henbit
Black-eyed Susan	Canada thistle	Dandelion	Hoary cress
Black medic	Carolina geranium	Dichondra	Hoary plantain
Black mustard	Carpetweed	Dogbane	Hoary vervain
Blackseed plantain	Catchweed bedstraw	Dogfennel	Horsenettle
Blessed thistle	Catnip	Dollarweed	Jimsonweed
Bloodflower milkweed	Catsear	Elderberry	Knawel
Blue lettuce	Chickweed, common	False dandelion	Knotweed
Blue vervain	Chickweed, mouseear	Falseflax	Kochia
Bracted plantain	Chicory	False sunflower	Lambsquarters
Brassbuttons	Cinquefoil	Fiddleneck	Lespedeza
Bristly oxtongue	Clover, crimson	Florida pusley	Mallow
Broadleaf dock	Clover, hop	Frenchweed	Matchweed
Broadleaf plantain	Clover, red	Galinsoga	Mexicanweed
Broomweed	Clover, strawberry	Goathead	Milk vetch
	Clover, sweet	Goldenrod	Morningglory

BROADLEAF WEEDS Cont.

Mouseear hawkweed	Prostrate pigweed	Spotted spurge	Wild lettuce
Mugwort	Prostrate spurge	Spurweed	Wild marigold
Musk thistle	Prostrate vervain	Stinging nettle	Wild mustard
Mustard	Puncturevine	Strawberry, India	Wild onion
Narrowleaf plantain	Purslane, common	mock	Wild parsnip
Narrowleaf vetch	Ragweed	Tall nettle	Wild radish
Nettle	Red sorrel	Tall vervain	Wild rape
Orange hawkweed	Redroot pigweed	Tansy ragwort	Wild strawberry
Oriental cocklebur	Redstem filaree	Tansy mustard	Wild sweet
Oxalis	Rough cinquefoil	Tanweed	potato
Parsley-piert	Rough fleabane	Thistle	Wild vetch
Parsnip	Russian pigweed	Trailing crownvetch	Wild violet
Pearlwort	Russian thistle	Tumble mustard	Woodsorrel
Pennycress	Scarlet pimpernel	Tumble pigweed	Woolly croton
Pennywort	Scotch thistle	Velvetleaf	Woolly
Peppergrass	Sheep sorrel	Venice mallow	morningglory
Pepperweed	Shepherdspurse	Virginia buttonweed	Woolly plantain
Pigweed	Slender plantain	Virginia creeper	Wormseed
Pineywoods bedstraw	Smallflower	Virginia pepperweed	Yarrow
Plains coreopsis (tickseed)	galinsoga	Water pennywort	Yellow rocket
Plantain	Smooth dock	Wavyleaf bullthistle	Yellowflower
Poison ivy	Smooth pigweed	Western clematis	pepperweed
Poison oak	Sorrel	Western salsify	and other
Pokeweed	Sowthistle	White mustard	broadleaf
Poorjoe	Spanishneedles	Wild aster	weeds
Poorjoe	Speedwell	Wild buckwheat	
Prairie sunflower	Spiny amaranth	Wild carrot	
Prickly lettuce	Spiny cocklebur	Wild four-o'clock	
Prickly sida	Spiny sowthistle	Wild garlic	
Prostrate knotweed	Spotted catsear	Wild geranium	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Avoid contamination of food or feedstuffs.

PESTICIDE DISPOSAL: If Empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

If in eyes:	<ul style="list-style-type: none">•Hold eye open and rinse slowly and gently with water for 15-20 minutes.•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.•Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">•Call a poison control center or doctor immediately for treatment advice.•Have person sip a glass of water if able to swallow.•Do not induce vomiting unless told to by a poison control center or doctor.•Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

LIMITED WARRANTY AND DISCLAIMER

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, seller's liability for default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, seller shall have no liability for consequential damages.