

1 Pint Treats
up to 2500 sq ft



Crab-E-Rad[®] Plus

**Kills Crabgrass &
All Major
Broadleaf Weeds**



Active Ingredients:

| | |
|-------------------|----------|
| 2,4-D DMA* | 6.557% |
| Quinclorac** | 4.085% |
| Dicamba Acid*** | 0.681% |
| Other Ingredients | 88.677% |
| TOTAL | 100.000% |

Isomer Specific Method, Equivalent to:

| | | |
|-----------------|--------|------------------|
| *2,4-D | 5.446% | 0.4591 lbs./gal. |
| **Quinclorac | 4.085% | 0.3443 lbs./gal. |
| ***Dicamba Acid | 0.681% | 0.0574 lbs./gal. |

EPA Reg. No. 228-595-54705 EPA Est. No. 48498-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION See Attached Booklet for Precautionary Statements, First Aid and Directions for Use

NET CONTENTS: 1 PINT/473.16 mL

(RV020510N) 0209/0817(02)

Crab-E-Rad[®] Plus

**For Selective Broadleaf and Certain
Grassy Weed Control In Residential and
Ornamental Turfgrasses**

**To Control Crabgrass, Dandelion, Clover,
Henbit, Plantains, Wild Onion and Many
Other Listed Lawn Weeds**

- **Rainproof in Hours – Won't Wash Away**

EPA Reg. No. 228-595-54705

EPA Est. No. 48498-CA-1



Distributed By:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745

(559) 499-2100

info@montereylawnngarden.com

(RV020510N) 0209/0817(02)



0 22179 10501 0

NET CONTENTS: 1 PINT / 473.16 mL

For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300
For Medical Emergencies Only,
Call (877) 325-1840

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves

USER SAFETY RECOMMENDATIONS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may

result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional).

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperatures over 90°F, turf injury may occur.

Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed the

maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

Do not apply this product by air or through any type of irrigation equipment.

Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

Make no more than 2 applications per year. For ornamental lawns and turf, apply no more than 0.5 oz. ae 2,4-D per 1,000 sq. ft. per application and no more than 1.0 oz. ae 2,4-D per 1,000 sq. ft. per year, excluding spot treatments.

This product combines three proven active ingredients [Quinclorac, 2,4-D, and Dicamba] to provide broad spectrum control of grassy and broadleaf weeds. This herbicide is absorbed by foliage and roots of weeds and translocated throughout the plant. **USE AS DIRECTED TO CONTROL OR SUPPRESS THE FOLLOWING WEEDS:**

WEEDS

GRASSY WEEDS CONTROLLED

| | |
|-------------------|-----------------|
| Crabgrass, Large | Foxtail, Green |
| Crabgrass, Smooth | Foxtail, Yellow |
| Barnyardgrass | Signalgrass, |
| Foxtail, Giant | Broadleaf |

BROADLEAF WEEDS CONTROLLED

| | |
|---------------------|--------------------|
| Alder | Broadleaf plantain |
| Annual yellow | Broomweed |
| sweet clover | Buckhorn |
| Artichoke | Buckhorn plantain |
| Aster | Bulbous buttercup |
| Austrian fieldcress | Bull nettle |
| Bedstraw | Bull thistle |
| Beggartick | Burdock |
| Biden | Burning nettle |
| Bindweed | Bur ragweed |
| Bird vetch | Burweed |
| Bitterweed | Buttercup |
| Bitter wintercress | Canada thistle |
| Black-eyed Susan | Carolina geranium |
| Black medic | Carpetweed |
| Black mustard | Catchweed |
| Blackseed plantain | bedstraw |
| Blessed thistle | Catsear |
| Blue lettuce | Catnip |
| Blue vervain | Chickweed |
| Box elder | Chicory |
| Bracted plantain | Cinquefoil |
| Brassbuttons | Common |
| Bristle oxtongue | chickweed |
| Broadleaf dock | Common mullein |

(continued)

| | |
|-------------------|-------------------|
| Common sowthistle | Ground ivy |
| Corn chamomile | Gumweed |
| Creeping Jenny | Hairy bittercress |
| Crimson clover | Hairy fleabane |
| Croton | Hawkweed |
| Cudweed | Healall |
| Curly dock | Heartleaf drymary |
| Curly indigo | Hedge bindweed |
| Dandelion | Hedge mustard |
| Dead nettle | Hemp |
| Dock | Henbit |
| Dollarweed | Hoary cress |
| Dogbane | Horsetail |
| Dogfennel | Indiana mallow |
| Elderberry | Ironweed |
| English daisy | Jewelweed |
| Fall dandelion | Jimsonweed |
| False dandelion | Kochia |
| False flax | Knawel |
| False sunflower | Knotweed |
| Fiddleneck | Lambsquarter |
| Field bindweed | Lespedeza |
| Field pansy | Locoweed |
| Flea bane (Daisy) | Lupine |
| Flixweed | Mallow |
| Florida betony | Marshelder |
| Florida pusley | Matchweed |
| Frenchweed | Mexicanweed |
| Galinsoga | Milk vetch |
| Garlic mustard | Milkweed |
| Goathead | bloodflower |
| Goatsbeard | Mugwort |
| Goldenrod | Morningglory |

| | |
|---------------------|--------------------|
| Mouseear chickweed | Redstem filaree |
| Musk thistle | Rough cinquefoil |
| Mustard | Rough fleabane |
| Narrowleaf plantain | Roundleafed |
| Narrowleaf vetch | marigold |
| Nettle | Rush |
| Orange hawkweed | Russian pigweed |
| Oxalis | Russian thistle |
| Oxeye daisy | St. Johnswort |
| Parsley-piert | Scarlet pimpernel |
| Parsnip | Scotch thistle |
| Pearlwort | Sheep sorrel |
| Peppergrass | Shepherdspurse |
| Pepperweed | Slender plantain |
| Pigweed | Smallflower |
| Pineywoods | galinsoga |
| bedstraw | Smartweed |
| Plains coreopsis | Smooth dock |
| Plantain | Smooth pigweed |
| Poison hemlock | Sneezeweed |
| Poison ivy | Southern wild rose |
| Poison oak | Sowthistle |
| Pokeweed | Spanishneedle |
| Poorjoe | Spatterdock |
| Povertyweed | Speedwell |
| Prostrate spurge | Spiny amaranth |
| Prostrate vervain | Spiny cocklebur |
| Puncture vine | Spotted catsear |
| Purslane | Spotted knapweed |
| Ragweed | Spotted spurge |
| Red clover | Spurge |
| Redroot pigweed | Spurweed |
| Red sorrel | Stinging nettle |

| | |
|----------------------|----------------------|
| Stinkweed | Wild aster |
| Stitchwort | Wild buckwheat |
| Strawberry clover | Wild carrot |
| Sumac | Wild four o'clock |
| Sunflower | Wild garlic |
| Sweet clover | Wild geranium |
| Tall nettle | Wild lettuce |
| Thistle | Wild marigold |
| Tick trefoil | Wild onion |
| Toadflax | Wild parsnip |
| Trailing crownvetch | Wild radish |
| Tumble mustard | Wild rape |
| Tumble pigweed | Wild strawberry |
| Tumbleweed | Wild sweet potato |
| Velvetleaf | Wild vetch |
| Venice mallow | Willow |
| Veronica | Witchweed |
| Vervain | Woodsorrel |
| Vetch | Woolly croton |
| Violet, Wild | Woolly morning-glory |
| Virginia buttonweed | Woolly plantain |
| Virginia creeper | Wormseed |
| Virginia pepperweed | Yarrow |
| Wavyleaf bullthistle | Yellow rocket |
| Western clematis | Yellowflower |
| Western salsify | pepperweed |
| White clover | and other broadleaf |
| White mustard | weeds |

Some of these species may require repeat spot applications and/or use of the maximum recommended rate on this product label even under ideal conditions for application.

USE TIPS

GENERAL TIPS

- Shake well before using.
- Spray during the growing season when weeds are young and growing actively.
- For best results, do not mow for 2 days before or after treating.
- Avoid entire lawn treatments when air temperature exceeds 90°F.
- Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees.
- Do not use on lawns with desirable clovers and legumes or on ornamentals.
- Avoid sprays which are too fine because they can drift and damage other plants.
- If certain tough weeds are not dead within 3 weeks, repeat with spot treatment application.

TIMING

Apply this product to young, actively growing weeds as a postemergence entire lawn or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds.

- Applications of this product made to annual grasses at 2 to 4 tiller may not provide complete control.

- **Newly Seeded Areas:** Application to grass seedlings is not recommended until after the third mowing.
- **Newly Sodded, Sprigged or Plugged Areas:** Application should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky bluegrass, perennial ryegrass and fine fescues.
- **Seeding:** Delay application for 4 weeks after seeding and emergence of turf species listed on this label.
- **Mowing:** Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

IRRIGATION AND RAINFALL

- If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results, do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

USE SITES

CRAB-E-RAD PLUS can be applied postemergence to the following turfgrasses.

| Highly Tolerant Turfgrasses | Moderately Tolerant Turfgrass | Do Not Apply To The Following Turfgrasses |
|--|--|---|
| COOL SEASON TURFGRASS Bluegrass, Kentucky Fescue, Tall Ryegrass, Annual Ryegrass, Perennial | COOL SEASON TURFGRASS Bluegrass, Annual (Poa Annua) (1) Bluegrass, Rough (Poa trivialis) Fescue, Chewing's Fescue, Fine (2) Fescue, Hard Fescue, Red WARM SEASON TURFGRASS Bermudagrass, Common (1) Bermudagrass (Hybrids) (1) Zoysiagrass | Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Buffalograss Carpetgrass Centipedegrass Dichondra Paspalum, Seashore St. Augustinegrass |
| <ol style="list-style-type: none"> 1. Temporary yellowing and discoloration on these species may occur after application. 2. Only apply to fine fescue when it is part of a blend. | | |

Warm season grasses such as Common Bermuda and Zoysia may see temporary discoloration. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

HOW MUCH TO USE

SPOT TREATMENT

Pressure Sprayer (Pump-up Sprayer):

Adjust spray nozzle to give a coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

| AMOUNT OF CONCENTRATE (Fluid Ounces) | AMOUNT OF WATER (Gallons) | AREA TREATED (SQUARE FEET) |
|--------------------------------------|---------------------------|----------------------------|
| 6.4 | 4 | 1,000 |
| 3.2 | 2 | 500 |
| 1.6 | 1 | 250 |

ENTIRE LAWN TREATMENT

Broadcast Application: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

| AREA TO BE TREATED (Square Feet) | AMOUNT OF PRODUCT NEEDED (Fluid Ounces) |
|-------------------------------------|--|
| 1,000 | 6.4 |
| 2,000 | 12.8 |
| 3,000 | 19.2 |
| 4,000 | 25.6 |
| 5,000 | 32.0 |

Ready-to-Spray Hose-End Sprayers:



[This product is packaged in a ready-to-spray packaging containing a plastic container fitted with a spray head that sprays the solution at a fixed concentrate.]

Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

This sprayer bottle will treat 5,000 sq. ft. when applied at 6.4 fluid ounces of this product per 1,000 sq. ft.

Instructions:

1. Attach garden hose to spray head.
2. Turn on water supply.
3. Bend safety tab down and turn control valve to "water".
4. Point nozzle toward lawn and turn control valve to "on". Product will automatically mix with water.
5. Walk back and forth at a steady pace and spray evenly over the lawn treatment area.



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Once clean, some agricultural pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE**

PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF

PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR USE BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Crab-E-Rad® Plus

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

EPA Reg. No. 228-595-54705
EPA Est. No. 48498-CA-1

Crab-E-Rad® is a registered trademark of Lawn and Garden Products, Inc.



Distributed By:
LAWN AND GARDEN PRODUCTS, INC.
P.O. Box 35000 • Fresno, CA 93745
(559) 499-2100
info@montereylawnngarden.com