

1 Pint Treats
up to 2500 sq ft



2,4-D, DICAMBA, QUINCLORAC GROUP 4 HERBICIDE

Crab-E-Rad[®] Plus

**Kills Crabgrass
& All Major
Broadleaf Weeds**



Active Ingredients:

2,4-D DMA*	6.557%
Quinclorac**	4.085%
Dicamba Acid***	0.681%
Other Ingredients	88.677%
TOTAL	100.000%

Isomer Specific Method, Equivalent to:

*2,4-D	5.446%	0.4591 lbs./gal.
**Quinclorac	4.085%	0.3443 lbs./gal.
***Dicamba Acid	0.681%	0.0574 lbs./gal.

EPA Reg. No.228-595-54705 EPA Est.No.48498-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION See Inside Booklet for First Aid and Additional
Precautionary Statements and Directions for Use

NET CONTENTS: 1 PINT/473.16 mL

(RV012424) 0209/0624(03)

Crab-E-Rad® Plus



**For Selective Broadleaf and Certain Grassy Weed
Control In Residential and Ornamental Turfgrasses
To Control Crabgrass, Dandelion, Clover, Henbit,
Plantains, Wild Onion and Many Other Listed
Lawn Weeds**

- **Rainproof in Hours – Won't Wash Away**

EPA Reg. No. 228-595-54705

EPA Est. No. 48498-CA-1

See Inside Booklet for First Aid and Additional
Precautionary Statements and Directions for Use

Distributed By:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745

(559) 499-2100

info@montereylawngarden.com

Crab-E-Rad® is a registered trademark of
Lawn and Garden Products, Inc.

(RV012424) 0209/0624(03)



NET CONTENTS: 1 PINT / 473.18 mL

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find someone to explain it to you in detail)

For Chemical Spill, Leak, Fire, or Exposure, Call
CHEMTREC (800) 424-9300

For Medical Emergencies Only,
Call (877) 325-1840

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Waterproof gloves

USER SAFETY RECOMMENDATIONS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Drift or runoff may adversely affect nontarget plants. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas

below the mean high water mark. **DO NOT** apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, **DO NOT** pour washwater on the ground; spray or drain over a large area away from wells and other water sources. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. **DO NOT** apply this product through any type of irrigation system. **DO NOT** contaminate water used for irrigation or domestic purposes.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional).

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Handheld Technology Applications

Take precautions to minimize spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with

altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

RESISTANCE MANAGEMENT

This product is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly on the same use site. Appropriate resistance-management strategies should be followed. To delay herbicide resistance take one or more of the following steps:

- Rotate this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.

- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other sites by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- For further information or to report suspected resistance, contact Lawn and Garden Products, Inc. at 559-499-2100.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action

thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

For Ornamental Turfgrass Sites

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash from treated areas onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

USE RESTRICTIONS

- **DO NOT** use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- **DO NOT** apply this product by air or through any type of irrigation equipment.
- **DO NOT** make more than 3 applications per year, with a minimum of 21 days between applications.
- **DO NOT** apply more than 6.4 fl oz. per 1000 sq. ft. (1.0 lb 2,4-D ae, 0.75 lb quinclorac, 0.125 lb dicamba per acre) in a single application.
- **DO NOT** apply more than 19.2 fl. oz per 1000 sq. ft. (3.0 lb 2,4-D ae, 2.25 lb quinclorac, 0.375 lb dicamba per acre) per year.
- **DO NOT** apply when wind speeds are greater than 10 mph at the application site.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperatures over 90°F, turf injury may occur.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre (or rate/1,000 sq. ft.) which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions

for use and precautionary statements of each product in the tank mixture.

Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

This product combines three proven active ingredients [Quinclorac, 2,4-D, and Dicamba] to provide broad spectrum control of grassy and broadleaf weeds. This herbicide is absorbed by foliage and roots of weeds and translocated throughout the plant. USE AS DIRECTED TO CONTROL OR SUPPRESS THE FOLLOWING WEEDS:

WEEDS:

GRASSY WEEDS

Crabgrass, Large

Crabgrass, Smooth

Barnyardgrass

Foxtail, Giant

Foxtail, Green

Foxtail, Yellow

Signalgrass, Broadleaf

BROADLEAF WEEDS

Alder	Burdock
Annual yellow sweet clover	Burning nettle
Artichoke	Bur ragweed
Aster	Burweed
Austrian fieldcress	Buttercup
Bedstraw	Canada thistle
Beggartick	Carolina geranium
Biden	Carpetweed
Bindweed	Catchweed bedstraw
Bird vetch	Catsear
Bitterweed	Catnip
Bitter wintercress	Chickweed
Black-eyed Susan	Chicory
Black medic	Cinquefoil
Black mustard	Common chickweed
Blackseed plantain	Common mullein
Blessed thistle	Common sowthistle
Blue lettuce	Corn chamomile
Blue vervain	Creeping Jenny
Box elder	Crimson clover
Bracted plantain	Croton
Brassbuttons	Cudweed
Bristle oxtongue	Curly dock
Broadleaf dock	Curly indigo
Broadleaf plantain	Dandelion
Broomweed	Dead nettle
Buckhorn	Dock
Buckhorn plantain	Dollarweed
Bulbous buttercup	Dogbane
Bull nettle	Dogfennel
Bull thistle	Elderberry
	English daisy

BROADLEAF WEEDS *(cont.)*

Fall dandelion	Jewelweed
False dandelion	Jimsonweed
False flax	Kochia
False sunflower	Knawel
Fiddleneck	Knotweed
Field bindweed	Lambsquarter
Field pansy	Lespedeza
Flea bane (Daisy)	Locoweed
Flixweed	Lupine
Florida betony	Mallow
Florida pusley	Marshelder
Frenchweed	Matchweed
Galinsoga	Mexicanweed
Garlic mustard	Milk vetch
Goathead	Milkweed bloodflower
Goatsbeard	Mugwort
Goldenrod	Morningglory
Ground ivy	Mousear chickweed
Gumweed	Musk thistle
Hairy bittercress	Mustard
Hairy fleabane	Narrowleaf plantain
Hawkweed	Narrowleaf vetch
Healall	Nettle
Heartleaf drymary	Orange hawkweed
Hedge bindweed	Oxalis
Hedge mustard	Oxeye daisy
Hemp	Parsley-piert
Henbit	Parsnip
Hoary cress	Pearlwort
Horsetail	Peppergrass
Indiana mallow	Pepperweed
Ironweed	Pigweed

BROADLEAF WEEDS *(cont.)*

Pineywoods bedstraw	Smooth dock
Plains coreopsis	Smooth pigweed
Plantain	Sneezeweed
Poison hemlock	Southern wild rose
Poison ivy	Sowthistle
Poison oak	Spanishneedle
Pokeweed	Spatterdock
Poorjoe	Speedwell
Povertyweed	Spiny amaranth
Prostrate spurge	Spiny cocklebur
Prostrate vervain	Spotted catsear
Puncture vine	Spotted knapweed
Purslane	Spotted spurge
Ragweed	Spurge
Red clover	Spurweed
Redroot pigweed	Stinging nettle
Red sorrel	Stinkweed
Redstem filaree	Stitchwort
Rough cinquefoil	Strawberry clover
Rough fleabane	Sumac
Roundleafed marigold	Sunflower
Rush	Sweet clover
Russian pigweed	Tall nettle
Russian thistle	Thistle
St. Johnswort	Tick trefoil
Scarlet pimpernel	Toadflax
Scotch thistle	Trailing crownvetch
Sheep sorrel	Tumble mustard
Shepherdspurse	Tumble pigweed
Slender plantain	Tumbleweed
Smallflower galinsoga	Velvetleaf
Smartweed	Venice mallow

BROADLEAF WEEDS *(cont.)*

Veronica	Wild onion
Vervain	Wild parsnip
Vetch	Wild radish
Violet, Wild	Wild rape
Virginia buttonweed	Wild strawberry
Virginia creeper	Wild sweet potato
Virginia pepperweed	Wild vetch
Wavyleaf bullthistle	Willow
Western clematis	Witchweed
Western salsify	Woodsorrel
White clover	Wooly croton
White mustard	Wooly morning-glory
Wild aster	Woolly plantain
Wild buckwheat	Wormseed
Wild carrot	Yarrow
Wild four o'clock	Yellow rocket
Wild garlic	Yellowflower pepperweed and other broadleaf weeds
Wild geranium	
Wild lettuce	
Wild marigold	

Some of these species may require repeat spot applications and/or use of the maximum recommended rate on this product label even under ideal conditions for application.

USE TIPS

GENERAL TIPS

- Shake well before using.
- Spray during the growing season when weeds are young and growing actively.
- For best results, do not mow for 2 days before or after treating.
- Avoid entire lawn treatments when air temperature exceeds 90°F.
- Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees.
- Do not use on lawns with desirable clovers and legumes or on ornamentals.
- Avoid sprays which are too fine because they can drift and damage other plants.
- If certain tough weeds are not dead within 3 weeks, repeat with spot treatment application.

TIMING

Apply this product to young, actively growing weeds as a postemergence entire lawn or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds.

- Applications of this product made to annual grasses at 2 to 4 tiller may not provide complete control.
- **Newly Seeded Areas:** Application to grass seedlings is not recommended until after the third mowing.
- **Newly Sodded, Sprigged or Plugged Areas:** Application should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging

operations. Delay applications for 4 weeks after seeding and emergence of Kentucky bluegrass, perennial ryegrass and fine fescues.

- **Seeding:** Delay application for 4 weeks after seeding and emergence of turf species listed on this label.
- **Mowing:** Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

IRRIGATION AND RAINFALL

- If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results, do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

USE SITES

This product can be applied postemergence to the following turfgrasses.

Highly Tolerant Turfgrasses	Moderately Tolerant Turfgrasses	Do Not Apply To The Following Turfgrasses
COOL SEASON TURFGRASS Bluegrass, Kentucky Fescue, Tall Ryegrass, Annual Ryegrass, Perennial	COOL SEASON TURFGRASS Bluegrass, Annual (Poa Annua) (1) Bluegrass, Rough Poa trivialis) Fescue, Chewing's Fescue, Fine (2) Fescue, Hard Fescue, Red WARM SEASON TURFGRASS Bermudagrass, Common (1) Bermudagrass (Hybrids) (1) Zoysiagrass	Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Buffalograss Carpetgrass Centipedegrass Dichondra Paspalum, Seashore St. Augustinegrass
<ol style="list-style-type: none"> 1. Temporary yellowing and discoloration on these species may occur after application. 2. Only apply to fine fescue when it is part of a blend. 		

Warm season turfgrasses such as Common Bermuda and Zoysia may see temporary discoloration. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

HOW MUCH TO USE

SPOT TREATMENT

Pressure Sprayer (Pump-up Sprayer): Adjust spray nozzle to give a coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

AMOUNT OF CONCENTRATE (Fluid Ounces)	AMOUNT OF WATER (Gallons)	AREA TREATED (SQUARE FEET)
6.4	4	1,000
3.2	2	500
1.6	1	250

ENTIRE LAWN TREATMENT

Broadcast Application: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

AREA TO BE TREATED (Square Feet)	AMOUNT OF PRODUCT NEEDED (Fluid Ounces)
1,000	6.4
2,000	12.8
3,000	19.2
4,000	25.6
5,000	32.0

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal.

(continued)

STORAGE AND DISPOSAL (cont.)

Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR USE BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Crab-E-Rad® Plus



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and gloves.

EPA Reg. No. 228-595-54705

EPA Est. No. 48498-CA-1

See Inside Booklet for First Aid and Additional
Precautionary Statements and Directions for Use

Distributed By:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745

(559) 499-2100

info@montereylawngarden.com

Crab-E-Rad® is a registered trademark of
Lawn and Garden Products, Inc.