

TRUSTED SINCE 1926  
**BONIDE**

**DuraTurf**

# INSECT & FEED

with *slow release* (N) FERTILIZER  
Treats up to 5,000 sq. ft.  
**12-0-10**

SUMMER  
PHASE  
**3**

Kills listed insects

Slow-release nitrogen-helps  
provide extended feeding

Phosphate-free fertilizer

Phase 3 of a 4 phase process *for best results*

**Kills listed insects  
and fertilizes the lawn**



Contains iron for deep greening

Keep Out Of Reach Of Children  
**CAUTION**  
Net Wt. 16.65 LBS (7.55 kg)

25 seal area -- one col common int ok -- no copy

TRUSTED SINCE 1926  
**BONIDE**

BOTTOM  
GUSSET

**DuraTurf**

# INSECT & FEED



## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

Do not make applications during the rain.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Do not allow children and pets to enter the area until dry.

Do not apply more than 13.3 lbs. per 1,000 square feet, of this product per year. Not for use on turf being grown for sale or other commercial use as sod, or for commercial greenhouses, nurseries, commercial seed production or research purposes.

This product should be applied when insects first appear. Application of this product should be applied evenly with a properly calibrated drop or rotary spreader over the entire turfgrass area at a rate of 3.33 to 6.66 pounds per 1,000 square feet. A uniform application is most desirable and is best achieved by spreading half of the required amount of products over the area and then applying the remaining half with swaths at a right angle to the first. A thorough irrigation of the treated area, immediately after application will provide best results.

### Restrictions concerning use near waters.

Do not apply this product within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, natural estuaries, and commercial fish ponds.

The contents of this 16.65 lb. bag feeds up to 5,000 sq. ft.

## LAWN AND PERIMETER APPLICATIONS

This product is intended to prevent and reduce insect entry to homes and businesses. When using this product as directed, apply a 6 to 10 feet wide band to the lawn areas around and adjacent to the structure. Observe and apply the following application rates.

### LAWN APPLICATION

Apply 3.33 lbs. per 1,000 sq.ft.	Apply 6.66 lbs. per 1,000 sq.ft.
Annual Bluegrass,	**Ants,
Weevils (Hyperodes Adult)	**Centipedes,
Armyworms,	*Fire Ants,
Billbugs (Adult),	**Fleas,
Black Turfgrass	Leafhoppers,
Ataenus (Adult),	**Millipedes,
Cutworms,	**Mole Crickets,
Mealybugs,	**Chinch Bugs,
Sod Webworms,	Ticks and Chiggers

\*This product will provide optimum control of Imported Fire Ants if combined with a mound treatment. A broadcast application of the area to be treated will control foraging worker ants and newly mated queens. In addition a mound treatment at the same time will aid in the destruction of the existing colony. For controlling mounds use a registered pesticide labeled for treatment of fire ant mounds.

\*\*The highest rate noted on the application table may be required to control Imported Fire Ants as well as Chiggers, Centipedes, Chinch Bugs, Fleas, Millipedes and other foraging ants.

In New York State this product may not be applied to any grass or turfgrass within 100 ft. of a water body (lake, pond, river, stream, wetland, drainage ditch). In New York State do make a single repeat application if there are signs of renewed insect activity, but not sooner than 2 weeks after the first application.

Florida Applications: Apply at the rate of 3.33 pounds per 1,000 sq. ft. Do not exceed 2 applications per year.

BACK PANEL

## SPREADER SETTINGS

These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

Drop Spreader	
Scotts AccuGreen	6.5
Republic EZ #530	6
Sears Drop	4
Jonathan Green/Precision	1.5
Rotary Spreaders	
Scotts Deluxe	5
Republic EZ #540	3
Earthway	11.5
Sears Rotary	5.5
Jonathan Green/Precision	3.5

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in its original container in a cool, dry, locked place out of the reach of children and animals. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Never pour unused product down any indoor or outdoor drain. **CONTAINER DISPOSAL:** If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not rinse unless required for recycling. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. If partly filled: Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain.

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco, or using toilet.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift and runoff from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

<b>Guaranteed Analysis</b>	<b>12-0-10</b>
Total Nitrogen (N) .....	12.00%
12.00% Urea Nitrogen*	
Soluble Potash (K <sub>2</sub> O) .....	10.00%
Boron (B) .....	0.02%
Copper (Cu) .....	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe) .....	1.00%
1.00% Water Soluble Iron (Fe)	
Manganese (Mn) .....	0.05%
0.05% Water Soluble Manganese (Mn)	
Molybdenum (Mo) .....	0.0005%
Zinc (Zn) .....	0.05%
0.05% Water Soluble Zinc (Zn)	

Derived from: Urea, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Copper Sulfate, Ferrus Sulfate, Manganese Sulfate, Zinc Sulfate, Sodium Borate, Sodium Molybdate

\*2.40% Slowly Available Nitrogen from Polymer Coated-Sulfur Coated Urea. F1290

**CAUTION -** Boron and molybdenum are recommended for all crops where boron and molybdenum may be deficient; however, excessive application to susceptible crops may cause damage. Crops high in molybdenum may be toxic to ruminant animals.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 0.4 to 0.8 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this container, and referring to the batch number found on this container.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Distributed and Guaranteed by  
Borde Products, Inc.  
6301 Sull Road  
Orlando, NY 13424  
©Borde Products, Inc.  
All Rights Reserved



25" seal area-- one col common ink ok -- no copy