ACE GREEN JURE. Grabgrass Preventer with Fertilizer

CE GREEN TURF. **COVERS 15,000 SQUARE FEET**

7229552

FACE PANEL GREEN TURF.

Crabgrass Preventer with Fertilizer

PREVENTS WEEDS* AND FERTILIZES THE LAWN

Fertilizante para Evitar el Garranchuelo

For control of crabgrass and control or suppression of many other listed annual grass and broadleaf weeds in established lawns and ornamental turf.

30-0-4

Net Wt. 40.5 lbs. (18.4 kg)

Prevents weeds* and feeds your lawn for up to 4 months

Promotes root growth for a thicker, greener lawn

KEEP OUT OF REACH OF CHILDREN
CAUTION
back panel for First Aid and additional Precautionary Statem

Net Wt. 40.5 lbs. (18.4 kg)

Crabgrass Preventer with Fertilizer

PREVENTS WEEDS* AND FERTILIZES THE LAWN

Fertilizante para Evitar el Garranchuelo

For control of crabgrass and control or suppression of many other listed annual grass and broadleaf weeds in established lawns and ornamental turf.

30-0-4

Net Wt. 40.5 lbs. (18.4 kg)

Prevents weeds*
and feeds your
lawn for up to
4 months

Promotes root growth for a thicker, greener lawn

 *See Use Directions for complete list of treated weeds.

In the state of New York, do not exceed 3.0 lb. of this product per 1000 sq. ft. per year (equivalent to 0.25 pounds of active ingredient per acre.)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for First Aid and additional Precautionary Statements.



PREVENTS WEEDS* AND FERTILIZES THE LAWN

30-0-4

Fertilizante para Evitar el Garranchuelo

Net Wt. 40.5 lbs. (18.4 kg)

NOTICE: Read the entire label. Use only according to label directions. Before buying this product, read the Warranty Section elsewhere on this label. If terms are not acceptable, return the unopened container at once.

Precautionary Statements

Hazards to Humans and Domestic Animals CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If in eves:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call (888) 208-1368.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies and drainage systems.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Intended for Outdoor Residential Use. **Read all directions for use carefully before applying.**

How This Product Works

This product can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control.

This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

COOL-SEASON GRASSES		WARM-SEASON GRASSES		
0 , 0	scue, tall egrass, perennial	Bahiagrass Bermudagrass Buffalograss Carpetgrass	Centipedegrass St. Augustinegrass Zoysiagrass	

USE PRECAUTIONS

This product must be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, use the initial application of this product only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

Do not use on turf grown for agricultural or commercial purposes (turf farms), or for grass seed production.

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress
 due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, are to be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product
 within 10 weeks after a single application of this product or within 16 weeks after a split application program
 totaling 6.0 lb./1000 sq. ft. (0.5 lb. active ingredient per acre) or more may inhibit the establishment of desirable
 turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding,
 irrigation and fertilization.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.

WEEDS CONTROLLED OR SUPPRESSED

Ace Green Turf Crabgrass Preventer with Fertilizer will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

Grasses: barley barnyardgrass bluegrass, annual brome	crabgrass, large crabgrass, smooth crabgrass, Southern crowfootgrass †	dallisgrass goosegrass foxtail, green foxtail, yellow	kikuyugrass † oats, wild ryegrass (annual & perennial)	sandbur smutgrass
Broadleaf Weeds: bittercress † carpetweed chickweed † geranium, Carolina † henbit	knotweed, prostrate lespedeza, common † marestail medic, black mustard	oxalis, buttercup pineappleweed pigweed, redroot parsley-piert † purslane, common	rocket, London shepherdspurse speedwell, corn † spurge, garden spurge, prostrate	spurge, spotted woodsorrel, creeping woodsorrel, yellow

[†] Suppression only

HOW TO APPLY

To optimize product performance apply this product within a few days after mowing and delay mowing again for a few days after the application. Most effective weed control will be obtained when this product is activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

SPREADER SETTINGS 2.7 lbs. per 1,000 sq. ft.						
Rotary Spreaders	Setting	Drop Spreaders	Setting			
Ace 2030AT Rotary	14	Scotts Accugreen 1000 Drop	7			
Earthway 2000A Rotary	14	Scotts Accugreen 2000 Drop	7			
Earthway 2050P Rotary	14	Scotts Accugreen 3000 Drop	7			
Scotts SpeedyGreen 1000 Rotary	4-1/2					
Scotts SpeedyGreen 2000 Rotary	4-1/2					
Scotts SpeedyGreen 3000 Rotary	4-1/2					

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

Determining Your Lawn's Square Footage

To determine how much of this product to apply, you must determine the square footage of your lawn. An easy way to do this is to walk the length of your lawn, figuring that one pace equals about 3 feet. Do the same with the width of the lawn. Then multiply the length by the width to arrive at the total. Make sure you subtract the square footage of your home and driveway when calculating the total square footage of your lawn. If your lawn has an irregular shape, try dividing it up into a number of square areas. Calculate the square footage for each area, then add the figures together to reach your total lawn size.

HOW MUCH TO APPLY

This product will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied.

Refer to the map to the right to determine the region of the U.S. that includes your location. Use the table below to determine the amount of this product to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.



MAXIMUM APPLICATION RATES

• Do not apply more than 6.0 lb. of this product per 1,000 sq. ft. per application and no more than 18.0 lb. of this product/1000 sq. ft. per year (equivalent to 0.5 and 1.5 lb. of active ingredient per acre, respectively).

In the state of New York, do not exceed 3.0 lb. of this product per 1000 sq. ft. per year (equivalent to 0.25 pounds of active ingredient per acre).

WHEN TO APPLY

Spring Application

Control of Crabgrass: This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

Fall Application

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.

Storage And Disposal

Do not contaminate water, food or feed by storage and disposal. Pesticide Storage: Store this product only in its original container in a cool, dry, locked place out of the reach of children and animals. Pesticide Disposal and Container Handling: Nonrefillable Container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

^{† 1,000} square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-lebsea.com.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.



QUESTIONS OR COMMENTS?

Call 1-800-233-1067

WARRANTY

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Lebanon Seaboard Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lebanon Seaboard Corporation DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

Manufactured by: Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042

EPA Reg. No. 961-417
EPA Est. No. 961-PA-01(), 961-IL-01()
Circled letter is first letter of lot number.

Made in the USA for Ace Hardware Corp., Oak Brook, Illinois, 60523, USA ©2011

Green Turf® is a registered trademark of United Industries Corporation.

* See Use Directions for complete list of treated weeds.

The Ace® Promise

If this Ace product when used for its intended purpose, fails to give you complete satisfaction, return the item to the Ace Hardware store where purchased for free replacement of the same or similar item.

El Compromiso de Ace®

Si este producto Ace, cuando se usa para los fines entendidos, no funciona a su entera satisfacción, devuélvalo a la ferretería de Ace Hardware donde lo compró para que se lo reemplacen sin costo por uno igual o similar.

