

OUR BEST ANNUAL PROGRAM

Scotts®

4STEP™ PROGRAM



4STEP™ PROGRAM

STEP 1 for Seeding

Starter® Lawn Food with Weed Preventer
Apply when Planting Grass Seed

Covers 5,000 SQ FT

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back for additional precautionary statements.

ACTIVE INGREDIENT:
Mesotrione 0.08%
OTHER INGREDIENTS..... 99.92%
Total 100.00%

21-22-4

NET WEIGHT 21.52 lb (9.76 kg)

Available only at authorized Scotts®


LawnPro®

Dealers

STEP 1® for Seeding

Starter® Lawn Food
with Weed Preventer

Apply when Planting
Grass Seed



Scotts® 4STEP PROGRAM

STEP 1 for Seeding

Starter® Lawn Food with Weed Preventer
Apply when Planting Grass Seed



EPA Reg No. 538-317 (5M)

Scotts Step 1 for Seeding Starter Lawn Food with Weed Preventer

NO VITAL COPY
BOTTOM SEAL AT 13



36905-1

100% SIZE
1.2-oz (35.23g) UPC-4
(Not for resale)

For Position Only

0 32247 36905 1

Scotts®

4 STEP™ PROGRAM

Plan your feeding schedule. Choose the right product for your lawn. Now it's Easy for You to Get a Better Lawn.



Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed. We recommend you apply the remaining products at two-month intervals.

Simple and easy

No more guesswork. You simply apply the right product – one at a time.

Guaranteed results

Each STEP® product is designed to the specific season to help give you a greener lawn.

SPREADING TIPS

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Shut off spreader or stop each time you reach a header strip. This package applies 4.3 lbs of product per 1,000 sq. ft. Covers 5,000 sq. ft.

Scotts® Broadcast/ Rotary Spreaders

Spreader Settings

4 1/2

Each pass should be within 5 feet of each other to ensure an overlap.



Scotts® Drop Spreaders

Spreader Settings

7

Make sure spreader wheel completely overlaps previous wheel marks



For additional spreader settings please call 1-800-543-TURF (8873)

GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.



Consumer Helpline:
1-800-543-TURF
(8873)



Visit our website:
www.scotts.com



Well-fed lawns are more efficient at using available water.

Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

PRECAUTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

INFORMATION

For use on newly seeded, overseeded, or newly planted Kentucky bluegrass, tall fescue, perennial ryegrass, centipede grass, as well as fine, spreading and creeping red fescues. This product is for application to home lawns and applies 0.15 pounds of mesotrione per acre per application. The maximum application rate of mesotrione per treatment site per year is 0.445 pounds. This package applies 4.3 pounds of product per 1,000 sq. ft.

Apply this product before watering newly seeded, overseeded or planted grass for the first time. New weeds will sprout white or tan white and quickly die. This product will not kill existing weeds.

USE PRECAUTIONS

Do not use this product in a way that will contact any person, or pet either directly or through drift. Keep people and pets out of the area during application. Do not enter or allow people or pets to enter the treated area (except those involved in the watering) until the watering-in is complete and the surface is dry. Do not use this product on flowers, shrubs, trees, fruit or vegetable plants, Canna, day lilies and irises are especially susceptible to injury. Do not apply another pesticide within seven days of an

application of this product as turf injury may occur. Do not apply to St. Augustinegrass, bermgrass, bermudagrass, zoysiagrass, dicentra, kikyo grass, bahiagrass or fescue.

APPLICATION INSTRUCTIONS

Apply on a calm day and lightly water to activate. Apply when temperatures are consistently between 60° and 90°. Avoid applying to adjacent flower beds when using a rotary spreader.

This product may cause temporary whitening of grass foliage. In general, symptoms appear five to seven days after application and may last up to 14 days. Discoloration is most likely to occur on ryegrass lawns.

WEEDS PREVENTED

Large crabgrass, Smooth crabgrass, Dandelion, Broadleaf plantain, Buckhorn plantain, Carpetweed, Common chickweed, Common lambsquarters, Common purslane, Corn speedwell, Ground ivy, Hairy bitertress, Henbit, Parsley plant, Pennsylvania smartweed, Persian speedwell, Pokedweed, Pursh speedwell, Redroot pigweed, Shepherd's-purse, Vervetleaf, White clover, Yellow woodsoot, Flatsedge, Yellow nutgrass.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place. Avoid bag by holding top down and securing. **CONTAINER DISPOSAL:** Material in this container, do not reuse or refill this container. If Empty: Place in trash or other for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never reuse unused product down any indoor or outdoor drain.

EPA Reg. No. 538-117
FQSA 104206-02-0

EPA Reg. No. 538-016-11, 538-016-11, 81665-016-11
Superscript used is first letter of ran code on end of container.

STEP 1 for Seeding Starter Lawn Food with Weed Preventer 21-42-4 F643

Guaranteed Analysis

Total Nitrogen (N) 21%

4.5% Ammoniacal Nitrogen

8.1% Urea Nitrogen

0.6% Other Water-Soluble Nitrogen*

0.6% Water-Soluble Nitrogen*

Available Phosphate (P2O5) 22%

Soluble Potash (K2O) 4%

Sulfur (S) 4%

2.0% Combined Sulfur (S)

Iron (Fe) Total 0.50%

0.50% Water-Soluble Iron (Fe)

Derived from: Methylenesulfone, Urea, Potassium Sulfate, Ammonium Phosphate and Ferrus Sulfate.

*Contains 5.0% slowly available nitrogen from methylenesulfone.

Information regarding the contents and level of metals in this product is available on the internet at <http://www.regulatore-info-sc.com>

Questions / Information

UMS and Manufactured by the Scotts Company
1415 Scotts Lane, Bay
Mansfield, Ohio 43041

Consumer Helpline: 1-800-543-TURF (8873)

©2012 The Scotts Company LLC.

World rights reserved.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to your money back. Simply send us proof of purchase and we will promptly mail you a refund.

SCOTTS LAWN FOOD STEP 1 FOR SEEDING STARTER LAWN FOOD WITH WEED PREVENTER 21-42-4 F643

120% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

0 32247 36905 3

PROOF OF PURCHASE
VP-00
BG1001

OUR BEST
ANNUAL
PROGRAM

Scotts[®]

4STEP[™]
PROGRAM

Covers 5,000 SQ FT

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back for additional precautionary statements.

ACTIVE INGREDIENT:

Mesotrione	0.09%
OTHER INGREDIENTS	99.91%
Total	100.00%

STEP[®]**1**
for Seeding

EPA Reg No. 538-317 (5M)

Scotts Step 1 for Seeding Starter Lawn Food with Weed Preventer

21-22-4

NET WEIGHT 21.52 lb (9.76 kg)


Available only at authorized Scotts®

LawnPro®

Dealers

Starter® Lawn Food with Weed Preventer

Apply when Planting
Grass Seed



4 STEP
PROGRAM

STEP 1 for Seeding

Starter® Lawn Food with Weed Preventer

Apply when Planting Grass Seed



EPA Reg No. 538-317 (5M)

Scotts Step 1 for Seeding Starter Lawn Food with Weed Preventer

5905-1

3/28/12 SSL

Page 4 of 6

NO VITAL COPY OF
BOTTOM SEAL IS AT 1/4

Plan your feeding schedule. Choose the right product for your lawn. Now it's Easy for You to Have a Better Lawn.



Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed. We recommend you apply the remaining products at two-month intervals.

Simple and easy

No more guesswork. You simply apply the right product – one at a time.

Guaranteed results

Each STEP® product is designed to the specific season to help give you a greener lawn.

SPREADING TIPS

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader. Shut off spreader or stop each time you reach a header strip. This package applies 4.3 lbs of product per 1,000 sq. ft. Covers 5,000 sq. ft.

Spreader Settings

Spreader Settings

Scott's® Broadcast/
Rotary Spreaders

4 1/2

Each pass should be within 5 feet of each other to ensure an overlap.



Scott's®
Drop Spreaders

7

Make sure spreader wheel completely overlaps previous wheel marks.



For additional spreader settings please call 1-800-543-TURF (8873)

GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.



Consumer Helpline:
1-800-543-TURF
(8873)



Visit our website:
www.scotts.com



Well-fed lawns are more efficient at using available water.

Feeding with Scott's® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

INFORMATION

For use on newly seeded, overseeded, or newly planted Kentucky bluegrass, tall fescue, perennial ryegrass, centipede grass, as well as fine, spreading and creeping red fescues.

This product is for application to home lawns and applies 0.15 pounds of meclozolin per acre per application. The maximum application rate of meclozolin per treatment site per year is 0.45 pounds. This package applies 4.3 pounds of product per 1,000 sq. ft.

Apply this product before watering newly seeded, overseeded or planted grass for the first time.

Now seeds will sprout white or turn white and quickly die. This product will not kill existing weeds.

USE PRECAUTIONS

Do not apply this product in a way that will contact any person, or pet either directly or through drift. Keep people and pets out of the area during application. Do not enter or allow people or pets to enter the treated areas (except those involved in the watering) until the watering is complete and the surface is dry. Do not use this product on flowers, shrubs, trees, fruit or vegetable plants. Cars, day lilies and irises are especially susceptible to injury. Do not apply another pesticide within seven days of an

application of this product as turf injury may occur. Do not apply to St. Augustinegrass, bermgrass, bermudagrass, myriophyllum, dionaea, kikuyagrass, bahiagrass or lippia.

APPLICATION INSTRUCTIONS

Apply on a calm day and lightly water to activate. Apply when temperatures are consistently between 60°F and 90°F. Avoid applying to adjacent flower beds when using a rotary spreader.

This product may cause temporary whitening of grass foliage. In general, symptoms appear five to seven days after application and may last up to 14 days. Discoloration is most likely to occur on ryegrass lawns.

WEEDS PREVENTED

Large crabgrass, Smooth crabgrass, Dandelion, Broadleaf plantain, Buckhorn plantain, Carpetweed, Common chickweed, Common lambsquarters, Common purslane, Corn speedwell, Ground ivy, Hairy bittercress, Henbit, Parsley plect, Pennycuaria smartweed, Persian speedwell, Pokeweed, Pursh speedwell, Redroot pigweed, Shepherd's-purse, Yellowweed, White clover, Yellow woodsoor, Flatsedge, Yellow nutedge

EPA Reg No. 538-317 (5M)

Scott's Step 1 for Seeding Starter Lawn Food with Weed Preventer

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run-off to water bodies or drainage systems.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place. Reseal bag by folding top down and securing.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

EPA Reg. No. 538-317 EPA Est. No. 538-011-1¹, 538-011-2¹, 85652-011-1¹
RCAS# 104206-82-8 Superscript used is first letter of run code on end of container.

3/28/12 SSL

Page 6 of 6

STEP 1 for Seeding Starter 21–22–4 F643
Lawn Food with Weed Preventer

Guaranteed Analysis

Total Nitrogen (N)	21%
4.6% Ammoniacal Nitrogen	
8.1% Urea Nitrogen	
6.0% Other Water Soluble Nitrogen*	
0.6% Water Insoluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	23%
Soluble Potash (K ₂ O)	4%
Sulfur (S)	1%
1.0% Combined Sulfur (S)	
Iron (Fe) Total	0.50%
0.50% Water Soluble Iron (Fe)	

Derived from: Methylenecreas, Urea, Potassium Sulfate, Ammonium Phosphate and Ferrous Sulfate.

*Contains 5.6% slowly available nitrogen from methylenecreas.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Questions / Information

SMG
Sulfur
Methyl
Iron
Sulfate
WISE
Sulfur
Methyl
Iron
Sulfate
Sulfur and Guaranteed by
The Scotts Company
14111 Scotts Lawn Rd.
Marysville, Ohio 43041

Consumer Helpline 1-800-543-TURF (8873)

©2012 The Scotts Company LLC.
World rights reserved.

EPA Reg No. 538-317 (5M)

Scotts Step 1 for Seeding Starter Lawn Food with Weed Preventer

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to your money back. Simply send us proof of purchase and we will promptly mail you a refund.

SCOTTS® LAWN FOOD-STEP 1 FOR SEEDING STARTERS®
LAWN FOOD WITH WEED PREVENTER 5M 36905

120% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

0 32247 36905 3
PROOF OF PURCHASE
VID-88
BG1801