

TRUSTED SINCE 1926
BONIDE[®]

DuraTurf[™]

LAWN WEED KILLER

Enjoy a beautiful weed free lawn

Controls over 200 listed weeds!
Including Dandelion, Chickweed, Knotweed



Ground Ivy



Henbit



Oxalis



Plantain



Purslane



Dandelion

Kills lawn weeds



Spotted spurge



Clover



Chickweed



Wild Onion



Thistle

Keep Out Of Reach Of Children

CAUTION

(See Back Panel for Additional Precautionary Statements)

ACTIVE INGREDIENTS:

| | |
|--|----------|
| 2,4-dichlorophenoxyacetic acid..... | 1.37% |
| (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid..... | 0.31% |
| Dicamba: 3,6-dichloro-o-anisic acid..... | 0.13% |
| OTHER INGREDIENTS:..... | 98.19% |
| TOTAL:..... | 100.000% |

THIS PRODUCT CONTAINS:

| |
|--|
| 27.4 lb 2,4-dichlorophenoxyacetic acid equivalent per ton. |
| 6.2 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton. |
| 2.6 lb 3,6-dichloro-o-anisic acid equivalent per ton. |

Isomer Specific by AOAC Methods
EPA Est. No. 32802-MO-1(B), 59144-GA-1(A), 37351-CO-1(C),
8378-IN-1(D), 9404-FL-1(E), 35497-OR-1(H) 49991-TX-1(Q), 4-NY-1
EPA Reg. No. 2217-640-4

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Apply when the air is calm. Do not apply to newly seeded lawns. Do not exceed specified dosages for any area. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

This product controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

For use on established turf of warm season grasses includes bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq.ft.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

DROP SPREADERS

| DROP SPREADERS | SETTING |
|------------------|---------|
| Republic EZ #530 | 3 |
| Scotts AccuGreen | 4 |
| Sears Drop | 4 |

ROTARY SPREADERS

| ROTARY SPREADERS | SETTING |
|--------------------|---------|
| Earthway | 7 |
| Republic EZ #540 | 2 |
| Scotts SpeedyGreen | 4 |
| Sears Rotary | 2 |

COVERAGE

To determine how much fertilizer you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 5,000 sq.ft. Length x Width = Total Sq.Ft. - Non-Lawn Areas = Total Sq.Ft. Coverage You Need

SPREADER SETTINGS

IMPORTANT: It is important to follow the directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

SPREADING HINTS

These granular settings are approximate. Calibrate your spreader on 100 sq.ft. and adjust to apply at the rate of 1 lb. of product per 500 sq.ft.. This bag covers 5,000 sq.ft. of lawn area.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

WEEDS

| | | | | | | | |
|---------------------------|---------------------|-------------------|---------------------|---------------------|------------------------|----------------------|---------------------|
| Annual yellow sweetclover | Bulbous buttercup | Common mullein | Hedge bindweed | Oriental cocklebur | Redstem filaree | Tansy ragwort | Wild radish |
| Aster | Bull thistle | Creeping jenny | Hedge mustard | Oxalis | Rough cinquefoil | Tansy mustard | Wild rape |
| Austrian fieldcress | Bullnettle | Cudweed | Hemp | Parsley-piert | Rough fleabane | Tanweed | Wild strawberry |
| Bedstraw | Burclover | Curly dock | Henbit | Parsnip | Russian pigweed | Thistle | Wild sweet potato |
| Beggarticks | Burdock | Daisy, English | Hoary cress | Pearlwort | Russian thistle | Trailing crownvetch | Wild vetch |
| Betony, Florida | Burning nettle | Daisy fleabane | Hoary plantain | Pennycress | Scarlet pimpernel | Tumble mustard | Woodsorrel |
| Bindweed, field | Burweed | Daisy, oxeye | Hoary vervain | Pepperweed | Scotch thistle | Tumble pigweed | Woolly croton |
| Bird vetch | Buttercup | Dandelion | Horsenettle | Pigweed | Sheep sorrel | Velvetleaf | Woolly morningglory |
| Bitter wintercress | Buttonweed | Dichondra | Jimsonweed | Pineywoods bedstraw | Shepherdspurse | Venice mallow | Woolly plantain |
| Bittercress, hairy | Canada thistle | Dogbane | Knawel | Plains coreopsis | Slender plantain | Virginia buttonweed | Wormseed |
| Bitterweed | Carolina geranium | Dogfennel | Knotweed | Plantain | Smallflower galinsoga | Virginia creeper | Yarrow |
| Black-eyed Susan | Carpetweed | Dollarweed | Kochia | Poison ivy | Smooth dock | Virginia pepperweed | Yellow rocket |
| Black medic | Catchweed bedstraw | False dandelion | Lambsquarters | Poison oak | Smooth pigweed | Wavyleaf bullthistle | Yellowflower |
| Black mustard | Catnip | Falseflax | Lespedeza | Pokeweed | Sorrel | Western clematis | pepperweed |
| Blackseed plantain | Catsear | False sunflower | Mallow | Poorjoe | Sowthistle | Western salsify | |
| Blessed thistle | Chickweed, common | Fiddleneck | Matchweed | Prairie sunflower | Spanishneedles | White mustard | |
| Bloodflower milkweed | Chickweed, mouseear | Florida pusley | Mexicanweed | Prickly lettuce | Speedwell | Wild aster | |
| Blue lettuce | Chicory | Frenchweed | Milk vetch | Prickly sida | Spiny amaranth | Wild buckwheat | |
| Blue vervain | Cinquefoil | Galinsoga | Morningglory | Prostrate knotweed | Spiny cocklebur | Wild carrot | |
| Bracted plantain | Clover, crimson | Goldenrod | Mouseear hawkweed | Prostrate pigweed | Spiny sowthistle | Wild four-o'clock | |
| Brassbuttons | Clover, hop | Ground ivy | Mugwort | Prostrate spurge | Spotted catsear | Wild garlic | |
| Bristly oxtongue | Clover, red | Gumweed | Musk thistle | Prostrate vervain | Spotted spurge | Wild geranium | |
| Broadleaf dock | Clover, strawberry | Hairy fleabane | Mustard | Puncturevine | Spurweed | Wild lettuce | |
| Broadleaf plantain | Clover, sweet | Hawkweed | Narrowleaf plantain | Purslane, common | Stinging nettle | Wild marigold | |
| Broomweed | Clover, white | Healall | Narrowleaf vetch | Ragweed | Strawberry, India mock | Wild mustard | |
| Buckhorn | Cockle | Heartleaf drymary | Nettle | Red sorrel | Tall nettle | Wild onion | |
| Buckhorn plantain | Cocklebur | Heath aster | Orange hawkweed | Redroot pigweed | Tall vervain | Wild parsnip | |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

First Aid

| | |
|-----------------------------------|--|
| If on skin or on clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none"> • Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.