

BIMART OWNER'S CHOICE™ LAWN WEED KILLER READY TO USE

- Kills Dandelion, Clover, Chickweed, Oxalis and over 200 other broadleaf weeds to the root!
- · Use on all types of Lawns won't harm lawn grasses when used as directed
- Starts working over night

STOPI READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Store and transport in an upright position.

EPA Reg. No. 4-401 EPA Est. No. 4-NY-1

Buyers guarantee limited to label claims. @Bonide Products, Inc. All Rights Reserved

Bonide Products, Inc.



WHAT IT IS & HOW TO USE IT

This product contains Trimec® turf herbicide, a combination of three proven weed killers 2,4-D, mecoprop and dicamba. Together they control a wide range of broadleaf lawn weeds; some of which are listed on this label.

Use to "spot treat" weeds in the lawn. For best results, apply when weeds are actively growing. Do not water within 24 hours after spraying. Use as is. Do not mix. Do not dilute. Aim at weed and spray. Only a short burst is needed. Overspraying only wastes the product and may cause damage to grasses such as centipedegrass, St. Augustinegrass, and Floratam. Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 30 days if necessary. The number of spot treatments is limited to 2 per year.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not exceed specified dosage. This product is intended for spot
- spraying individual weeds only.

 Do not apply as a fine mist because of potential for injury to
- desirable plants,

 Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when spiriting as when rein is expected that day. when raining or when rain is expected that day.

CONTROLS - WEEDS			
Annual yellow sweetclover Aster Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field	Bulbous buttercup Bull thistle Bull nettle Burclover Burdock Burning nettle Burweed Buttercup	Clover, white Cockle Cocklebur Common mullein Creeping charlie Creeping jenny Cudweed Curly dock	Heartleaf drymary Heath aster Hedge bindweed Hedge mustard Henbit Hoary cress Hoary plantain Hoary vervain Horsenettle
Bird vetch Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower	Buttonweed Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catnip Catsear Chickweed,	Daisy, English Daisy fleabane Daisy, oxeye Dandelion Dichondra Doglennel Dollarweed False dandelion Falseflax False sunflower	Jimsonweed Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed
milkweed Blue lettuce Blue vervain Bracted plantain Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain	common Chickweed, mouseear Chicary Cinquefoil Clover, crimson Clover, hop Clover, red Clover, strawberry Clover, strawberry	Fiddleneck Florida pusley Frenchweed Galinsoga Goldenrod Ground ivy Gurmweed Hairy fleabane Hawkweed Healall	Milk vetch Morringglory Mouseear hawkweed Mugwort Musk thistle Mustard Narrowleaf plantain Narrowleaf yetch

CONTROLS - WEEDS Cont. Nettle Orange hawkweed Oriental cocklebur Oxalis Parsley-piert Parsnip Pearlwort Stinging nettle Strawberry, India Wild carrot Wild four-o'clock Wild garlic Red sorrel Redroot pigweed mock Redstern filaree Rough cinquefoil fall nettle Wild geranium Wild lettuce Tall vervain Wild marigold Wild mustard Tansy ragwort Tansy mustard Rough fleabane Russian pigweed Russian thistle Pennycress Pepperweed Pigweed Pineywoods Tanweed Wild onion Scarlet pimpernel Scotch thistle Thistle Wild parsnip Trailing crownvetch Wild radish Wild rape Wild strawberry Sheep sorrel bedstraw Plains coreopsis Plantain Shepherdspurse Tumble mustard Wild sweet potato Wild vetch Tumble pigweed Slender plantain Smallflower Velvetleaf galinsoga Smooth dock Venice mallow Woodsorrel Woolly croton Poison ivy Virginia Poison oak Pokeweed Poorjoe Prairie sunflower Prickly lettuce Woolly morningglory Woolly plantain Wormseed Smooth pigweed buttonweed Virginia creeper Sorrel Virginia
pepperweed
Wavyleaf
bullthistle Sowthistle panishneedles Prickly sida Prostrate knotwee needwell Yarrow piny amaranth Yellow rocket Western clematis Prostrate pigweed Prostrate spurge Spiny cocklebur Spiny sowthistle Yellowflower Western salsify White mustard pepperweed Yellow sweetclove Prostrate vervain Spotted catsear potted spurge Wild aster and other broad-leaf weeds uncturevine Wild buckwheat Purslane, common Spurweed

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION - Harmful if absorbed through skin. Harmful if
swallowed. Avoid contact with skin, eyes, or clothing. Causes
moderate eye irritation. Wash thoroughly with soap and water
after handling and before eating, drinking, chewing gum, using
tobacco, or using the toilet. Wear long-sleeved shirt, long pants,
socks, shoes, and gloves. Remove and wash contaminated
clothing before reuse.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and pets.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container.
Do not reuse or refill this container. If Empty - Do not reuse this container.
Place in trash or offer for recycling if available. If Partiy Filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

If in eyes:

- Hold eye open and rinse stowly and gently with water for 15 - 20 minutes.

- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

- Call a poison control center or doctor for treatment advice.

If swallowed:

- Have person sip a glass of water if able to swallow.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor.

- Do not induce vomiting unless told to do so by a poison control center or doctor or an unconscious person.

If on skin

- Take off contaminated clothing.

- Rinse skin immediately with plenty of water for 15 - 20 minutes.

- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticle Information Center at 1-800-858-7378 for emergency medical treatment information.

LIMITED WARRANTY AND DISCLAIMER -

For use only as directed. To the extent consistent with applicable law, the manufacturer neither makes nor intends any express or implied warranties, including any warranty of merchantability or fitness for a particular purpose, which are hereby expressly disclaimed. To the extent consistent with applicable law, in no case shall the manufacturer be liable for incidental, consequential, or special damages resulting from the use or handling of this product. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

For information on pesticide products (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.