10001513324-01 | Released - 05/08/2012 by , Connie Christian| Printed on 05/08/2012 - Printed By Con

Scotts®

Turf Builder®
WEED & FEED I

100% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only
32247 24990 4

PROOF OF PURCHASE
UPC 0-32247-24990-4

24990
1

4/27/12 SSL
Page 1 of 6

For Best Results Apply to a Wet Lawn

KILLS WEEDS DOWN TO THE ROOTS

Scotts®

Turf Builder®

WEED & FEED I

Fertilizante para césped con control de malezas

KILLS DANDELIONS AND FEEDS GRASS

TO CROWD OUT FUTURE WEEDS

Builds Strong, Deep Grass Roots

COVERS 5,000 SQ FT

NET WT./PESO NETO
14.29 lb (6.48 kg)

ACTIVE INGREDIENTS:	
2,4-D[1][2]	1.21%
Mecoprop-p[1][3][4]	0.61%
OTHER INGREDIENTS	98.18%
TOTAL	100.00%

[2]0.121 lbs 2,4-D acid equivalent per pound of product
[3]0.061 lbs MCPP-p acid equivalent per pound of product

Do not use on St Augustinegrass, Floratam, Dichondra, Lippia, Carpetgrass, or Bentgrass Lawns

KEEP OUT OF REACH OF CHILDREN
CAUTION SEE BACK FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

No aplique en césped San Agustín, floratam, oreja de ratón, lippia, césped alfombra ni agrostis.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION CONSULTE EL PANEL POSTERIOR PARA CONOCER LOS AVISOS DE PRECAUCION ADICIONALES

Scotts®

Turf Builder®
WEED & FEED I

28-0-3

EPA Reg No. 538-282 (5M)
Scotts Turf Builder Weed and Feed

10001513324_901_000_0100p70 r3pep2z CCHRISTIAN 2012-05-08T11:39:35 0.945

Guaranteed!

KILLS DANDELIONS AND FEEDS GRASS

TO CROWD OUT FUTURE WEEDS

Scotts®

DIRECTIONS FOR USE
INSTRUCCIONES DE USO

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This package applies 3.2 pounds of product per 1,000 sq. ft. This product applies 0.84 pounds of 2,4-D acid equivalent, 0.17 pounds of MCPP-p acid equivalent and 0.10 pounds of Dichlorprop-p acid equivalent per acre per application.

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta. Este producto aplica 1.4 kg de producto por 92.9 m2. Este producto aplica 0.38 kg de 2,4-D ácido equivalente, 77 g de MCPP-p ácido equivalente y 45 g de Dichlorprop-p ácido equivalente por acre por aplicación.

Scotts® QUICK START GUIDE: You're Just Three Simple Steps Away From A Beautiful Scotts® Lawn

Guía de inicio rápido de Scotts®: ¡Está a sólo tres sencillos pasos de tener un hermoso césped con Scotts®!

FILL & SET SPREADER
Llenar y ajustar el esparcidor

Scotts® Broadcast / Rotary Spreaders **4¾**

Scotts® Drop Spreaders **8¼**

Scotts® products are designed to work with Scotts spreaders. For additional spreader settings, please call 1-800-543-8873 This package applies 3.2 lbs of product per 1,000 sq. ft.

Los productos Scotts® están diseñados para funcionar con los esparcidores Scotts. Para ajustes adicionales del esparcidor, llame al 1-800-543-8873. Con este paquete, se aplican 1.3 kg de producto por cada 93 m².

APPLY
Aplicar

- Apply on a calm day.
- Apply to moist grass.
- Make sure that the grass is wet from dew or sprinkling before applying. This helps particles adhere to weed leaves to produce best results.
- Apply when weeds are actively growing (when daytime temperatures are consistently between 60°F and 90°F).
- Tall grass should be mowed a day or two before application.
- For even application walk at a steady pace slightly overlapping each pass.
- Scotts® Drop Spreaders: Align hopper arrow with previous wheel track.
- Scotts Broadcast/Rotary Spreaders: Line up each pass approximately 5 ft. from wheel track.

• Aplíquelo un día que no haya brisa. • Aplíquelo sobre el césped humedo. • Asegúrese de que el césped está húmedo, de lo contrario, rocíelo antes de aplicar el producto. Esto ayuda a que las particulas se adhieran a las hojas para obtener los mejores resultados. • Aplíquelo cuando las malezas estén creciendo activamente (cuando las temperaturas diurnas estén constantemente entre los 15,5 °C y los 32 °C). • Se debe cortar el césped alto uno o dos días antes de la aplicación. • Para lograr una aplicación uniforme, camine a un ritmo regular superponiendo ligeramente cada pasada. • Esparcidores de goteo Scotts®: Alinee la flecha de la tolva con la huella anterior de la rueda. • Esparcidores a voleo/giratorios Scotts®: Alinee cada pasada, aproximadamente, a 1,5 m de la huella de la rueda.

CLEAN-UP
Limpiar

Sweep product from hard surfaces onto the lawn to keep valuable feeding nutrients on the grass.

Barra el producto derramado en las superficies duras hacia el césped para mantener los nutrientes valiosos en el jardín.

YOU'RE DONE! WHAT'S NEXT?
¡Listo! ¿Qué sigue?

Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.

No permita que el producto entre en contacto con personas o mascotas, de forma directa ni a través de la desviación. Mantenga a las personas y a las mascotas lejos del césped durante la aplicación. No permita que las personas o las mascotas ingresen al área tratada hasta que el polvillo del producto se haya asentado.

Questions? Call **1-800-543-TURF**

¿Preguntas? Llame al 1-800-543-TURF

www.scotts.com

RESTRICTIONS

- Do not apply more than twice per year.
- A minimum of 30 days between applications is required.
- Apply only to grass. Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetable plants.
- Do not apply on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass. On Bermudagrass, bahia, centipedegrass, zoysia lawns, under certain conditions, some temporary discoloration may result.
- On bahia and centipede lawns, wait until after the second mowing of the year but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

Watering or Rain

- Do not water your lawn for 24 hours after application or apply if rain is expected within the next 24 hours.

Seeding/Sodding/Sprigging

- Delay application on newly seeded, sodded, or sprigged areas until new grass has been mowed 4 times.
- Wait at least one month after using this product to seed, sod or plant grass sprigs.

RESTRICCIONES

- No aplique el producto más de dos veces por año.
- Espere, al menos, 30 días antes de realizar una nueva aplicación.
- Sólo aplique al césped. No permita que los gránulos o el polvillo entren en contacto con las flores, arbustos, plantas frutales o vegetales.
- No aplique en césped San Agustín, floratam, oreja de ratón, lippia, césped alfombra ni agrostis. En céspedes Bermuda, Bahía, ciempiés y zoysia, en determinadas circunstancias, puede producirse una decoloración temporal.
- En césped Bahía y césped ciempiés, espere al segundo corte del año, pero NO antes de mediados de marzo, En el otoño, NO trate el césped ciempiés después del 1 de octubre y NO trate el césped Bahía después del 1 de noviembre.
- No aplique el producto directamente sobre agua ni cerca de lugares donde haya agua, como drenajes de tormenta, canalones, alcantarillas o zanjas de drenaje. No aplique el producto a menos de 8 m de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías y océanos.
- No aplique el producto cuando hay viento. Aplique este producto directamente sobre su césped, y barra el producto que caiga sobre la entrada de automóviles, la acera o la calle hacia el área tratada. Para evitar el escurrimiento del producto, no riegue el área tratada en exceso, ni lo aplique en condiciones de lluvia o en días con pronóstico de lluvia.

Riego o lluvia

- No riegue su césped durante 24 horas después de la aplicación ni aplique si se prevén lluvias en el plazo de las próximas 24 horas.

Sembrar/Cubrir de césped/Plantar retoños

- Retrase la aplicación en áreas recientemente sembradas, cubiertas de césped o donde se plantaron retoños hasta que haya cortado el césped cuatro veces.
- Espere, al menos, un mes después de utilizar este producto para sembrar, cubrir de césped o plantar retoños.

WEED CONTROL

- Kills listed weeds.
- Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control.

Aster	Burclover	Chicory	Curly Dock	Fleabane	Lambsquarters	Pennywort	Red Sorrel (Sheep Sorrel)	Wild Carrot
Bindweed	Buttonweed	Cinquefoil	Dandelion	Ground Ivy	Lespedeza	Pepperweed	Scarlet Pimpernel	Yarrow
Bittercress	Canada Thistle	Clover (Hop)	Dichondra	Hawkweed	Lippia (Matchweed)	Pigweed	Shepherd's-purse	Yellow Rocket
Black Medic	Carpetweed	Clover (Red)	Dollarweed	Healall	Mallow	Plantain	Sow Thistle	
Buckhorn	Catsear	Clover (White)	English Daisy	Henbit	Morningglory	Pusley	Spurweed	
Bull Thistle	Chickweed	Cudweed	Filaree	Knotweed	Mustard	Ragweed	Vetch	

TIPS FOR LAWNCARE SUCCESS

Feed Regularly

Regular feedings provide the nutrients your lawn needs to look its best and withstand the stresses of weeds, heat, drought & family activity.

Benefits of Multiple Feedings

1 ANNUAL FEEDING | 2-3 ANNUAL FEEDINGS | 4 ANNUAL FEEDINGS

Mow and Leave the Grass Clippings on Your Lawn

Mowing high allows the grass to develop a deep root system & helps your lawn use water more effectively. Grass clippings recycle the plant nutrients back into the soil.

Conserve Water

Well-fed lawns are more efficient at using available water. Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat drought and other stresses. Remember your lawn will begin to wilt when water is needed. As much as possible take advantage of nature's sprinkler and rely on the rain to water your lawn.

SCOTTS® RECOMMENDED FEEDING ROUTINE

EARLY SPRING
An early spring feeding helps grass green-up after winter. You should feed around the first time you mow your grass.

LATE SPRING
Feeding now prepares grass for summer stress by thickening the blades on top and building deep roots below.

EARLY FALL
An early fall feeding replenishes the nutrients lost in the harsh summer months for quicker recovery. You should feed when your grass comes out of dormancy.

LATE FALL
Late fall is the best feeding time because it builds strong deep roots and prepares grass for winter. You should feed when all the leaves have fallen.

Florida Annual Fertilizer Applications: 4 annual feedings with the exception of bahiagrass and centipedegrass (2 feedings in the north and 3 feedings in the central and south).

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

Scotts® & Your Environment

We care. Scotts® products are designed with care to grow thicker, greener lawns. **A Green Lawn Is a Good Lawn -** A healthy lawn can clean the air, produce oxygen and prevent runoff and soil erosion. **You can make a difference -** Help keep our water resources clean. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk or street back onto your lawn.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment. Call: 1-800-543-TURF (8873).

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.
This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry area. Always store pesticides in the original container. Store away from food and pet food. Reseal opened bag by folding top down and securing.
PESTICIDE CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Scotts®Turf Builder® Weed & Feed B 28-0-3 F643

GUARANTEED ANALYSIS

Total Nitrogen (N)	28%
28% Urea Nitrogen*	
Soluble Potash (K_2O)	3%

Derived from: Urea, Polymer coated Urea, Potassium Chloride.
*7.5% slowly available Nitrogen from coated Urea.

EPA Reg. No. 538-314
EPA Est. No. 538-OH-1[1], 85653-OH-1[1], 9198-OH-2[2], 9198-OH-1[4], 9198-AE-001[1], 084175-TX-001[1], 034637-OH-002[2], 034637-OH-007[1], 034637-OH-003[2], 034637-OH-006[1], 538-SC-1[1], 35512-FL-2[2], 538-OH-2[2], 82757-FL-001[1], 82757-OH-001[4], 82757-MA-002[2], 35497-OR-1[1],
Superscript used is first letter of run code on end of container.

QUESTIONS / INFORMATION

Preguntas/Información

Sold and Guaranteed By:
The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

Questions? Call **1-800-543-8873**

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-sc.com

La información relacionada con los contenidos y los niveles de metales de este producto está disponible en Internet en http://www.regulatory-info-sc.com

©2013 The Scotts Company LLC
World rights reserved.

AVISOS DE PRECAUCION

Peligroso para humanos y animales domésticos

PRECAUCION: Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Use camisa de mangas largas, pantalones largos, zapatos y calcetines. En algunas personas, el contacto prolongado o frecuente con la piel puede provocar reacciones alérgicas.

PRIMEROS AUXILIOS

SI ENTRA EN CONTACTO CON LOS OJOS: Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos. Si usa lentes de contacto, retírelos después de los primeros 5 minutos y luego continúe enjuagando los ojos.
Llame a un centro de control de envenenamiento o a un médico para recibir instrucciones de tratamiento. Cuando llame a un centro de control de envenenamiento o a un médico, o intente obtener tratamiento, tenga a mano el envase o la etiqueta del producto. Llame al 1-800-543-TURF (8873).

RIESGOS AMBIENTALES: Este pesticida es tóxico para los peces y otros invertebrados acuáticos, y puede dañar las plantas no indicadas para el tratamiento. El desplazamiento y el escurrimiento del producto pueden ser peligrosos para los organismos acuáticos en aguas adyacentes a las áreas tratadas. Para proteger el medio ambiente, no permita que el pesticida ingrese o discurra a desagües de tormenta, zanjas de drenaje, canalones o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no haya un pronóstico de lluvia para las siguientes 24 horas, asegurará que el viento o la lluvia no arrastren o laven el pesticida del área de tratamiento. Barra el producto que caiga sobre la entrada de automóviles, la vereda o la calle hacia el área tratada del césped para evitar que discurra a cuerpos de agua o sistemas de drenaje.
Este producto químico tiene propiedades y características asociadas con las sustancias químicas detectadas en aguas subterráneas. La utilización de este producto químico en áreas donde los suelos son permeables, en particular donde el manto freático es superficial, puede derivar en la contaminación de las aguas subterráneas. La aplicación en torno a una cisterna o a un pozo puede provocar la contaminación del agua potable o las aguas subterráneas.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO DE PESTICIDAS: Guarde el producto en un lugar seco. Almacene el pesticida en su envase original en un lugar seguro, lejos de niños y mascotas.
ELIMINACION Y MANEJO DEL ENVASE DEL PESTICIDA: Envase no rellenable. No vuelva a utilizar ni a llenar este envase. Si está vacío: Colóquelo en la basura o recíclelo, si corresponde. Si está parcialmente lleno: Llame a su agencia de desechos sólidos local para obtener instrucciones sobre cómo eliminar el producto. Nunca elimine el producto no utilizado en drenajes internos o externos.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

NOTA: En conformidad con la ley aplicable, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones correspondientes.

SCOTTS NO-QUIBBLE GUARANTEE®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund promptly.

Si por alguna razón usted, el consumidor, no está completamente satisfecho con este producto, le devolveremos su dinero. Sólo deberá enviarnos el comprobante de compra y le enviaremos un cheque de reembolso por correo inmediatamente.

PROOF OF PURCHASE UPC 0-32247-25000-9
25000-1 VID-BB BG2290

115% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

0 32247 25000 9

SCOTTS® TURF BUILDER® WEED & FEED I
5M 25000-1

100% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

32247 25000 9

PROOF OF PURCHASE
UPC 0-32247-25000-9

25000
1

NO VITAL COPY ON THE B
BOTTOM SEAL IS AT 1/4" +/- 1/8"

EPA Reg. No. 538-314
Scotts Turf Builder Weed & FeedB

KILLS WEEDS DOWN TO THE ROOTS

For Best Results Apply to a Wet Lawn

Scotts®

Turf Builder®

WEED & FEED B

Fertilizante para césped con control de malezas

EPA Reg. No. 538-314
Scotts Turf Builder Weed & FeedB

KILLS DANDELIONS AND FEEDS GRASS

TO CROWD OUT FUTURE WEEDS

Builds Strong, Deep Grass Roots

COVERS 5,000 SQ FT

NET WT./PESO NETO
16.00 lb (7.25 kg)

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt ‡	0.723%
Mecoprop-p, dimethylamine salt ‡‡	0.148%
Dichlorprop-P, dimethylamine salt ‡‡‡	0.088%
OTHER INGREDIENTS	99.041%
TOTAL	100.000%

This product contains:
‡ 0.601% 2,4-dichlorophenoxyacetic acid equivalent
‡‡ 0.122% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent
‡‡‡ 0.074% (+)-(R)-2-(2,4-dichlorophenoxy) propionic acid equivalent

Do not use on St. Augustinegrass including floratam, dichondra, lippia, carpetgrass or bentgrass lawns.

KEEP OUT OF REACH OF CHILDREN

CAUTION SEE BACK FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

No aplique en césped San Agustín, floratam, oreja de ratón, lippia, césped alfombra ni agrostis.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCION CONSULTE EL PANEL POSTERIOR PARA CONOCER LOS AVISOS DE PRECAUCION ADICIONALES.

Scotts®

Turf Builder®

WEED & FEED$_B$

28-0-3

NO VITAL COPY ON THE BOTTOM 1/2"
BOTTOM SEAL IS AT 1/4" +/- 1/8" INK IS ALLOWED

EPA Reg. No. 538-314
Scotts Turf Builder Weed & FeedB

Scotts®

DIRECTIONS FOR USE
INSTRUCCIONES DE USO

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This package applies 3.2 pounds of product per 1,000 sq. ft. This product applies 0.84 pounds of 2,4-D acid equivalent, 0.17 pounds of MCPP-p acid equivalent and 0.10 pounds of Dichlorprop-p acid equivalent per acre per application.

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta. Este producto aplica 1.4 kg de producto por 92.9 m2. Este producto aplica 0.38 kg de 2,4-D ácido equivalente, 77 g de MCPP-p ácido equivalente y 45 g de Dichlorprop-p ácido equivalente por acre por aplicación.

Scotts® QUICK START GUIDE: You're Just Three Simple Steps Away From A Beautiful Scotts® Lawn

Guía de inicio rápido de Scotts®: ¡Está a sólo tres sencillos pasos de tener un hermoso césped con Scotts®!

FILL & SET SPREADER
Llenar y ajustar el esparcidor

Scotts® Broadcast / Rotary Spreaders	4¾
Scotts® Drop Spreaders	8¼

Scotts® products are designed to work with Scotts spreaders. For additional spreader settings, please call 1-800-543-8873 This package applies 3.2 lbs of product per 1,000 sq. ft.

Los productos Scotts® están diseñados para funcionar con los esparcidores Scotts. Para ajustes adicionales del esparcidor, llame al 1-800-543-8873. Con este paquete, se aplican 1.3 kg de producto por cada 93 m².

APPLY
Aplicar

- Apply on a calm day.
- Apply to moist grass.
- Make sure that the grass is wet from dew or sprinkling before applying. This helps particles adhere to weed leaves to produce best results.
- Apply when weeds are actively growing (when daytime temperatures are consistently between 60°F and 90°F).
- Tall grass should be mowed a day or two before application.
- For even application walk at a steady pace slightly overlapping each pass.
- Scotts® Drop Spreaders: Align hopper arrow with previous wheel track.
- Scotts Broadcast/Rotary Spreaders: Line up each pass approximately 5 ft. from wheel track.

• Aplíquelo un día que no haya brisa. • Aplíquelo sobre el césped húmedo. • Asegúrese de que el césped esté húmedo, de lo contrario, rocíelo antes de aplicar el producto. Esto ayuda a que las partículas se adhieran a las hojas para obtener los mejores resultados. • Aplíquelo cuando las malezas estén creciendo activamente (cuando las temperaturas diurnas estén constantemente entre los 15.5 °C y los 32 °C). • Se debe cortar el césped alto uno o dos días antes de la aplicación. • Para lograr una aplicación uniforme, camine a un ritmo regular superponiendo ligeramente cada pasada. • Esparcidores de goteo Scotts®: Alinee la flecha de la tolva con la huella anterior de la rueda. • Esparcidores a voleo/giratorios Scotts®: Alinee cada pasada, aproximadamente, a 1.5 m de la huella de la rueda.

CLEAN-UP
Limpiar

Sweep product from hard surfaces onto the lawn to keep valuable feeding nutrients on the grass.

Barra el producto derramado en las superficies duras hacia el césped para mantener los nutrientes valiosos en el jardín.

YOU'RE DONE! WHAT'S NEXT?
¡Listo! ¿Qué sigue?

Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.

No permita que el producto entre en contacto con personas o mascotas, de forma directa ni a través de la desviación. Mantenga a las personas y a las mascotas lejos del césped durante la aplicación. No permita que las personas o las mascotas ingresen al área tratada hasta que el polvillo del producto se haya asentado.

Questions? Call **1-800-543-TURF**

¿Preguntas? Llame al 1-800-543-TURF

www.scotts.com

RESTRICTIONS

- Do not apply more than twice per year.
- A minimum of 30 days between applications is required.
- Apply only to grass. Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetable plants.
- Do not apply on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass. On Bermudagrass, bahia, centipedegrass, zoysia lawns, under certain conditions, some temporary discoloration may result.
- On bahia and centipede lawns, wait until after the second mowing of the year but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat bahia after the first of November.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

Watering or Rain

- Do not water your lawn for 24 hours after application or apply if rain is expected within the next 24 hours.

Seeding/Sodding/Sprigging

- Delay application on newly seeded, sodded, or sprigged areas until new grass has been mowed 4 times.
- Wait at least one month after using this product to seed, sod or plant grass sprigs.

RESTRICCIONES

- No aplique el producto más de dos veces por año.
- Espere, al menos, 30 días antes de realizar una nueva aplicación.
- Sólo aplique al césped. No permita que los gránulos o el polvillo entren en contacto con las flores, arbustos, plantas frutales o vegetales.
- No aplique en césped San Agustín, floratam, oreja de ratón, lippia, césped alfombra ni agrostis. En céspedes Bermuda, Bahía, ciempiés y zoysia, en determinadas circunstancias, puede producirse una decoloración temporal.
- En césped Bahía y césped ciempiés, espere al segundo corte del año, pero NO antes de mediados de marzo. En el otoño, NO trate el césped ciempiés después del 1 de octubre y NO trate el césped Bahía después del 1 de noviembre.
- No aplique el producto directamente sobre agua ni cerca de lugares donde haya agua, como drenajes de tormenta, canalones, alcantarillas o zanjas de drenaje. No aplique el producto a menos de 8 m de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías y océanos.
- No aplique el producto cuando hay viento. Aplique este producto directamente sobre su césped, y barra el producto que caiga sobre la entrada de automóviles, la acera o la calle hacia el área tratada. Para evitar el escurrimiento del producto, no riegue el área tratada en exceso, ni lo aplique en condiciones de lluvia o en días con pronóstico de lluvia.

Riego o lluvia

- No riegue su césped durante 24 horas después de la aplicación ni aplique si se prevén lluvias en el plazo de las próximas 24 horas.

Sembrar/Cubrir de césped/Plantar retoños

- Retrase la aplicación en áreas recientemente sembradas, cubiertas de césped o donde se plantaron retoños hasta que haya cortado el césped cuatro veces.
- Espere, al menos, un mes después de utilizar este producto para sembrar, cubrir de césped o plantar retoños.

WEED CONTROL

- Kills listed weeds.
- Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control.

Aster	Burclover	Chicory	Curly Dock	Fleabane	Lambsquarters	Pennywort	Red Sorrel (Sheep Sorrel)	Wild Carrot
Bindweed	Buttonweed	Cinquefoil	Dandelion	Ground Ivy	Lespedeza	Pepperweed	Scarlet Pimpernel	Yarrow
Bittercress	Canada Thistle	Clover (Hop)	Dichondra	Hawkweed	Lippia (Matchweed)	Pigweed	Shepherd's-purse	Yellow Rocket
Black Medic	Carpetweed	Clover (Red)	Dollarweed	Healall	Mallow	Plantain	Sow Thistle	
Buckhorn	Catsear	Clover (White)	English Daisy	Henbit	Morningglory	Pusley	Spurweed	
Bull Thistle	Chickweed	Cudweed	Filaree	Knotweed	Mustard	Ragweed	Vetch	

TIPS FOR LAWNCARE SUCCESS

Feed Regularly

Regular feedings provide the nutrients your lawn needs to look its best and withstand the stresses of weeds, heat, drought & family activity.

Benefits of Multiple Feedings

1 ANNUAL FEEDING | 2-3 ANNUAL FEEDINGS | 4 ANNUAL FEEDINGS

Mow and Leave the Grass Clippings on Your Lawn

Mowing high allows the grass to develop a deep root system & helps your lawn use water more effectively. Grass clippings recycle the plant nutrients back into the soil.

Conserve Water

Well-fed lawns are more efficient at using available water. Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat drought and other stresses. Remember your lawn will begin to wilt when water is needed. As much as possible take advantage of nature's sprinkler and rely on the rain to water your lawn.

SCOTTS® RECOMMENDED FEEDING ROUTINE

EARLY SPRING
An early spring feeding helps grass green-up after winter. You should feed around the first time you mow your grass.

LATE SPRING
Feeding now prepares grass for summer stress by thickening the blades on top and building deep roots below.

EARLY FALL
An early fall feeding replenishes the nutrients lost in the harsh summer months for quicker recovery. You should feed when your grass comes out of dormancy.

LATE FALL
Late fall is the best feeding time because it builds strong deep roots and prepares grass for winter. You should feed when all the leaves have fallen.

Florida Annual Fertilizer Applications: 4 annual feedings with the exception of bahiagrass and centipedegrass (2 feedings in the north and 3 feedings in the central and south).

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

Scotts & Your Environment

We care. Scotts® products are designed with care to grow thicke[illegible] greener lawns. **A Green Lawn Is a Good Lawn -** A healthy lawn can clean the a[illegible] and prevent runoff and soil erosion. **You can make a difference** water resources clean. Apply this product only to your lawn, and sweep any

EPA Reg. No. 538-314
Scotts Turf Builder Weed & FeedB

wilt when water is needed. As much as possible take advantage of nature's sprinkler and rely on the rain to water your lawn.

Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

Scotts® & Your Environment

We care. Scotts® products are designed with care to grow thicker, greener lawns. **A Green Lawn Is a Good Lawn -** A healthy lawn can clean the air, produce oxygen and prevent runoff and soil erosion. **You can make a difference -** Help keep our water resources clean. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk or street back onto your lawn.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment. Call: 1-800-543-TURF (8873).

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.
This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry area. Always store pesticides in the original container. Store away from food and pet food. Reseal opened bag by folding top down and securing.
PESTICIDE CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Scotts®Turf Builder® Weed & Feed B 28-0-3 F643

GUARANTEED ANALYSIS

Total Nitrogen (N).....28%
28% Urea Nitrogen*
Soluble Potash (K_2O).....3%

Derived from: Urea, Polymer coated Urea, Potassium Chloride.
*7.5% slowly available Nitrogen from coated Urea.

EPA Reg. No. 538-314
EPA Est. No. 538-OH-1[S], 85653-OH-1[R], 9198-OH-2[Q], 9198-OH-1[M], 9198-AL-001[Y], 084175-TX-001[B], 034637-OH-002[T], 034637-OH-007[I], 034637-OH-003[Q], 034637-OH-006[X], 538-SC-1[P], 35512-FL-2[H], 538-OH-2[V], 82757-FL-001[T], 82757-OH-001[K], 82757-MA-002[E], 35497-OR-1[W].
Superscript used is first letter of run code on end of container.

QUESTIONS / INFORMATION

Preguntas/Información

SMG Listed NYSE

Sold and Guaranteed By:
The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

Questions? Call
1-800-543-8873

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-sc.com

La información relacionada con los contenidos y los niveles de metales de este producto está disponible en Internet en http://www.regulatory-info-sc.com

©2013 The Scotts Company LLC
World rights reserved.

AVISOS DE PRECAUCION

Peligroso para humanos y animales domésticos

PRECAUCION: Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Use camisa de mangas largas, pantalones largos, zapatos y calcetines. En algunas personas, el contacto prolongado o frecuente con la piel puede provocar reacciones alérgicas.

PRIMEROS AUXILIOS

SI ENTRA EN CONTACTO CON LOS OJOS: Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos. Si usa lentes de contacto, retírelos después de los primeros 5 minutos y luego continúe enjuagando los ojos.
Llame a un centro de control de envenenamiento o a un médico para recibir instrucciones de tratamiento. Cuando llame a un centro de control de envenenamiento o a un médico, o intente obtener tratamiento, tenga a mano el envase o la etiqueta del producto. Llame al 1-800-543-TURF (8873).

RIESGOS AMBIENTALES: Este pesticida es tóxico para los peces y otros invertebrados acuáticos, y puede dañar las plantas no indicadas para el tratamiento. El desplazamiento y el escurrimiento del producto pueden ser peligrosos para los organismos acuáticos en aguas adyacentes a las áreas tratadas. Para proteger el medio ambiente, no permita que el pesticida ingrese o discurra a desagües de tormenta, zanjas de drenaje, canalones o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no haya un pronóstico de lluvia para las siguientes 24 horas, asegurará que el viento o la lluvia no arrastren o laven el pesticida del área de tratamiento. Barra el producto que caiga sobre la entrada de automóviles, la vereda o la calle hacia el área tratada del césped para evitar que discurra a cuerpos de agua o sistemas de drenaje.
Este producto químico tiene propiedades y características asociadas con las sustancias químicas detectadas en aguas subterráneas. La utilización de este producto químico en áreas donde los suelos son permeables, en particular donde el manto freático es superficial, puede derivar en la contaminación de las aguas subterráneas. La aplicación en torno a una cisterna o a un pozo puede provocar la contaminación del agua potable o las aguas subterráneas.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO DE PESTICIDAS: Guarde el producto en un lugar seco. Almacene el pesticida en su envase original en un lugar seguro, lejos de niños y mascotas.
ELIMINACION Y MANEJO DEL ENVASE DEL PESTICIDA: Envase no rellenable. No vuelva a utilizar ni a llenar este envase. **Si está vacío:** Colóquelo en la basura o recíclelo, si corresponde. **Si está parcialmente lleno:** Llame a su agencia de desechos sólidos local para obtener instrucciones sobre cómo eliminar el producto. Nunca elimine el producto no utilizado en drenajes internos o externos.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

NOTA: En conformidad con la ley aplicable, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones correspondientes.

SCOTTS NO-QUIBBLE GUARANTEE®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund promptly.

Si por alguna razón usted, el consumidor, no está completamente satisfecho con este producto, le devolveremos su dinero. Sólo deberá enviarnos el comprobante de compra y le enviaremos un cheque de reembolso por correo inmediatamente.

PROOF OF PURCHASE UPC 0-32247-25000-9
25000-1 VID-BB BG2290

115% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

0 32247 25000 9

SCOTTS® TURF BUILDER®
WEED & FEED I
5M 25000-1

12/5/2012 SSL
Page 6 of 6

NO VITAL COPY ON THE BOTTOM 1/2"

EPA Reg. No. 538-314
Scotts Turf Builder Weed & FeedB