



538-283_Scotts LawnPro Super Turf Builder Winterguard Fall Weed And Feed 1 27-0-12_20170109_145_538_.pdf

LawnPro

Water Smart

SCOTT'S BEST FERTILIZER!

80% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

29206

1



Scott's

SUPER Turf Builder®


**WINTERGUARD®
FALL WEED AND FEED!**

SUPER Turf Builder®

**WINTERGUARD®
FALL WEED AND FEED!**



Kills dandelions, chickweed, white clover and other broadleaf weeds



Protects and Strengthens for a Better Spring Lawn



Improves Lawn's Ability to Absorb Water & Nutrients

DO NOT USE ON ST. AUGUSTINE, FLORATAM, BENTGRASS, CARPETGRASS, DICHOCHORDA OR LIPPIN.
NO SE DEBE USAR EN CESPEDES DE GRAMA SAN AGUSTIN, FLORATAM, AGROSTIS, GRAMA ALFOMBRERA, OREJA DE RATON O LIPPIN.

NET WT./PESO NETO 16.67 lb (7.56 kg)

ACTIVE INGREDIENTS:
2, 4-D* 1.03%
Mecoprop-p** 0.52%
OTHER INGREDIENTS 98.45%
TOTAL 100.00%

*0.0103 lbs 2,4-D acid equivalent per pound of product
**0.0052 lbs MCP-P acid equivalent per pound of product

KEEP OUT OF REACH OF CHILDREN
CAUTION

MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION

Won't Burn Lawn Guaranteed!

WINTERGUARD® FALL WEED AND FEED!
27-0-12 Covers 5,000 SQ FT

29206
1



SUPER Turf Builder

Scotts® & Your Environment

We care.
Scotts® products are designed with care to grow thicker, greener lawns.

A green lawn is a good lawn.
A healthy lawn can clean the air, produce oxygen and prevent runoff and soil erosion.

You can make a difference.
Help keep our water resources clean. Apply this product only to your lawn, and sweep up any product that lands in the driveway, sidewalk or street back onto your lawn.

Tips for Lawncare Success:

- Feed regularly for your best lawn.** Regular feedings of Scotts® Super Turf Builder® will help build a lawn you and your family can enjoy.
- Scotts Super Turf Builder provides the nutrients your lawn needs to look its best and withstand the stresses of weeds, heat, drought and family activity.
- Mow your lawn high.** Mowing high allows the grass to develop a deep root system & helps your lawn use water more effectively.
- Leave the grass clippings on your lawn.** This recycles the plant nutrients back into the soil.
- Conserve water.** Your lawn will begin to wilt when water is needed. As much as possible, take advantage of nature's sprinkler and rely on the rain to water your lawn.

Plan your feeding schedule. Choose the right product for your lawn.

ANY SEASON	EARLY SPRING February-April	LATE SPRING April-June	SUMMER June-August	FALL September-November
Apply any season if weeds or bugs are not a problem: Scotts® Super Turf Builder® Lawn Food 	To prevent crabgrass: Scotts® Super Turf Builder® with HALT® Crabgrass Preventer 	To kill weeds: Scotts® Super Turf Builder® Weed and Feed 	To kill bugs: Scotts® Super Turf Builder® with SummerGuard® 	Scotts® Super Turf Builder® WinterGuard® Fall Lawn Food To kill fall weeds: Scotts® Super Turf Builder® WinterGuard® Fall Weed and Feed  

The SUPER Turf Builder Advantage

Quick Greening and Extended Long Lasting Feeding

Your lawn requires essential nutrients to thrive.

Builds a thicker, sturdier lawn this fall. Better greenup next spring.



Well-fed lawns are more efficient at using available water.

Feeding with Scotts® helps grow a thick lawn with more roots to better absorb water and nutrients than unweaned lawns. This helps protect your lawn against heat, drought and other stresses.

Get expert advice for your lawn NOW!

Want information on how to use and when?

Sign up today to get free expert advice specifically for your lawn.

Visit our website: www.scotts.com Consumer Helpline: **1-800-543-TURF (8873)**

Application	Aplicación
Directions for Use It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds (0.7 kg) of 2-4-D per acre per application. The maximum number of broadcast applications per treatment site is two per year. A minimum of 30 days between applications is required. Super Turf Builder® WinterGuard® Fall Weed and Feed helps you get a thicker, sturdier lawn with increased root development. Apply it as your last feeding of the year to help prepare your lawn for winter and provide better greenup next spring. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until 24 hours have passed. Do not apply directly to or near water, stream drains, ditches, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and coasts. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when rain is expected that day. - Apply anytime weeds are actively growing (when daytime temperatures are consistently above 60°F). - Apply on a calm day. - For best results apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Super Turf Builder® WinterGuard® Fall Weed and Feed. - On bahia and centipede lawns, wait until after the second mowing of the year, but NOT before mid-March. In the fall, DO NOT treat centipede after the first of October and DO NOT treat after the first of November. - Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustinegrass, Floratam, Carpetgrass, Ligula and Dichroa lawns. On bermudagrass, bahiagrass, centipede grass, zoysiagrass lawns, under certain conditions, some temporary discoloration may result.	Instrucciones de uso Es una violación de la ley federal usar este producto con otras líneas que las indicadas en el etiqueto. Este producto aplica 1.5 libras (0.7 kg) de 2-4-D por acre por aplicación. El número máximo de aplicaciones (separadamente) por área de tratamiento es dos por año. Se requiere un mínimo de 30 días entre aplicaciones. El Super Turf Builder® WinterGuard® Fall Weed and Feed ayuda a obtener un césped más tupido y fuerte con un mayor desarrollo de la raíz. Se aplica como la última fertilización del año para ayudar a preparar su césped para el invierno y obtener un mejor verdeamiento la siguiente primavera. No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o a través de la deriva del producto. Durante la aplicación, mantenga alejados a las personas y las mascotas. No permita que las personas o las mascotas ingresen al área tratada hasta que el producto se haya asentado. No aplique el producto directamente sobre o cerca de agua, drenajes de alcantarillas, alcantarías, arroyos o canales. No aplique el producto a menos de 8 metros de ríos, estanques de peces, lagos, embalses, pantanos, estuarios, bahías y costas. No aplique el producto cuando hay viento. Aplique este producto directamente sobre el césped o jardín y barra hacia el área tratada el producto que caiga sobre la entrada de automóviles, la vereda, canchales o el calle. Para evitar el escurrimiento, no riegue en exceso el área tratada ni la aplique en condiciones de lluvia o en días con pronóstico de lluvia. - Se debe aplicar en cualquier momento en que las malezas estén creciendo activamente (cuando las temperaturas del día estén arriba de 16°C consistentemente). - Se debe aplicar en un día tranquilo. - Se debe aplicar a la grama húmeda para obtener mejores resultados. Para ayudar a que las partículas se adhieran a las hojas de las malezas, la grama debe estar completamente mojada o rociada con rocío la noche antes de aplicar el Super Turf Builder® WinterGuard® Fall Weed and Feed. - En céspedes con grama bahia y céspedes se debe aplicar hasta la segunda vez que corte la grama en el año pero NO antes de mediados de marzo. En el otoño NO se debe tratar la grama centipede después del primer de octubre y NO tratar la grama bahia después del primer de noviembre. - No se debe usar este producto en flores, arbustos, frutas, plantas de vegetales, ni en céspedes con grama agrostis, estentorina, grama San Agustín, Floratam, grama de alfombra, Ligula y orégano de ratón. En céspedes con grama Bermuda, grama bahia, grama centipede, grama zoysa puede ocurrir alguna decoloración temporal bajo ciertas condiciones.

Watering or Rain Do not water your lawn for 24 hours after application. Water will wash the weed control off the weeds' leaves and reduce the effectiveness of this product. This product should not be applied if rain is expected within the next 24 hours. IMPORTANT What to Expect Wind stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control. This product will not prevent weed seeds from producing new plants. Extreme Temperatures Avoid applying when temperatures are consistently above 90°F. Seeding/sodding/sprigging Wait until the new grass has been mowed at least four times before applying Super Turf Builder® WinterGuard® Fall Weed and Feed. If you are planning to seed, sod, or sprig, you should wait at least one month after you have applied Super Turf Builder® WinterGuard® Fall Weed and Feed. Easy to Apply For use on bahiagrass, bermudagrass, bluegrass, centipedegrass, fescues, ryegrass, or zoysiagrass lawns. • Do not contaminate water used for irrigation or domestic purposes. • Do not graze treated areas. • Do not feed clippings to livestock. Weeds controlled: <table><tr><td>aster bitter black medic buckhorn buckwheat buckflower butterweed</td><td>carpetweed chicory chickweed cinquillo cudweed curly dock dandelion</td><td>dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy</td><td>healall horseweed knotted (seedling) lambquarter lippia</td><td>mallow purslane puncture mormongrass mustard pennwort peppercorn shepherd's purse smartweed</td><td>plantain purslane pusley (Florida) ragweed saffron shepherd's smartweed</td><td>sow thistle vetch white clover yellow rocket</td></tr></table>	aster bitter black medic buckhorn buckwheat buckflower butterweed	carpetweed chicory chickweed cinquillo cudweed curly dock dandelion	dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy	healall horseweed knotted (seedling) lambquarter lippia	mallow purslane puncture mormongrass mustard pennwort peppercorn shepherd's purse smartweed	plantain purslane pusley (Florida) ragweed saffron shepherd's smartweed	sow thistle vetch white clover yellow rocket	Riego o lluvia No debe regar ni lavar las siguientes 24 horas después de la aplicación. El agua lavará el herbicida de las hojas de las malezas y reducirá la efectividad de este producto. No se debe aplicar este producto si se espera lluvia dentro de las siguientes 24 horas. IMPORTANTE Qué debe esperar Los tallos y hojas de las malezas se empezarán a retorcer y a poner rizadas en pocos días, pero puede ser necesario un mes o más para controlarlas. Este producto no previene que las semillas de las malezas produzcan nuevas plantas. Temperaturas extremas Evite aplicar cuando las temperaturas estén arriba de 32°C consistentemente. Siembra/cobertura de tepales/rotación Espera a que la nueva grama haya sido cortada por lo menos cuatro veces antes de aplicar el Super Turf Builder® WinterGuard® Fall Weed and Feed. Si planea sembrar, colocar nuevos tepales o nuevos rotaciones, debes esperar al menos un mes después de haber aplicado el Super Turf Builder® WinterGuard® Fall Weed and Feed. Fácil de aplicar Para uso en céspedes con grama bahia, grama Bermuda, pasto azul (Bluegrass), grama centipede, festuca, riego, o grama zoysa. • No contaminar el agua usada para irrigación ni con fines domésticos. • No usar las áreas tratadas para pastar. • No dar de los recortes de césped a los animales. Las malezas que controla: <table><tr><td>aster malva bitter black medic buckhorn buckwheat buckflower butterweed</td><td>carpetweed chicory chickweed cinquillo cudweed curly dock dandelion</td><td>dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy</td><td>healall horseweed knotted (seedling) lamb's-quarter lipia</td><td>plantain purslane puncture (Florida) ragweed saffron shepherd's smartweed</td><td>sow thistle vetch white clover yellow rocket</td></tr></table>	aster malva bitter black medic buckhorn buckwheat buckflower butterweed	carpetweed chicory chickweed cinquillo cudweed curly dock dandelion	dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy	healall horseweed knotted (seedling) lamb's-quarter lipia	plantain purslane puncture (Florida) ragweed saffron shepherd's smartweed	sow thistle vetch white clover yellow rocket
aster bitter black medic buckhorn buckwheat buckflower butterweed	carpetweed chicory chickweed cinquillo cudweed curly dock dandelion	dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy	healall horseweed knotted (seedling) lambquarter lippia	mallow purslane puncture mormongrass mustard pennwort peppercorn shepherd's purse smartweed	plantain purslane pusley (Florida) ragweed saffron shepherd's smartweed	sow thistle vetch white clover yellow rocket								
aster malva bitter black medic buckhorn buckwheat buckflower butterweed	carpetweed chicory chickweed cinquillo cudweed curly dock dandelion	dichondra dandelion English daisy eveningprimrose filaree flexibane ground ivy	healall horseweed knotted (seedling) lamb's-quarter lipia	plantain purslane puncture (Florida) ragweed saffron shepherd's smartweed	sow thistle vetch white clover yellow rocket									

Spreader Settings

Scotts® Broadcast/Rotary Spreaders

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Shut off spreader or stop each time you reach a header strip. Each pass should be within 5 feet of each other to ensure an overlap.

Scotts® Drop Spreaders

Make sure spreader wheel completely overlaps previous wheel marks.

For additional spreader settings, please call 1-800-543-TURF (8873).

Covers 5,000 sq. ft.

Posiciones del esparcidor

Esparcidores al voleo/giratorios de Scotts®

Aplique franjas guía. Luego aplique el producto en la dirección más larga del césped. Para obtener los mejores resultados, utilice un esparcidor Scotts®. Apague el esparcidor o pare cada vez que llegue a una franja guía. Cada pasada debe ser a menos de 1.5 metros de la otra para asegurar un traslape.

Esparcidores de goteo Scotts®

Asegúrese de que la rueda del esparcidor traslape completamente las marcas previas de ruedas.

Para conocer las posiciones adicionales del esparcidor, comuníquese al 1-800-543-TURF (8873).

First Aid & Environmental Information

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION, Causes moderate eye irritation, Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, shoes, and socks. Wash hands and exposed skin with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

First Aid

- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
- IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not get anything by mouth to an unconscious person.
- IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards: 2-4-D is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. McCropoxy may adversely affect non-target plants. Do not contaminate water when disposing of equipment washwater or rinsate. 2-4-D has properties and characteristics associated with chemicals detected in groundwater. The use of 2-4-D in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Storage and Disposal

Do not contaminate food or foodstuffs.

PESTICIDE STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE CONTAINER DISPOSAL: Nonrecyclable container. Do not reuse or refill this container. IF EMPTY: Place in trash or offer for recycling if available.

PARTICLY FILLER: Call your local solid waste disposal agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Primeros auxilios e información ambiental

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN

Avisos de Precaución

Riesgos para humanos y animales domésticos

PRECAUCIÓN, Causa irritación moderada en los ojos y en la ropa. Evite el contacto con los ojos o la ropa. Para utilizar este producto, debe llevar camisa de manga larga, pantalones largos, calcetines y zapatos. Lávese las manos y la piel después con agua y jabón después de manipularlo. Evite el contacto con el producto o frecuente con la piel, puede provocar reacciones alérgicas en algunas personas.

Primeros Auxilios

- SI ENTRA EN LOS OJOS: Mantenga el ojo abierto y enjuague con agua después de con cuidado por 15 a 20 minutos. Si tuviera lentes de contacto, quítelos después de lavarse con agua los primeros 5 minutos y después continúe enjuagando el ojo.
- SI ENTRA EN CONTACTO CON LA PIEL O LA ROPA: Quite la ropa contaminada. Enjuague la piel inmediatamente con mucha agua por 15 a 20 minutos.
- SI SE TRAGÓ: Llame inmediatamente al CENTRO de control de envenenamiento o al médico para obtener recomendaciones de tratamiento. Haga que la persona tome sorbos de un vaso con agua, si puede tragar. No induzca al vómito a menos que lo indique un centro de control de envenenamiento o el médico. No le dé nada por la boca a una persona inconsciente.
- SI SE INHALA: Lleve a la persona al aire fresco. Si la persona no respira llame a 911 o a una ambulancia, luego de respiración artificial, preferentemente de boca a boca, si es posible.

Llame al centro de control de envenenamiento o al médico para obtener recomendaciones de tratamiento. Tenga a la mano el envase del producto o la etiqueta cuando llame al centro de control de envenenamiento o al médico, o al buscar tratamiento.

Scotts® Super Turf Builder® WinterGuard® Fall Weed and Feed 27-4-12 F 643

Guaranteed Analysis

Total Nitrogen (N)	27%
16.0% Free Nitrogen	
11.0% Other Water Soluble Nitrogen	
1.0% Water Insoluble Nitrogen†	
Soluble Potash (K ₂ O)	12%
Sulfur (S)	4%
4.0% Combined Sulfur (S)	
Derived from: Methylenesulfonate, Urea and Potassium Sulfate.	
†Contains 93% slowly available nitrogen from methylenesulfonate.	
EPA Reg. No. 538-283	EPA Est. No. 538-0H-11 86565-OH-11
Supersoluble used is first letter of run code on end of container.	

Questions / Information

Sold and Guaranteed by: The Scotts Company

14111 ScottsLawn Rd.
Marysville, Ohio 43041

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Metals in this product are in compliance with the law and do not pose a health risk to humans or animals. Metals in this product are in compliance with the law and do not pose a health risk to humans or animals.

©2012 The Scotts Company LLC. World rights reserved.

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

AVISO: En forma consistente con la ley aplicable, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you may return it for a full refund of your money. Simply send us evidence of purchase and we will mail you a refund check promptly. Si por cualquier razón usted, el consumidor, no está satisfecho después de usar este producto, tiene derecho a que se le devuelva su dinero. Simplemente envíenos su evidencia de compra y nosotros le enviaremos rápidamente por correo un cheque de reembolso.

120" SIZE 12-digit UPC-A (non-suppressed) For Position Only

0 32247 29206 1

PROOF OF PURCHASE BG1896 I VID-BB

Scotts®
LawnPro®

**Water
Smart®**

SCOTTS® BEST FERTILIZER!




SUPER ***Turf Builder®***

WINTERGUARD®
FALL WEED AND FEED!



Kills dandelions, chickweed, white clover and other broadleaf weeds



Protects and Strengthens for a Better Spring Lawn



Improves Lawn's Ability to Absorb Water & Nutrients

DO NOT USE ON ST. AUGUSTINE, FLORATAM, BENTGRASS, CARPETGRASS, DICHONDRA OR LIPPIA.
NO SE DEBE USAR EN CESPEDES DE GRAMA SAN AGUSTIN, FLORATAM, AGROSTIS, GRAMA ALFOMBRA, OREJA DE RATON O LIPPIA.

NET WT./PESO NETO 16.67 lb (7.56 kg)

ACTIVE INGREDIENTS:	
2, 4-D*	1.03%
Mecoprop-p**	0.52%
OTHER INGREDIENTS:	98.45%
TOTAL:	100.00%

*0.0103 lbs 2,4-D acid equivalent per pound of product
**0.0052 lbs MCPP-p acid equivalent per pound of product

KEEP OUT OF REACH OF CHILDREN
CAUTION SEE BACK FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION CONSULTE EL REVERSO PARA CONOCER LOS AVISOS DE PRECAUCION ADICIONALES.



SUPER Turf Builder®

WINTERGUARD® FALL WEED AND FEED,
27-0-12 Covers 5,000 SQ FT **29206**
1

Won't Burn Lawn Guaranteed!



& Your Environment >>

We care.

Scotts® products are designed with care to grow thicker, greener lawns.

A green lawn is a good lawn.

A healthy lawn can clean the air, produce oxygen and prevent runoff and soil erosion.

You can make a difference.

Help keep our water resources clean. Apply this product only to your lawn, and sweep up any product that lands in the driveway, sidewalk or street back onto your lawn.

Tips for Lawn Care Success:

- **Feed regularly for your best lawn.** Regular feedings of Scotts® Super Turf Builder® will help build a lawn you and your family can enjoy. Scotts Super Turf Builder provides the nutrients your lawn needs to look its best and withstand the stresses of weeds, heat, drought and family activity.
- **Mow your lawn high.** Mowing high allows the grass to develop a deep root system & helps your lawn use water more effectively.
- **Leave the grass clippings on your lawn.** This recycles the plant nutrients back into the soil.
- **Conserve water.** Your lawn will begin to wilt when water is needed. As much as possible, take advantage of nature's sprinkler and rely on the rain to water your lawn.

Plan your feeding schedule. Choose the right product for your lawn.

ANY SEASON	EARLY SPRING February-April	LATE SPRING April-June	SUMMER June-August	FALL September-November
Apply any season if weeds or bugs are not a problem: Scotts® Super Turf Builder® Lawn Food	To prevent crabgrass: Scotts® Super Turf Builder® with HALTS® Crabgrass Preventer	To kill weeds: Scotts® Super Turf Builder® Weed and Feed	To kill bugs: Scotts® Super Turf Builder® with SummerGuard®	Scotts® Super Turf Builder® WinterGuard® Fall Lawn Food To kill fall weeds: Scotts® Super Turf Builder® WinterGuard® Fall Weed and Feed

The SUPER Turf Builder Advantage

Quick Greening and Extended Long Lasting Feeding

Your lawn requires essential nutrients to thrive.

Builds a thicker, sturdier lawn this fall. Better greenup next spring.



Well-fed lawns are more efficient at using available water.

Feeding with Scotts® helps grow a thick lawn with more roots to better absorb water and nutrients than unfed lawns. This helps protect your lawn against heat, drought and other stresses.



Get expert advice for your lawn NOW!

Want information on what to use and when?

Sign up today to get free expert advice specifically for your lawn.

Visit our website:

www.scotts.com

Consumer Helpline:

1-800-543-TURF (8873)

Application

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product applies 1.5 pounds of 2,4-D per acre per application.

The maximum number of broadcast applications per treatment site is two per year.

A minimum of 30 days between applications is required.

Super Turf Builder® WinterGuard® Fall Weed and Feed helps you get a thicker, sturdier lawn with increased root development. Apply it as your last feeding of the year to help prepare your lawn for winter and provide better greenup next spring.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. Do not prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

• **Apply anytime weeds are actively growing (when daytime temperatures are consistently above 60°F).**

• **Apply on a calm day.**

• **For best results apply to moist grass.** To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Super Turf Builder® WinterGuard® Fall Weed and Feed.

• **On Bahia and centipede lawns, wait until after the second mowing of the year, but NOT before mid-March.** In the fall, **DO NOT** treat centipede after the first of October and **DO NOT** treat Bahia after the first of November.

• **Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustinegrass, Floratam, Carpetgrass, Lippia and Dichondra lawns.** On bermudagrass, bahiagrass, centipedegrass, zoysiagrass lawns, under certain conditions, some temporary discoloration may result.

Watering or Rain

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weeds' leaves and reduce the effectiveness of this product. This product **should not** be applied if rain is expected within the next 24 hours.

IMPORTANT

What to Expect

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for control. This product will not prevent weed seeds from producing new plants.

Extreme Temperatures

Avoid applying when temperatures are consistently above 90°F.

Seeding/sodding/sprigging

Wait until the new grass has been mowed at least four times before applying Super Turf Builder® WinterGuard® Fall Weed and Feed. If you are planning to seed, sod, or sprig, you should wait at least one month after you have applied Super Turf Builder® WinterGuard® Fall Weed and Feed.

Easy to Apply

For use on bahiagrass, bermudagrass, bluegrass, centipedegrass, fescues, ryegrass, or zoysiagrass lawns.

• **Do not contaminate water used for irrigation or domestic purposes.**

• **Do not graze treated areas.**

• **Do not feed clippings to livestock.**

Weeds controlled:

aster	carpetweed	dichondra	heala	mallow	plantain	sow thistle
bitter cross	dollarweed	henbit	horseweed	pursh	plantain	velvet
black medic	chickweed	English daisy	knotted	ragweed	shepherd's purse	white clover
blackhorn	cinefoil	evening primrose	mustard	rusch	shepherd's purse	yellow rocket
bull thistle	cudweed	filaree	(seeding)	shepherd's purse	shepherd's purse	
buckhorn	curly dock	flaxseed	lamb's quarter	shepherd's purse	shepherd's purse	
bottomweed	dandelion	ground ivy	lippia	shepherd's purse	shepherd's purse	

Aplicación

Instrucciones de uso

Es una violación a la ley Federal usar este producto con otros fines que los indicados en la etiqueta. Este producto brinda 0.7 kg de 2,4-D por acre por aplicación.

El número máximo de aplicaciones (esparcimientos) por área de tratamiento es dos por año.

Se requiere un mínimo de 30 días entre aplicaciones.

El Super Turf Builder® WinterGuard® Fall Weed and Feed le ayuda a obtener un césped más tupido y fuerte con un mayor desarrollo de la raíz. Se aplica como la última fertilización del año para ayudarle a preparar su césped para el invierno y obtener un mejor reverdecimiento la siguiente primavera.

No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o a través de la desviación del producto. Durante la aplicación, mantenga alejadas a las personas y las mascotas. No permita que las personas o las mascotas ingresen al área tratada hasta que el producto se haya asentado.

No aplique el producto directamente sobre o cerca de agua, drenajes de tormenta, candelones, alcantarillas o zanjas de drenaje. No aplique el producto a menos de 8 metros de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías y océanos. No aplique el producto cuando hay viento. Aplique este producto directamente sobre el césped o jardín y barra hacia el área tratada el producto que caiga sobre la entrada de automóviles, la vereda, candelones o la calle. Para evitar el escurrimiento, no riegue en exceso el área tratada ni lo aplique en condiciones de lluvia o en días con pronostico de lluvia.

• **Se debe aplicar en cualquier momento en que las malezas estén creciendo activamente (cuando las temperaturas del día estén arriba de 16°C consistentemente).**

• **Se debe aplicar en un día tranquilo.**

• **Se debe aplicar a la grama húmeda para obtener mejores resultados.** Para ayudar a que las partículas se adhieran a las hojas de las malezas, la grama debe estar completamente regada o mojada con rocío de la noche antes de aplicar el Super Turf Builder® WinterGuard® Fall Weed and Feed.

• **En céspedes con grama bahía y ciempies se debe esperar hasta la segunda vez que corte la grama en el año pero NO antes de mediados de marzo.** En el otoño NO se debe tratar la grama ciempies después del primero de octubre y NO trate la grama bahía después del primero de noviembre.

• **No se debe usar este producto en flores, arbustos, frutas, plantas de vegetales, ni en céspedes con grama agrostis estolonifera, grama San Agustín, Floratam, grama alombra, Lippia y oreja de ratón.** En céspedes con grama Bermuda, grama bahía, grama ciempies, grama zoysia puede ocurrir alguna decoloración temporal bajo ciertas condiciones.

Riego o lluvia

No debe regar en las siguientes 24 horas después de la aplicación. El agua lavará el herbicida de las hojas de las malezas y reducirá la efectividad de este producto. **No se debe** aplicar este producto si se espera lluvia dentro de las siguientes 24 horas.

IMPORTANTE

Qué debe esperar

Los tallos y hojas de las malezas se empezarán a retorcer y a poner rizados en pocos días, pero puede ser necesario un mes o más para controlarlas. Este producto no previene que las semillas de las malezas produzcan nuevas plantas.

Temperaturas extremas

Debe evitar aplicar cuando las temperaturas estén arriba de 32°C consistentemente.

Siembra/colocación de tepes/retoños nuevos

Espera a que la nueva grama haya sido cortada por lo menos cuatro veces antes de aplicar el Super Turf Builder® WinterGuard® Fall Weed and Feed. Si tiene planeado sembrar, colocar nuevos tepes o nuevos retoños deberá esperar por lo menos un mes después de haber aplicado el Super Turf Builder® WinterGuard® Fall Weed and Feed.

Fácil de aplicar

Para uso en céspedes con grama bahía, grama Bermuda, pasto azul (Bluegrass), grama ciempies, festucas, raigrás, o grama zoysia.

• **No contamine el agua usada para irrigación ni con fines domésticos.**

• **No use las áreas tratadas para pastar.**

• **No les des los recortes de césped a los animales.**

Las malezas que controla:

áster	anicolis	oreja de ratón	hierba mora	malva	lantén	persicaria
maizuelo	achicoria	acariaca	origan	hierba de la moneda	chicaballo	chicaballo
alfalfa	alsine	margarita inglesa	erigeron canadensis	ipomoea	veza	veza
lantén	potentilla	primula nocturna	cien nudos	mostaza	trebol blanco	trebol blanco
cardo negro	siempreviva	alfalfa	(plantillas)	centaurea asiática	jaramago	jaramago
trébol erizo	romaza	erigeron	quinta	lepidio	pimpinela escarlata	pimpinela escarlata
hierba de botón	diente de león	hierba terrestre	lippia	bledo	bolsa del pastor	bolsa del pastor

Spreader Settings

This package applies 3.3 lbs.
of product per 1,000 sq. ft.

Covers 5,000 sq. ft.

Scotts® Broadcast/Rotary Spreaders

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Shut off spreader or stop each time you reach a header strip. Each pass should be within 5 feet of each other to ensure an overlap.

Scotts® Drop Spreaders

Make sure spreader wheel completely overlaps previous wheel marks.

For additional spreader settings, please call 1-800-543-TURF (8873).

Posiciones del esparcidor

Este paquete aplica 1.5 kg
de producto por cada 93
metros cuadrados.

Cubre 93
metros cuadrados

Esparcidores al voleo/giratorios de Scotts®

Aplique franjas guía. Luego aplique el producto en la dirección más larga del césped. Para obtener los mejores resultados, utilice un esparcidor Scotts®. Apague el esparcidor o pare cada vez que llegue a una franja guía. Cada pasada debe ser a menos de 1.5 metros de la otra para asegurar un traslape.

Esparcidores de goteo Scotts®

Asegúrese de que la rueda del esparcidor traslape completamente las marcas previas de ruedas.

Para conocer las posiciones adicionales del esparcidor, comuníquese al 1-800-543-TURF (8873).

First Aid & Environmental Information

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, shoes, and socks. Wash hands and exposed skin with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

First Aid

- **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
- **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS: 2,4-D is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Mecoprop-p may adversely affect non-target plants. Do not contaminate water when disposing of equipment washwater or rinsate. 2,4-D has properties and characteristics associated with chemicals detected in groundwater. The use of 2,4-D in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Storage and Disposal

Do not contaminate feed or foodstuffs.

- **PESTICIDE STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.
- **PESTICIDE CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. IF EMPTY: Place in trash or offer for recycling if available.
- **IF PARTLY FILLED:** Call your local solid waste disposal agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Primeros auxilios e información ambiental

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN

Avisos de Precaución

Riesgos para humanos y animales domésticos

PRECAUCIÓN. Causa irritación moderada en los ojos. Evite el contacto con los ojos o la ropa. Para utilizar este producto, debe llevar camisa de manga larga, pantalones largos, calcetines y zapatos. Lave completamente las manos y la piel expuesta con agua y jabón después de manipularlo. En algunas personas, el contacto prolongado o frecuente con la piel, puede provocar reacciones alérgicas.

Primeros Auxilios

- **SI ENTRA EN LOS OJOS:** Mantenga el ojo abierto y enjuague con agua despacio y con cuidado por 15 a 20 minutos. Si tuviera lentes de contacto, quítelos después de lavarse con agua los primeros 5 minutos y después continúe enjuagando el ojo.
- **SI ENTRA EN CONTACTO CON LA PIEL O LA ROPA:** Quite la ropa contaminada. Enjuague la piel inmediatamente con mucha agua por 15 a 20 minutos.
- **SI SE TRAGA:** Llame inmediatamente al centro de control de envenenamiento o al médico para obtener recomendaciones de tratamiento. Haga que la persona tome sorbos de un vaso con agua, si puede tragar. No induzca el vómito a menos que lo indique un centro de control de envenenamiento o el médico. No le dé nada por la boca a una persona inconsciente.
- **SI SE INHALA:** Lleve a la persona al aire fresco. Si la persona no respira llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible. Llame al centro de control de envenenamiento o al médico para obtener recomendaciones de tratamiento. Tenga a la mano el envase del producto o la etiqueta cuando llame al centro de control de envenenamiento o al médico, o al buscar tratamiento.

RIESGOS AMBIENTALES: 2,4-D es tóxico para los peces e invertebrados acuáticos. No lo aplique directamente en el agua, en áreas donde hay agua superficial o en áreas entre mareas por debajo de la marca de marea alta promedio. El desplazamiento o el escurrimiento del producto pueden ser peligrosos para los organismos acuáticos en aguas adyacentes a las áreas tratadas. Mecoprop-p puede afectar de manera adversa a plantas no indicadas para el tratamiento. No contamine el agua cuando elimine la que utilizó para lavar o enjuagar el equipo. 2,4-D tiene propiedades y características asociadas con las sustancias químicas detectadas en aguas subterráneas. La utilización de 2,4-D en áreas donde los suelos son permeables, en particular donde el manto freático es superficial, puede derivar en la contaminación de las aguas subterráneas. La aplicación en torno a un cisterna o a un pozo puede provocar la contaminación del agua potable o las aguas subterráneas.

Almacenamiento y Eliminación

No contamine comestibles ni alimentos para animales.

- **ALMACENAMIENTO DEL PESTICIDA:** Guardar en lugar limpio y seco. Cierre la bolsa abierta doblando la parte superior y asegurándola.
- **ELIMINACIÓN DEL ENVASE DEL PESTICIDA:** Envase no reutilizable. No vuelva a utilizar o llenar este envase. **SI ESTA VACÍO:** Tírela a la basura o, si está disponible, ofrezcámosla para reciclar. **SI ESTA PARCIALMENTE LLENTO:** Llame a su agencia de eliminación de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto no utilizado en drenajes internos o externos.

Scotts® Super Turf Builder® Winterguard® Fall Weed and Feed 27-0-12 F 643

Guaranteed Analysis

Total Nitrogen (N)	27%
15.0% Urea Nitrogen	
11.0% Other Water Soluble Nitrogen†	
1.0% Water Insoluble Nitrogen†	
Soluble Potash (K ₂ O)	12%
Sulfur (S)	4%
4.0% Combined Sulfur (S)	

Derived from: Methyleneureas, Urea and Potassium Sulfate.

†Contains 9% slowly available nitrogen from methyleneureas.

EPA Reg. No. 538-283 EPA Est. No. 538-OH-1^S, 85653-OH-1^A
Superscript used is first letter of run code on end of container.

Questions / Information

Sold and Guaranteed by:
The Scotts Company
14111 Scottslawn Rd.
Marysville, Ohio 43041

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Información relacionada con el contenido y con los niveles de metales en este producto se encuentra disponible en Internet en <http://www.regulatory-info-sc.com>

©2012 The Scotts Company LLC.
World rights reserved.

NOTICE: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

AVISO: En forma consistente con la ley aplicable, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly. Si por cualquier razón usted, el consumidor, no está satisfecho después de usar este producto, tiene derecho a que se le devuelva su dinero. Simplemente envíenos su evidencia de compra y nosotros le enviaremos rápidamente por correo un cheque de reembolso.

120% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only

0 32247 29206 1

PROOF OF PURCHASE

B61896 1 VID-BB

SCOTT'S® SUPER TURF BUILDER® WINTERGUARD® FALL WEED AND FEED 27-0-12 F 643