



DISINFECTANT – SANITIZER – FUNGICIDE – VIRUCIDE* – DEODORIZER

For Hospital, Institutional, Industrial, School, Restaurant, Farm, Food Handling and Processing Areas, Bar and Institutional Kitchen Use

Active Ingredients:	
Octyl decyl dimethyl ammonium chloride	2.25%
Diocetyl dimethyl ammonium chloride	0.90%
Didecyl dimethyl ammonium chloride	1.35%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	3.00%
Other Ingredients.....	92.50%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

See side panel for precautionary statements and first aid.

ChemStation International, Inc.
3400 Encrete Lane, Dayton OH 45439 USA
(937) 294-8265

EPA Registration No. 6836-70-65072

EPA Establishment No.

<input type="checkbox"/> 65072-CO-001	<input type="checkbox"/> 65072-IL-001	<input type="checkbox"/> 65072-OH-005
<input type="checkbox"/> 65072-OH-006	<input type="checkbox"/> 65072-PA-001	<input type="checkbox"/> 65072-PA-002
<input type="checkbox"/> 65072-PA-003	<input type="checkbox"/> 65072-TX-001	<input checked="" type="checkbox"/> 65072-WA-001

Net Contents: _____

KP 3510 per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution) for at least 60 seconds. Articles too large for immersing should be thoroughly wetted by rinsing, spraying, or swabbing. 5. Remove immersed items from solution to drain and then air dry. Non-immersed items must also be allowed to air dry. Do not rinse.

TO SANITIZE SOFT SERVE, FOOD AND BEVERAGE DISPENSING EQUIPMENT:
1. Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing. 2. Fill equipment with a solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallon of water (0.25-0.68 oz. of **KP 3510** per gallon of water) (150-400 ppm active quat). 3. Allow solution to remain in equipment for at least 60 seconds. 4. Drain thoroughly and allow to air dry before reuse. Do not rinse.

TO SANITIZE SANITARY FILLING EQUIPMENT:

Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing. Prepare a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallons of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution) for final washer and rinser applications. Allow surfaces to remain wet for at least 60 seconds. Drain thoroughly and allow to air dry before reuse. Do not rinse.

TO SANITIZE BEER FERMENTATION AND STORAGE TANKS:

Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing. Prepare a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallons of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution) for mechanical or automated systems. Allow surfaces to remain wet for at least 60 seconds. Drain thoroughly and allow to air dry before reuse. Do not rinse.

TO SANITIZE EGG SHELLS INTENDED FOR FOOD:

To sanitize previously cleaned food-grade eggs in shell egg and egg product processing plants, spray with a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallons of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution). The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product shall be subjected to a potable water rinse only if they are to be broken

This product is not for use on medical device surfaces.

DISINFECTION / VIRUCIDAL*/ FUNGICIDAL DIRECTIONS:

Add 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water (0.78 oz. of **KP 3510** per gallon of water) (or equivalent dilution) to disinfest hard, nonporous surfaces. Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. When used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities, this product is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus). Apply use-solution with a cloth, mop, sponge, sprayer or by immersion, thoroughly wetting surfaces. For heavily soiled areas, preclean first. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray. Note: For spray applications, cover or remove all food products. Treated surfaces must remain wet for 10 minutes. For Human Coronavirus treated surfaces must remain wet for 1 minute. Wipe dry with a clean cloth, sponge or mop or allow to air dry.

Rinse food contact surfaces such as counter tops, tables, picnic tables, appliances and/or stove tops with potable water prior to reuse. Do not use on glasses, dishes or utensils as a disinfectant. For heavily soiled areas, preclean first.

BACTERICIDAL STABILITY OF USE-DILUTION: Tests confirm that the use solution of this product, when stored in a sealed container such as a spray bottle, remains effective for up to 5 months. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, properly labeled containers when diluting this product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails.

NON-ACID TOILET BOWL AND URINAL DISINFECTANT DIRECTIONS:

Remove heavy soil prior to disinfection. **From concentrate:** Add 3¹/₂ oz. of **KP 3510** to the toilet bowl or urinal and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop or sponge. Allow to stand for 10 minutes and flush.

From use-solution: Empty toilet bowl or urinal and apply a use-solution of 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water (0.78 oz. of **KP 3510** per gallon of water) (or equivalent dilution) to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly, then allow to stand for 10 minutes and flush.

FOR USE ON FINISHED FLOORS: To limit gloss reduction use 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water. Apply with a damp mop or autoscrubber.

ULTRASONIC BATH DISINFECTANT DIRECTIONS:

Use **KP 3510** to disinfect hard nonporous non-critical instruments / objects compatible with Ultrasonic cleaning units. Pour fresh use-solution of 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water (0.78 oz. per gallon of water) or equivalent dilution directly into bath chamber. Preclean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturer's use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Prepare a fresh solution for each use.

TO SANITIZE NON-FOOD CONTACT SURFACES:

Add 2 oz. of **KP 3510** per 5¹/₈ gallons of water (0.34 oz. of **KP 3510** per gallon of water) (200 ppm active) (or equivalent dilution) to sanitize hard, nonporous surfaces. Apply sanitizer use-solution with a cloth, mop, sponge, sprayer or by immersion, thoroughly wetting surfaces. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray. Note: For sprayer applications, cover or remove all food products. Treated surfaces must remain wet for 1 minute. Wipe dry with a sponge, mop or cloth or allow to air dry. Prepare fresh solution daily or more often if solution becomes visibly diluted, cloudy or soiled.

FOR FOOD CONTACT SURFACES, FOOD PROCESSING EQUIPMENT AND UTENSILS IN DAIRIES AND PUBLIC EATING ESTABLISHMENTS AND FOOD CONTACT SURFACES, FOOD PROCESSING EQUIPMENT AND UTENSILS IN FOOD PROCESSING PLANTS (40 CFR 180.940 (a)):

2 oz. of **KP 3510** per 2 7/8 - 7 3/4 gallons of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat)

TO SANITIZE FOOD CONTACT SURFACES, FOOD PROCESSING EQUIPMENT AND OTHER HARD SURFACES IN FOOD PROCESSING LOCATIONS, DAIRIES, RESTAURANTS AND BARS (IN A THREE COMPARTMENT SINK):

Use **KP 3510** to sanitize hard nonporous surfaces of food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage areas and display equipment and other hard nonporous surfaces. Prior to application, remove gross food particles and soil by a pre-flush, pre-scrape or, when necessary, pre-soak. Then thoroughly wash or flush objects with a good detergent or compatible cleaner followed by a potable water rinse before application of the sanitizing solution. Articles too large for immersing, apply a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallon of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution) to pre-cleaned hard surfaces thoroughly wetting surfaces with a cloth, mop, sponge, or coarse sprayer. Surfaces must remain wet for at least 60 seconds (one minute) followed by adequate draining and air drying. Do not rinse. Immerse pre-cleaned glassware, dishes, silverware, cooking utensils and other similar size food processing equipment in a solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallons of water (0.25 - 0.68 oz. of **KP 3510** per gallon of water) (150 - 400 ppm active quat) (or equivalent dilution) for at least 60 seconds. Drain thoroughly (and allow to air dry) before reuse. Do not rinse. Prepare a fresh sanitizing solution daily or more often if the use solution becomes diluted or soiled. For mechanical application, use-solution may not be reused for sanitizing applications but may be reused for other purposes such as cleaning. Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard nonporous surfaces by cloth, sponge or brush or coarse spray.

TO SANITIZE FOOD PROCESSING EQUIPMENT, UTENSILS, AND OTHER FOOD CONTACT ARTICLES REGULATED BY 40 CFR § 180.940(a) (IN A THREE COMPARTMENT SINK):

1. Scrape, flush, or presoak articles (whether mobile or stationary) to remove gross food particles and soil. 2. Thoroughly wash articles with an appropriate detergent or cleaner. 3. Rinse articles thoroughly with potable water. 4. Sanitize by immersing articles with a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallons of water (0.25 - 0.68 oz. of

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through the skin. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective gloves (rubber, chemical resistant) when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

As a disinfectant, KP 3510 kills:	
Pseudomonas aeruginosa	Staphylococcus aureus
Salmonella enterica	Brevibacterium ammoniagenes
Campylobacter jejuni	Escherichia coli
Escherichia coli pathogenic strain 0157:H7	Klebsiella pneumoniae
Listeria monocytogenes	Pseudomonas cepacia
Salmonella typhi	Human Coronavirus
Yersinia enterocolitica	*Herpes Simplex Virus Type 1
*Herpes Simplex Virus Type 2	*Influenza A2 / Hong Kong Virus
*Vaccinia Virus	*Norovirus - Feline Calicivirus
Aspergillus niger	
Trichophyton mentagrophytes (Athlete's Foot Fungus) (Cause of Ringworm)	

KP 3510 is effective against: Avian Infectious Laryngotracheitis Virus; Arkansas '99 (Infectious Bronchitis Virus); Avian Influenza Virus (Turkey/Wisconsin); Avian Influenza Virus (H5N1); Canine Distemper Virus; Marek's Disease Virus; Newcastle's Disease Virus; Pseudorabies Virus

KP 3510 is an effective sanitizer at 150 ppm active quat for use on food contact surfaces in 400 ppm hard water against: Escherichia coli; Staphylococcus aureus **KP 3510** is an effective sanitizer at 200 ppm active quat for use on food contact surfaces in 500 ppm hard water against: Shigella sonnei; Staphylococcus aureus - Community Associated Methicillin-Resistant [CA-MRSA] [NRS384] [USA300]; Staphylococcus aureus - Community Associated Methicillin-Resistant [CA-MRSA] [NRS123] [USA400]; Staphylococcus aureus - Methicillin-Resistant [MRSA]; Vibrio cholerae

KP 3510 is an effective sanitizer at 400 ppm active quat for use on food contact surfaces in 1000 ppm hard water against: Escherichia coli; Staphylococcus aureus **KP 3510** meets Sanitizer Test standards for non-food contact surface sanitizers. At 200 ppm active, **KP 3510** is an effective one-step sanitizer against Staphylococcus aureus and Klebsiella pneumoniae in 60 seconds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Refillable container. Refill this container with **KP 3510** only. Do not reuse this container for any other purpose.

Pesticide Storage:

Open dumping is prohibited. Store in original container in areas inaccessible to children.

Pesticide Disposal:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

FINAL CONTAINER DISPOSAL

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before refilling is the responsibility of the refiller. Cleaning the container before final disposal is the responsibility of the person disposing of the container.

(Plastic or Metal Containers:) To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in sanitary landfill. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations. (For metal containers only: DO NOT cut or weld metal containers.)

KP 3510 has been cleared under 40 CFR 180.940 (a) for use on food Processing Equipment and Utensils in Dairies and Public Eating Establishments and Food Contact Surfaces, Food Processing Equipment and Utensils in Food Processing Plants and other food-contact articles at a concentration of 150 - 400 ppm active.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

immediately for use in the manufacture of egg products. Eggs must be reasonably dry before casing or breaking. The solution must not be re-used for sanitizing eggs. **Note:** Only clean, whole eggs can be sanitized. Dirty, cracked or punctured eggs cannot be sanitized.

U.S. PUBLIC HEALTH SERVICE FOOD SERVICE SANITIZATION

RECOMMENDATIONS CLEANING AND SANITIZING: Equipment and utensils must be thoroughly pre-flushed or pre-scraped and, when necessary, pre-soaked to remove gross food particles and soil. 1. Thoroughly wash equipment and utensils in hot detergent solution. 2. Rinse utensils and equipment thoroughly with potable water. 3. Sanitize equipment and utensils by immersion in a use solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallon of water (0.25-0.68 oz. of **KP 3510** per gallon of water) (150-400 ppm active quat) (or equivalent dilution) for at least 60 seconds at a temperature of 75°F. 4. For equipment and utensils too large to sanitize by immersion, apply a use-solution of 2 oz. of **KP 3510** per 2¹/₈ - 7³/₈ gallon of water (0.25-0.68 oz. of **KP 3510** per gallon of water)(150-400 ppm active quat) (or equivalent dilution) by rinsing, spraying or swabbing until thoroughly wet. 5. Allow sanitized surface to drain and air dry. Do not rinse.

WATERPROOF GLOVE SANITIZING DIRECTIONS

To prevent cross contamination into processing areas of food plants, waterproof gloves must be sanitized prior to entering or re-entering those areas. Remove gross contamination from gloves before sanitizing. Then place gloved hand in a use-solution of 2 oz of **KP 3510** per 5¹/₈ gallons of water (200 ppm quat) for sixty seconds. Change the solution in the bath at least daily or more often if the solution appears visibly diluted or soiled.

WISCONSIN STATE BOARD OF HEALTH DIRECTIONS FOR EATING

ESTABLISHMENTS: 1. Scrape and pre-wash utensils and glasses whenever possible. 2. Wash with a good detergent or compatible cleaner. 3. Rinse with potable water. 4. Sanitize in a solution of 2 oz. of **KP 3510** per 2¹/₈ - 5¹/₈ gallons of water (0.34 - 0.68 oz. of **KP 3510** per gallon of water) (200 - 400 ppm active quat). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code. 5. Place sanitized utensils on a rack or drain board to air-dry. 6. Prepare a fresh sanitizing solution at least daily or when visibly soiled or diluted.

Note: A clean potable water rinse following sanitization is not permitted under HFS 196, Appendix 7-204.11 of the Wisconsin Administrative Code (reference 40 CFR 180.940 (a)).

(For food processing or other facilities that have installed entryway sanitizing systems.)

ENTRYWAY SANITIZING SYSTEMS:

To prevent cross contamination from area to area, set the system to deliver (0.68 - 1.36 oz. per gallon of water) (2 - 4 oz. per 2¹/₈ per gallon of water) (400 - 800 ppm active) of sanitizing solution. The spray (foam) must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor. Do not mix other foam additives to the sanitizing solution.

SHOE BATH SANITIZER DIRECTIONS:

To prevent cross contamination into animal areas, (and the packaging and storage areas of food plants), shoe baths containing one inch of freshly made sanitizing solution must be placed at all entrances to buildings, and hatcheries, (and production and packaging rooms). Scrape waterproof work boots (shoes) and place in a use-solution of (2 oz. of **KP 3510** per 5¹/₈ gallons of water) (0.34 oz. of **KP 3510** per gallon of water) (200 ppm active quat) (or equivalent dilution) for 60 seconds prior to entering area. If there is a heavy soil load or excessive traffic, place work boots (shoes) in a use solution of (0.68 - 1.36 oz. per gallon of water) (2 - 4 oz. per 2¹/₈ gallons of water) (400 - 800 ppm active) for 60 seconds prior to entering area. Change the solution in the bath at least daily or more often if the solution appears diluted or soiled.

SHOE FOAM SANITIZER DIRECTIONS:

To prevent cross contamination into animal areas, and the packaging and storage areas of food plants. Apply a foam layer approximately 1/2 to 2 inches thick made from a solution of 8 to 12 oz. of **KP 3510** per 5¹/₈ gallons of water (or equivalent dilution) (800 to 1200 ppm active), at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply the foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator or aerator. Scrape waterproof work boots (shoes). Stand or walk through foamed area for 60 seconds prior to entering area. Foam area must be washed and foam replaced at least daily or more often if the foam appears visibly diluted or soiled.

"SPECIAL INSTRUCTIONS FOR INACTIVATING AVIAN INFLUENZA A" AND OTHER ANIMAL VIRUSES LISTED ON THIS LABEL

VETERINARY, ANIMAL CARE AND ANIMAL LABORATORY FACILITIES / ZOOS / PET SHOPS / KENNELS AND FARM PREMISE DISINFECTION / VIRUCIDAL DIRECTIONS:

For cleaning and disinfecting hard nonporous surfaces: equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries pens, stalls and etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, pens, stalls and etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all (troughs, racks and) feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 3 1/2 oz. of **KP 3510** per 4 1/2 gallons of water [or equivalent dilution] for a period of 10 minutes. Wipe or allow to air dry. Immerse all animal handling and restraining equipment as well as forks, shovels, and scrapers used to remove litter and manure. Thoroughly scrub all treated surfaces, then rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. For Veterinary Practice, Animal Care and Animal Laboratory Facilities/Zoos/ Pet Shops/Kennels and Farm Premise Disinfection/Virucidal Directions: Thoroughly scrub all treated feeding and watering appliances with soap or detergent, and rinse with potable water before reuse. Ventilate buildings, animal enclosures, vehicles and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

MUSHROOM FARM PREMISE USE:

Site Preparation: The first step in any on-going sanitation program must be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

Cleaning and Disinfection: For general cleaning and disinfection, use 3¹/₂ oz. of **KP**

3510 per 4¹/₂ gallons of water (0.78 oz. of **KP 3510** per gallon of water) (or equivalent dilution). Apply use solution with a cloth, mop, sponge, sprayer or by immersion to thoroughly wet the surfaces. Treated surfaces must remain wet for 10 minutes. Wipe or allow to air dry. For heavily soiled areas, preclean first. Prepare a fresh solution for each use or more often if solution becomes visibly diluted, clouded or soiled.

For Heavy Duty Cleaning: When greater cleaning is desired, use 7 oz. of **KP 3510** per 4¹/₂ gallons of water. Heavily soiled areas may require repeated cleaning before treatment.

DO NOT APPLY THE USE-SOLUTION TO THE MUSHROOM CROP, COMPOST OR CASING. Rinse treated surfaces with potable water before they contact the crop, compost or casing.

POULTRY AND SWINE PREMISE DISINFECTION DIRECTIONS:

Remove all animals and feeds from premises, vehicles and enclosures such as coops, and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water (0.78 oz. of **KP 3510** per gallon of water) (or equivalent dilution). Saturate surfaces with the disinfecting solution for 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, trucks, coops, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

Hatcheries:

Use to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Use 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water. Leave all treated surfaces wet for 10 minutes or more. Allow to air dry.

Vehicles:

Clean all vehicles including mats, crates, cabs, and wheels with water and **KP 3510**. Use 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water. Apply use solution to treat all vehicles. Leave treated surfaces wet for 10 minutes or more. Allow to air dry.

Dressing Plant:

Disinfect equipment, utensils, walls and floors in poultry and animal dressing plants. Disinfect ofal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all heavy soils prior to application. Saturate all surfaces with a solution of 3¹/₂ oz. of **KP 3510** per 4¹/₂ gallons of water (0.78 oz. of **KP 3510** per gallon of water) and scrub to loosen all soils. Surfaces must remain wet for 10 minutes, and then thoroughly rinsed with potable water before operations are resumed.

CITRUS CANKER DISEASE CONTROL:

For prevention of Citrus Canker Disease through treatment of precleaned equipment. Effective against Xanthomonas axonopodis pv. citri (Citrus Canker Disease) at 2000 ppm active quat. Treatments can be applied to trucks, attached trailers and field harvesting equipment including cargo area, wheels, tires, undercarriage, hood, roof, fenders, and any other hard nonporous part of the equipment that can be taken into infested areas. Thoroughly clean all surfaces with soap or detergent and rinse with water. Then saturate surfaces with a 1:37 use-dilution (6 oz. of **KP 3510** per 1¹/₄ gallons of distilled water or equivalent dilution) (2000 ppm active quat) for a period of 10 minutes. Allow to air dry. All surfaces that come in contact with food or crop must be rinsed with potable water before reuse. To prevent the spread of Citrus Canker Disease by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing. Clothing must be either rinsed or laundered before reuse. Footwear must be rinsed before reuse.

FOGGING DEVICE HATCHERY ROOM SANITIZER DIRECTIONS:

All surfaces must be cleaned and disinfected in accordance with label directions prior to fogging. Fogging is an adjunct or supplement to normal cleaning and disinfectant procedures and practices. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Mix two parts of **KP 3510** to five parts water (1 gallon of **KP 3510** to 2.5 gallons water). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, for one minute for each 4000 cubic feet of space in the room.