

0.875"

**Scotts**  
**4 STEP**  
PROGRAM

**STEP 2**

Weed Control plus Lawn Food  
Apply in Late Spring: APR | MAY | JUN

23615-1  
85% SIZE  
12-digit UPC-A  
(non-suppressed)  
For Position Only  
32247 23615

23615-1

OUR BEST  
ANNUAL  
PROGRAM

**Scotts**

**4 STEP**  
PROGRAM



Improves Lawn's Ability to  
Absorb Water & Nutrients

Covers 5,000 SQ FT

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
See back for additional precautionary statements.

ACTIVE INGREDIENTS:  
2,4-D<sup>1</sup> ..... 1.18%  
Mecoprop-p<sup>11</sup> ..... 0.59%  
OTHER INGREDIENTS: ..... 98.23%  
Total 100.00%

<sup>1</sup>0.01% to 2.5-0.01% depending on grade of product.  
<sup>11</sup>0.01% to 2.5-0.01% depending on grade of product.

**28-0-3**

NET WT. 14.63 lb (6.64 kg)

Available only at authorized Scotts

**LawnPro**  
Dealers

**Scotts** **4 STEP**  
PROGRAM



**STEP 2**

Weed Control plus  
Lawn Food

DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM,  
DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS.



Apply in Late Spring:  
**APR | MAY | JUN**

**STEP 2** Weed Control plus Lawn Food  
Apply in Late Spring: APR | MAY | JUN





# Scotts®

## 4 STEP™ PROGRAM

### Now it's Easy for You to Have a Better Lawn



#### Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed.

#### Simple and easy

No more guesswork. You simply apply the right product – one STEP® at a time.

#### Guaranteed results

Each STEP® product is designed to the specific season to help give you a greener lawn.

#### Florida Annual Fertilizer Applications

On Centipede lawns, do not exceed rate of 2 lbs of Nitrogen per 1,000 sq. ft. in the north and 3 lbs of Nitrogen per 1,000 sq. ft. in the central and south. On Bahia lawns, do not exceed rate of 3 lbs of Nitrogen per 1,000 sq. ft. in the north.

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

#### SPREADER SETTINGS

Spreader Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. Covers 5,000 sq. ft. This package applies 2.9 lbs of product per 1,000 sq. ft.

	Spreader Settings	Directions		Spreader Settings	Directions
Scotts® Broadcast Rotary Spreaders	3 1/2	Each pass should be within 5 feet of each other to ensure overlap.		Drop Spreaders	5 3/4
					Make sure spreader wheel completely overlaps previous wheel marks.

For additional spreader settings please call 1-800-543-TURF (8873)

#### GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline:  
**1-800-543-TURF**  
(8873)

Visit our website:  
**www.scotts.com**



#### Well-fed lawns are more efficient at using available water.

Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product applies 1.5 pounds of 2,4-D per acre per application.

#### APPLICATION

- Apply on a calm day.
- Make sure that the grass is wet from dew or sprinkling before applying. (This helps particles adhere to weed leaves to produce best results.)
- Set spreader to the setting indicated by the chart above before applying.
- Kills weeds including dandelion, white clover, chickweed, plantain and ragweed.
- Apply when daytime temperatures are between 60 and 90 degrees F.
- Apply only to grass. Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetables.
- Grass should not touch the bottom of a drop spreader. Tall grass should be mowed a day or two before application.
- Delay application on newly seeded, sodded or sprigged areas until new grass has been mowed four times.
- Wait at least one month after using this product before you seed, sod or plant grass sprigs.
- Not for use on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass lawns.
- On bahia or centipede grass lawns, wait until after the second mowing of the year, but not before mid-March.
- In fall, **DO NOT** treat centipede grass after Oct. 1, nor bahia grass after Nov. 1.
- Wait at least 24 hours after application before you water or mow the lawn (Granules must remain on weed leaves for 24 hours to be effective.)

#### USE RESTRICTIONS

- Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.
- Do not apply more than twice per year.
- A minimum of 30 days between applications is required.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### KEEP OUT OF REACH OF CHILDREN

#### CAUTION

#### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical resistant gloves made of any waterproof material.

#### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling poison control center or doctor or going for treatment.

#### ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

EPA Reg. No. 538-270 EPA Est. No. 538-OH-1 #CAS #94-75-7 #CAS #16484-77-8

STEP® 2 Weed Control Plus Lawn Food 20-0-3 F 643

#### Guaranteed Analysis

Total nitrogen (N)..... 28%  
5.2% ammoniacal nitrogen  
12.7% urea nitrogen  
9.3% other water soluble nitrogen\*  
0.8% water insoluble nitrogen\*  
Soluble potash (K<sub>2</sub>O)..... 3%  
Sulfur (S)..... 7%  
7.0% combined sulfur (S)

Derived from: Methylenesulfate, potassium sulfate, ammonium sulfate, urea.

\*Contains 7.5% slowly available nitrogen from methylenesulfate.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Notice: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling if this product not in accordance with directions.

SMG Solid and Guaranteed by

The Scotts Company  
14111 Scottsblaw Rd.  
Mansville, Ohio 43041

©2012 The Scotts Company LLC.  
World rights reserved.

#### Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTT'S LAWN FOOD STEP 2 WEED CONTROL PLUS LAWN FOOD 97236151

120% SIZE  
12-digit UPC-A  
(non-suppressed)  
For Position Only

0 32247 23615 7  
PROOF OF PURCHASE  
VID-BB  
BG1884



# OUR BEST ANNUAL PROGRAM



## 4STEP<sup>TM</sup> PROGRAM



Improves Lawn's Ability to  
Absorb Water & Nutrients

Covers 5,000 SQ. FT

KEEP OUT OF REACH OF CHILDREN  
CAUTION

See back for additional precautionary statements.

ACTIVE INGREDIENTS:

2,4-D <sup>1</sup> .....	1.18%
Mecoprop-P <sup>1</sup> .....	0.59%
OTHER INGREDIENTS: .....	98.23%
Total .....	100.00%

<sup>1</sup>0.0118 lb's 2,4-D and equivalent per pound of product  
<sup>2</sup>0.0058 lb's MCP-P and equivalent per pound of product



# STEP<sup>®</sup> 2

## Weed Control plus Lawn Food

DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM,  
DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS.

EPA Reg No. 538-270 (5M)

Scotts Step 2 Weed Control plus Lawn Food





\*0.0118 lbs 2,4-D acid equivalent per pound of product  
\*\*0.0059 lbs MCPP-p acid equivalent per pound of product

28-0-3

NET WT. 14.63 lb (6.64 kg)

Available only at authorized Scotts

**LawnPro**®

Dealers



**4 STEP**™  
PROGRAM

DO NOT USE ON ST. AUGUSTINEGRASS, FLORATAM,  
DICHONDRA, LIPPIA, CARPETGRASS OR BENTGRASS LAWNS.



Apply in Late Spring:

APR | MAY | JUN


**STEP 2**

Weed Control plus Lawn Food

Apply in Late Spring: APR | MAY | JUN







# Now it's Easy for You to Have a Better Lawn



## Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® product listed.

## Simple and easy

No more guesswork. You simply apply the right product – one STEP® at a time.

## Guaranteed results

Each STEP® product is designed to the specific season to help give you a greener lawn.

## Florida Annual Fertilizer Applications

On Centipede lawns, do not exceed rate of 2 lbs of Nitrogen per 1,000 sq. ft. in the north and 3 lbs of Nitrogen per 1,000 sq. ft. in the central and south. On Bahia lawns, do not exceed rate of 3 lbs of Nitrogen per 1,000 sq. ft. in the north.

North Florida is north of Ocala. Central Florida is defined as south of Ocala to a line extending from Vero Beach to Tampa. South Florida includes the remaining southern portion of the state.

## SPREADER SETTINGS

Spreader Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader. Covers 5,000 sq. ft. This package applies 2.9 lbs of product per 1,000 sq. ft.

	Spreader Settings	Directions		Spreader Settings	Directions
Scott's® Broadcast Rotary Spreaders	3 1/2	Each pass should be within 5 feet of each other to ensure overlap.		Drop Spreaders	5 3/4
					Make sure spreader wheel completely overlaps previous wheel marks.



For additional spreader settings please call 1-800-543-TURF (8873)





## GET EXPERT ADVICE FOR YOUR LAWN NOW!



Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline:  
**1-800-543-TURF**  
(8873)



Visit our website:  
**www.scotts.com**



Well-fed lawns are more efficient at using available water.

Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn against heat, drought and other stresses.

## DIRECTIONS FOR USE *It is a violation of federal law to use this product in a manner inconsistent with its labeling.*

This product applies 1.5 pounds of 2,4-D per acre per application.

### APPLICATION

- Apply on a calm day.
- Make sure that the grass is wet from dew or sprinkling before applying. (This helps particles adhere to weed leaves to produce best results.)
- Set spreader to the setting indicated by the chart above before applying.
- Kills weeds including dandelion, white clover, chickweed, plantain and ragweed.
- Apply when daytime temperatures are between 60 and 90 degrees F.
- Apply only to grass. Do not allow granules or dust to drift onto flowers, shrubs, fruit or vegetables.
- Grass should not touch the bottom of a drop spreader. Tall grass should be mowed a day or two before application.
- Delay application on newly seeded, sodded or sprigged areas until new grass has been mowed four times.
- Wait at least one month after using this product before you seed, sod or plant grass sprigs.
- Not for use on St. Augustinegrass, Floratam, dichondra, lippia, carpetgrass or bentgrass lawns.
- On bahia or centipedegrass lawns, wait until after the second mowing of the year, but not before mid-March.
- In fall, **DO NOT** treat centipedegrass after Oct. 1, nor bahiagrass after Nov. 1.
- Wait at least 24 hours after application before you water or mow the lawn (Granules must remain on weed leaves for 24 hours to be effective.)

### USE RESTRICTIONS

- Do not allow product to contact people or pets directly or through drift. Keep people and pets off lawn during application. Do not allow people or pets to enter treated areas until dusts have settled.
- Do not apply more than twice per year.
- A minimum of 30 days between applications is required.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a dry place. Reseal opened bag by folding top down and securing. Store in original container in a secure area away from children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes and socks and chemical resistant gloves made of any waterproof material.

### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling poison control center or doctor or going for treatment.

### ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

EPA Reg. No. 538-270 EPA Est. No. 538-OH-1 †CAS #94-75-7 ††CAS #16484-77-8

STEP® 2 Weed Control Plus Lawn Food 28-0-3 F 643

### Guaranteed Analysis

Total nitrogen (N).....	28%
5.2% ammoniacal nitrogen	
12.7% urea nitrogen	
9.3% other water soluble nitrogen*	
0.8% water insoluble nitrogen*	
Soluble potash (K <sub>2</sub> O).....	3%
Sulfur (S) .....	7%
7.0% combined sulfur (S)	

Derived from: Methyleneureas, potassium sulfate, ammonium sulfate, urea.

\*Contains 7.5% slowly available nitrogen from methyleneureas.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Notice: To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling if this product not in accordance with directions.



Sold and Guaranteed by:  
The Scotts Company  
14111 Scottslawn Rd.  
Marysville, Ohio 43041

©2012 The Scotts Company LLC.  
World rights reserved.

### Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

SCOTTS® LAWN FOOD STEP® 2 WEED CONTROL PLUS LAWN FOOD 5M 23615-1

120% SIZE  
12-digit UPC-A  
(non-suppressed)  
For Position Only

0 32247 23615 7  
PROOF OF PURCHASE  
VID-BB  
BG1884