

**GOT PESTS, get...**

**REVENGE®**



# Horse & Stable Fly Spray

Ready to Use

*No oily residue*

**Protect horses, beef & dairy cattle, goats, sheep & swine from horse flies, stable flies and face flies**

**ACTIVE INGREDIENT:**

Permethrin (CAS No. 52645-53-1) ..0.50%

**OTHER INGREDIENTS** ..... 99.50%

**TOTAL** ..... 100.00%

EPA Reg No. 4-358 EPA Est No. 4-NY-1



**Keep Out Of Reach Of Children**

**CAUTION** (See back panel for additional Precautionary Statements)

Net Contents 128 FL. OZ. (3.784 L)

#46173 v7/29

TRUSTED SINCE 1926  
**BONIDE®**

*Keeps working  
up to 4 weeks*



GOT PESTS, get... REVENGE  
**Horse & Stable  
Fly Spray**

Ready to Use

- ✓ Insecticide for horses, cattle (beef & dairy), goats, sheep, and hogs
- ✓ Controls fleas and ticks on dogs for 35 days
- ✓ For use in poultry houses and on birds
- ✓ No oily residue

Store and transport in an upright position.  
EPA Reg. No. 4-358 EPA Est. No. 4-NY-1  
Buyers Guarantee Limited to Label Claims.  
©Bonide Products, LLC All Rights Reserved  
Manufactured By:  
©Bonide Products, LLC  
6301 Sutliff Road, Oriskany, NY 13424  
#46173 Gallon v9/25

BN



**READ ENTIRE LABEL BEFORE EACH USE.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For both indoor and outdoor use.

**SHAKE WELL BEFORE USING**

**Use Restrictions**

Except when applying to pets, companion animals or non-commercial livestock, do not enter or allow others to enter until sprays have dried. Except when applying to pets, companion animals or non-commercial livestock, do not apply this product in a way that will contact adults, children or pets, either directly or through drift. Avoid contamination of feed and foodstuffs. Remove pets and birds, and cover fish aquaria before use.

Affix trigger sprayer to container for application. Do not transfer to another container.

**SPRAYER DIRECTIONS**



- Remove sprayer from storage holster.
- Slide safety lock to the LOCK position.



- Unwind pickup tube on the sprayer.
- Remove cap and seal from bottle.
- Insert pickup tube in bottle and screw sprayer cap onto bottle.



- Slide safety lock to the ON position.
- Gently squeeze the trigger to dispense fluid. The sprayer may require several seconds for the fluid to prime.
- After use, return to LOCK position.

For travelers who want to protect their luggage from “picking up” listed insects such as bedbugs, spray the outer and inner surfaces of luggage with a light spray mist of this product. Allow the luggage to completely dry before use.

### **INDOOR APPLICATION**

For spot application only. Not for broadcast use. Do not use as a space spray indoors. Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.



Do not allow the product to enter any drain during or after application.

Do not allow to enter indoor or outdoor drains.

No permita la entrada a desagües internos o externos.

#### **ON COMPANION ANIMALS AND NON-COMMERCIAL LIVESTOCK**

To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes, and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control Stable Flies, Horse Flies, and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control Bedbugs and Mites in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**USE ONLY ON DOGS. DO NOT USE ON CATS.** Do not repeat treatment for seven (7) days. Do not use on puppies under (12) weeks. Consult a veterinarian before using this product on debilitated,

aged, pregnant, nursing or medicated animals. Application of product on pets must only be done indoors.

To control Fleas, Ticks, and Lice: Start spraying at the tail moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat as necessary.

In case of an emergency, call 1-800-548-2423 (Animal Poison Control Center).

## **SURFACE SPRAYING**

To control Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetle, Meal Worms, Grain Mites and Cadelles:

Direct the spray into hiding places, cracks, crevices, behind shelves and drawers. For silverfish, spray bookcase. For ants, spray trails, nests and points of entry.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor covering and baseboard, molding, and floor. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Allow all treated articles to dry thoroughly before use.

To control Fleas (adult and larvae) and Ticks, including Ticks that may carry Lyme disease (adult and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboard, molding, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

## OUTDOOR APPLICATION

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- 1) Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
- 2) Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch or lawn);
- 3) Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
- 4) Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application. Do not apply when windspeed is greater than 15 mph.

Spot treatments must not exceed two square feet in size (e.g., 2 ft x 1 ft or 4 ft x 0.5 ft.).

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

## **OUTDOOR APPLICATION CONT.**

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds. Do not

apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Do not apply or irrigate to the point of runoff.

**TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):**

Use this product as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each area listed below apply until surface is wet (approximately 1 gal/750-1000 sq. ft.).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around and the outside of buildings to help control ticks occurring there.

## **OUTDOOR APPLICATION CONT.**

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 to 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**TO CONTROL FIRE ANTS:** Apply 1 gallon as a gentle rain to each fire ant mound using a sprinkler can or other applicator. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65-80 degrees F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

**PESTS ON OUTSIDE SURFACES OF BUILDINGS:** Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 3 feet where pests are and may find entrance. Also, apply as a spot treatment to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

**HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FLYING MOTHS OUTDOORS:** For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and

## **OUTDOOR APPLICATION CONT.**

other places where these insects alight or congregate. Repeat as necessary.

### **ANTS, LONE STAR TICKS, CRICKETS, FLEAS AND EARWIGS OUTDOORS:**

For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat applications as infestations warrant and as reinfestations occur.

**WASPS, HORNETS, YELLOW JACKETS, BEES OUTDOORS:** Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

### **STORAGE AND DISPOSAL**

**Pesticide Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

**Pesticide Disposal:** If partly filled - Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for disposal options.

**Container Disposal:** If empty - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Wash thoroughly after handling and before smoking or eating. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface water. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

For Emergency Medical Treatment Information call SafetyCall® at 1-833-972-1102

For Spill, Leak, Fire or Accident Information call Chemtrec at 1-800-424-9300

To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

# Horse & Stable Fly Spray

Got pests...get

# REVENGE®

Ready to Use

**ACTIVE INGREDIENT:**

Permethrin (CAS No. #52645-53-1) ..... 0.50%

**OTHER INGREDIENTS:** ..... 99.50%

Total ..... 100%

## Keep Out Of Reach Of Children **CAUTION**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Wash thoroughly after handling and before smoking or eating.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

### **STORAGE AND DISPOSAL**

**Pesticide Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

**Pesticide Disposal:** If partly filled – Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for disposal options.

**Container Disposal:** If empty – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

See inside the booklet for additional Precautionary Statements, First Aid, and Storage and Disposal.

©Bonide Products, LLC All Rights Reserved

Manufactured By:

Bonide Products, LLC  
6301 Suttiff Road, Oriskany, NY 13424

EPA Reg. No. 4-358  
EPA Est. No. 4-NY-1

#46173 v8/25

**Net Contents: 128 FL OZ (3.785L)**

Lift & Peel Here