KILLS OVER 250 WEEDS

KILLS DOWN TO THE ROOT



CLOVER



DANDELION

RESEALABLE PACKAGING



PENNINGTON®

ULTRAGREEN®

WEED & FEED

30-0-4

QUICK GREENING
FEEDS FOR UP TO 3 MONTHS



CREATES THICK LUSH LAWNS



EASY APPLICATION



CONTAINS 5% IRON FOR A DEEP GREEN LAWN



100% SATISFACTION GUARANTEED OR YOUR MONEY BACK

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p acid	0.167%
Dicamba acid	0.071%
OTHER INGREDIENTS:	98.654%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9.
Isomer Specific by AOAC Methods

NET WEIGHT **12.5 LB** | COVERS UP TO **5,000 SQ. FT.**

5.7 KG | 464.5 M²

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.

MANTÉNGALO LEJOS DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN Vea las precauciones completas y las indicaciones para el uso, incluyendo la información sobre primeros auxilios y almacenaje y desecho en el panel posterior.

CUT HERE TO OPEN

PENNINGTON® ULTRAGREEN® WEED & FEED 30-0-4

PENNINGTON® ULTRAGREEN® WEED & FEED 30-0-4

- KILLS OVER 250 WEEDS INCLUDING HENBIT, DOLLAR WEED, WHITE CLOVER, PLANTAIN AND CHICKWEED
- TREATS NORTHERN AND SOUTHERN GRASSES
- HIGHER NITROGEN CONTENT FOR GREATER COVERAGE

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use on dichondra, carpetgrass, 'Floratam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
- Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Weeds:

Amaranth, purple
Amaranth, slender
Aster, heath
Aster, white heath
Aster, purple
Aster, white prairie
Artichoke
Austrian fieldcress
Bedstraw, Catchweed
Bedstraw, smooth
Bedstraw, Northern
Bedstraw, pineywoods
Beggarticks, hairy
Betony, Florida
Bindweed, Field
Bindweed, Hedge
Bitter Wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black Medic
Blood flower milkweed
Blue lettuce
Brassbuttons
Bristly oxtongue
Broomweed
Buckhorn
Bullnettle
Burclover
Burdock
Burweed, lawn (spurweed)
Buttercup, bulbous
Buttercup, creeping
Buttonweed, common (poorjoe)
Buttonweed, virginia
Carolina geranium (wild)
Carpetweed
Catnip
Catsear, spotted (false dandelion)
Chamber bitter
Chickweed, common
Chickweed, mouseear
Chickweed, sticky
Chicory
Cinquefoil
Clover, crimson
Clover, alyce
Clover, hop
Clover, rabbit foot
Clover, red
Clover, strawberry
Clover, yellow sweet
Clover, white sweet
Clover, white
Cockle
Cocklebur
Cocklebur, oriental
Cocklebur, spiny
Common mullein
Corn Spurry
Creeping Beggarweed
Creeping Jenny (creeping charlie)
Cupid shaving Brush
Cudweed
Daisy, English
Daisy, oxeye
Dandelion, common
Dandelion, false
Day flower
Deadnettle
Dichondra
Dock, broadleaf
Dock, curly
Dogbane
Dogfennel
Dollarweed (Pennywort)
Dove weed
Elderberry
Falseflax
False sunflower
Fiddleneck
Filaree, redstem
Filaree, whitestem
Fleabane, daisy (annual)
Fleabane, rough
Galinsoga, hairy
Galinsoga, smallflower
Goldenrod
Ground ivy
Groundsel
Gumweed
Hairy fleabane
Hawkweed, meadow
Hawkweed, mouseear
Hawkweed, orange
Hawkweed, yellow
Healall
Heartleaf drymary
Hemp
Henbit
Hoary cress
Horsenettle
Horseweed
Jimsonweed
Johnny-Jumpup violet
Knawel
Knotweed, prostrate
Kochia
Lambsquarters
Lespedeza
Mallow, alkali
Mallow, bristly
Mallow, common
Mallow, dwarf
Mallow, venice
Marcela
Matchweed
Mexicanweed
Moneywort
Morning glory, bigroot
Morning glory, ivyleaf
Morning glory, mexican
Morning glory, red
Morning glory, tall
Morning glory, wolly
Mugwort
Mustard, ball
Mustard, black
Mustard, blue
Mustard, hedge
Mustard, Indian
Mustard, tansy
Mustard, tumble
Mustard, white
Mustard, wormseed
Nettle, stinging (burning)
Nettle, tall
Nettle, wood
Parsley-piert
Parsnip
Pearlwort
Pennycress (frenchweed)
Pepperweed, perennial
Pepperweed, virginia (Peppergrass)
Pigweed, prostrate
Pigweed, redroot
Pigweed, russian
Pigweed, smooth
Pigweed, tumble
Pineapple weed
Plains coreopsis (tickseed)
Plantain, blackseed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Plantain, hoary
Plantain, narrowleaf
Plantain, slender
Plantain, woolly
Poison ivy
Poison oak
Pokeweed
Prairie Sunflower
Prickly lettuce (compass plant)
Prickly sida
Prostrate knotweed
Purslane, common
Pusley, brazil
Pusley, florida
Pusley, large flower
Ragweed, bur
Ragweed, common
Ragweed, lanceaf
Ragweed, western
Redstem filaree
Rough cinquefoil
Scarlet pimpernel
Shepardspurse
Smartweed, ladysthumb
Smartweed, pale
Smartweed, pennsylvania
Smooth chaff-flower
Smoothdock
Sorrel, red (sheep)
Sowthistle, annual
Sowthistle, spiny
Spanish needles
Speedwell, bi-lobed
Speedwell, common
Speedwell, corn
Speedwell, ivyleaf
Speedwell, persian
Speedwell, purslane
Speedwell, slender
Speedwell, snow
Speedwell, thymeleaf
Spiny amaranth
Spurge, spotted
Spurge, prostrate
Spurweed
Star of Bethlehem
Strawberry, India mock
Tansy ragwort
Tanweed
Thistle, blessed
Thistle, bull
Thistle, canada
Thistle, flodman (prairie)
Thistle, musk
Thistle, platte
Thistle, plumeless
Thistle, russian
Thistle, scotch
Thistle, wavyleaf
Thistle, yellow (horrible)
Trailing crownvetch
Velvetleaf
Vervain, blue
Vervain, hoary
Vervain, prostrate
Vervain, tall
Vetch, bird
Vetch, hairy
Vetch, milk
Vetch, narrowleaf
Vetch, wild
Virginia creeper
Western clematis
Western salsify
Wild aster
Wild buckwheat
Wild carrot
Wild four-o-clock
Wild garlic
Wild lettuce
Wild marigold
Wild mustard (yellow rocket)
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild violet
Woodsorrel, creeping
Woodsorrel, violet
Woodsorrel, yellow (Oxalis)
Woolly croton
Wormseed
Yarrow, common
Yarrow, western
Yarrow, yellow

INSTRUCTIONS:

Pennington® UltraGreen® Weed & Feed 30-0-4 is a combination of three proven weed killers -- 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit and spurge. In addition, a 30-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS:

- Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
- Reseeding or over seeding areas may occur three (3) weeks after the application of this product.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass (not 'Floratam'), centipedegrass, hybrid bermudagrass, and colonial bentgrass:

Apply 2.5 pounds of product per 1,000 square feet of established turfgrass. Do not apply more than 2.5 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 5 pounds of product per 1000 square feet per year.

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see Step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTINGS

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

Coverage – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

BAG SIZE (LBS)	APPLICATION RATE (LBS / 1,000 SQ. FT.)	TREATED AREA FOR THIS BAG (SQ.FT.)
12.5	2.5	5,000

SINGLE PASS DISTRIBUTION:

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions. To provide proper distribution, calibrate spreader before application.

- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

SPREADER	SETTING DELIVERS 2.5 LB / 1,000 SQ. FT.
SCOTTS® CLASSIC DROP SPREADER	5¾
SCOTTS® EDGEGUARD® DLX BROADCAST SPREADER	5¼
SCOTTS® MINI EDGEGUARD® DLX BROADCAST SPREADER	4½
STA-GREEN® BROADCAST SPREADER	2¾
EARTHWAY® EV-N-SPRED® BROADCAST SPREADER	12

STATE RESTRICTIONS:

- Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html
- DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN.
- Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 0.75 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.
- Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.
- Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.pennington.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY & DISCLAIMER: To the extent consistent with applicable law, Central Garden & Pet, Garden Division guarantees this fine product to the extent of the purchase price, only when used according to directions.

30-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	30%
30% Urea Nitrogen*	
Soluble Potash (K_2O)	4%
Calcium (Ca)	2%
Iron (Fe)	5%
0.003% Water Soluble Iron (Fe)	

Derived from Urea, Polymer Coated Urea, Iron Sucrate, Ground Limestone, and Muriate of Potash.

***10.56% Slowly Available urea nitrogen from polymer coated urea.**

MONEY BACK GUARANTEE: If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explanation for seeking a refund.

HERBICIDA Y NUTRIENTE PENNINGTON® ULTRAGREEN® DE 30-0-4

- ELIMINA MÁS DE 250 MALEZAS INCLUYENDO ORTIGA MANSA, OMBLIGO DE VENUS, TRÉBOL BLANCO, LLANTÉN Y ÁLCINE
- PARA EL TRATAMIENTO DE HIERBAS SUREÑAS Y NORTEÑAS
- MÁS CONTENIDO DE NITRÓGENO PARA MAYOR CUBRIMIENTO

¡DETÉNGASE! LEA PRIMERO TODA LA ETIQUETA. RESPETE TODAS LAS PRECAUCIONES Y SIGA DETENIDAMENTE LAS INDICACIONES.

PRECAUCIONES

Peligros para las personas y animales domésticos

PRECAUCIÓN: Causa irritación moderada de los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de manipularlo y antes de comer, beber, mascar chicle, consumir tabaco o usar el baño.

PRIMEROS AUXILIOS

En caso de contacto con los ojos:	• Mantenga el ojo abierto y enjuague lentamente con agua durante 15-20 minutos. Si usa lentes de contacto, retírelos después de los primeros 5 minutos y continúe enjuagándose. • Llame a un centro de control de envenenamiento o doctor para recibir consejos sobre el tratamiento.

Tenga a mano el recipiente o la etiqueta del producto cuando llame al centro de control de envenenamiento o doctor, o cuando acuda a recibir el tratamiento. También puede llamar al 1-800-265-0761 para recibir información sobre el tratamiento médico de emergencia.

PELIGROS AMBIENTALES

Este pesticida es tóxico para los peces e invertebrados acuáticos y puede afectar de manera adversa a las plantas para las que no está destinado. Para proteger el ambiente, no permita que el pesticida penetre o sea arrastrado hacia desagües para tormentas, zanjas de drenaje, canaletas o aguas de superficie. Aplique el producto en tiempo de calma cuando no se pronostique lluvia por las próximas 24 horas para evitar que la lluvia o el viento arrastren el pesticida fuera del área de tratamiento. Barra hacia el área de tratamiento el producto que caiga en la entrada para carros, la acera o la calle para evitar que sea arrastrado hacia cuerpos de agua o sistemas de drenaje.

Este producto químico tiene propiedades y características asociadas con sustancias químicas detectadas en las aguas subterráneas. El uso de este producto en áreas con suelos permeables, particularmente donde el nivel acuífero o freático es poco profundo, puede ocasionar la contaminación de las aguas subterráneas. La aplicación cerca de cisternas o pozos puede ocasionar la contaminación del agua potable o el agua subterránea.

INDICACIONES DE USO

El uso de este producto de manera diferente a lo indicado en su etiqueta es una violación de las leyes federales.

RESTRICCIONES DE USO:

- No aplique este producto de manera que entre en contacto con personas o animales, ya sea directa o indirectamente. Evite la entrada al área de personas y mascotas durante la aplicación y no permita que entren al área tratada hasta que el polvo se haya asentado.
- No lo use en céspedes cultivados para la venta u otro uso comercial como los tepes, o para producción comercial de semillas ni para fines investigativos.
- No lo use en golondrina, zacate amargo, hierba de San Agustín 'Floratam', nombre de dios ni tepes donde desee la presencia de trébol.
- No use este producto en plantas de flores, vegetales, cobertura de terreno, huertos ni jardines, incluyendo arbusto y árboles. No exceda la dosis especificada para cualquier área y no lo aplique dentro del área de goteo próxima al tronco o sobre las raíces de los árboles, arbustos y otras plantas ornamentales.
- No use este producto en o cerca de plantas deseables y evite esparcir los gránulos sobre los sistemas de raíces expuestas o brotes adventicios (rizomas) dentro del área de goteo de las raíces de árboles y arbustos deseables ya que puede ocasionar daños a la planta.
- No lo aplique directamente a o cerca del agua, desagües para tormentas, canaletas, alcantarillas o zanjas de drenaje. No lo aplique a menos de 25 pies (7,62 m) de los ríos, estanques de peces, lagos, arroyos, reservas de agua, pantanos, estuarios, bahías ni océanos. No lo aplique en viento. Aplique este producto directamente a su césped y barra cualquier producto que caiga sobre la entrada para carros, acera, canaleta o calle hacia el área tratada. Para evitar que el producto sea arrastrado, no riegue el área tratada de manera que el agua pueda correr ni lo aplique tampoco cuando esté lloviendo o se espere lluvia ese día.

INSTRUCCIONES:

El herbicida y nutriente Pennington® UltraGreen® de 30-0-4 es una combinación de tres herbicidas: 2,4-D, Mecoprop-p y Dicamba. Este herbicida controla una amplia gama de malezas del césped, incluyendo el diente de león, álcine, sanguinaria, llantén, ortiga mansa y golondrina de jardín. Además, el fertilizante de 30-0-4 brinda abundantes nutrientes para las plantas.

PRECAUCIONES DE USO:

- No aplique este producto a las hierbas recién sembradas hasta que se asienten bien o hasta después de la segunda o la tercera poda.
- La resiembra o sobresiembra de semillas en el área puede hacerse tres (3) semanas después de la aplicación de este producto.

CUÁNTO APLICAR:

Para pasto poa, festuca, ballico perenne, hierba Bermuda común, hierba bahía, zoysia, hierba de San Agustín (no 'Floratam'), ciempiés, híbridos de hierba Bermuda y hierba fina:

Aplique 2,5 libras del producto por 1000 pies cuadrados de césped asentado. No aplique más de 2,5 libras de producto por 1000 pies cuadrados de césped por aplicación. Si es necesario, se puede realizar una segunda aplicación a no menos de 30 días después de la primera. No aplique más de 5 libras de producto por 1000 pies cuadrados por año.

CUÁNDO APLICARLO:

Aplíquelo cuando las malezas de hojas anchas estén creciendo activamente en la primavera o comienzos del otoño.

Aplíquelo cuando el viento esté calmado para asegurar una cobertura uniforme y evitar que los gránulos se esparzan hacia flores, vegetales y arbustos ornamentales.

Es posible que sea necesario repetir el tratamiento si el césped está invadido por las malezas, si prevalecen las condiciones adversas de poca humedad o si las malezas están en un estado de poco crecimiento.

INSTRUCCIONES PARA LA APLICACIÓN:

1. Pode el césped a una altura normal 1 o 2 días antes de la aplicación.
2. Riegue el césped abundantemente al menos 1 o 2 días antes de la aplicación para mantener la humedad hasta la próxima sesión de riego (vea el paso 5).
3. Aplíquelo cuando las malezas de hojas anchas estén jóvenes y creciendo activamente, preferiblemente en la mañana cuando el rocío esté aún sobre la hierba. En el momento de la aplicación, la humedad del rocío, la lluvia o el riego sobre las hojas de la maleza ayudará al control. Si la hierba no está húmeda al momento de la aplicación, rocíela ligeramente con agua antes de aplicar el producto.
4. Evite aplicarlo cuando las temperaturas no sean consistentemente superiores a los 90°F (32°C).
5. No permita que el agua retire el producto de las hojas de la maleza por 1 o 2 días después de la aplicación. Solo entonces, se debe aplicar una sesión de riego abundante.

AJUSTES DEL ESPARCIDOR

Aplíquelo siempre con un esparcidor giratorio o por goteo para distribuir los gránulos uniformemente. No lo aplique nunca a mano.

Cobertura – Para determinar la cantidad que necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale al total de pies cuadrados. No incluya las áreas sin césped (casa, jardín, etc.). Una bolsa cubre hasta 5000 pies cuadrados.

IMPORTANTE: Es importante que cumpla con las indicaciones que se incluyen con este producto de calidad para el cuidado del césped, ya que una aplicación excesiva puede dañar el césped. Los ajustes del esparcidor son aproximados y puede haber diferencias debido a las condiciones del esparcidor, la velocidad con la que lo aplique y las irregularidades del área tratada. Asegúrese de revisar el esparcidor antes de comenzar y ajustarlo si fuera necesario. Consulte el manual del propietario. Nunca lo aplique a mano.

TAMAÑO DE LA BOLSA (LBS)	TASA DE APLICACIÓN (LBS / 1000 pies²)	ÁREA TRATADA POR UNA BOLSA DE ESTE TAMAÑO (pies²)
12,5	2,5	5,000

DISTRIBUCIÓN CON UN SOLO PASE:

- Para esparcidores giratorios: Trabaje en intervalos correspondientes con el ancho de la franja de tierra.
- Para esparcidores por goteo: Solape la trayectoria de las ruedas para garantizar una cobertura completa.
- Para lograr mejores resultados, aplique el producto uniformemente y trate de no dejar franjas sin cubrir, saltarse espacios o aplicar dos veces en una misma área. Camine con un paso estable para asegurar una cobertura uniforme. Cierre la abertura del esparcidor cuando gire para cambiar de dirección o cuando se detenga.

INSTRUCCIONES PARA LA CALIBRACIÓN Y AJUSTES DEL ESPARCIDOR:

Calibre el esparcidor según las instrucciones del fabricante. El ajuste inicial del esparcidor pudiera requerir ser regulado para aplicar la tasa recomendada bajo condiciones reales de aplicación. Para garantizar una distribución apropiada, calibre su esparcidor antes de la aplicación.

- Para 2,5 libras de producto por 1000 pies cuadrados de césped: Verifique su esparcidor en 400 pies cuadrados y ajústelo para aplicar a una tasa de 1 libra de producto por 400 pies cuadrados.

ESPARCIDORES POR DISPERSIÓN: Para obtener mejores resultados, aplique en trayectoria circular empezando desde afuera y moviéndose hacia el centro superponiendo ligeramente para asegurar un cubrimiento completo.
ESPARCIDORES POR GOTEO: Aplique 2 franjas a lo largo de cada extremo. Luego aplique a lo largo de la longitud del césped superponiendo ligeramente y cerrando el esparcidor antes de cada giro al borde de las franjas.

ESPARCIDOR	AJUSTE PARA APLICAR 2,5 LB / 1000 pies²
SCOTTS® CLASSIC DROP SPREADER	5¾
SCOTTS® EDGEGUARD® DLX BROADCAST SPREADER	5¼
SCOTTS® MINI EDGEGUARD® DLX BROADCAST SPREADER	4½
STA-GREEN® BROADCAST SPREADER	2¾
EARTHWAY® EV-N-SPRED® BROADCAST SPREADER	12

RESTRICCIONES ESTATALES:

- La información acerca del contenido y los niveles de metales en este producto está disponible por internet en http://www.aapfco.org/metals.html
- NO LO APLIQUE CERCA DEL AGUA, DESAGÜES PARA TORMENTAS NI ZANJAS DE DRENAJE. NO LO APLIQUE CUANDO SE ESPERE LLUVIA FUERTE. APLIQUE ESTE PRODUCTO EN EL CÉSPED SOLAMENTE Y BARRA HACIA EL CÉSPED/JARDÍN EL PRODUCTO QUE CAIGA SOBRE LA ENTRADA PARA CARROS, LA ACERA.
- Aleje el producto de las aceras, patios, entradas para carros, etc. Este fertilizante contiene hierro que puede manchar las superficies pintadas y de concreto. Si entra en contacto con la superficie de concreto, bárralo inmediatamente antes de limpiar con agua. Si el concreto está mojado cuando se produce el contacto, puede que resulte imposible impedir que se manche. Tenga cuidado al usar esparcidores giratorios que esparcen el fertilizante más allá del borde del césped y hacia el área con concreto.
- Generalmente, un césped consume de 1 a 4 libras de nitrógeno por cada 1000 pies cuadrados por cada temporada de crecimiento. Cuando se aplica como se indica, este producto brinda 0,75 libras de nitrógeno por cada 1000 pies cuadrados así como herbicida para controlar las malezas mencionadas. Comuníquese por favor con el servicio de extensión local para recibir información sobre el programa regular de fertilización.
- Consulte con el Servicio de Extensión Cooperativa de su estado para conocer la información actualizada sobre los límites anuales de aplicación de fertilizantes de césped en su región.
- La información acerca de las materias primas que componen este producto se puede obtener escribiendo al avalador mencionado en esta etiqueta y refiriendo el número de lote que se encuentra en este recipiente.

ALMACENAJE Y DESECHO

No contamine el agua, los alimentos ni el pienso al almacenar y desechar el producto.

ALMACENAJE DE PESTICIDAS: Guárdelo en el recipiente original y mantenga las bolsas que contienen el producto sin usar en un área de almacenaje cerrada e inaccesible para niños y animales domésticos.

DESECHO DE PESTICIDAS Y MANIPULACIÓN DEL RECIPIENTE: Este recipiente no es rellenable. No lo reutilice ni lo rellene. **Si está vacío:** Tírelo a la basura o recíclelo si el servicio está disponible. **Si está parcialmente lleno:** Llame a su agencia local para el desecho de sustancias sólidas para recibir instrucciones en cuanto al desecho. No arroje nunca el producto sin usar en inodoros ni desagües interiores o exteriores.

GARANTÍA Y RENUNCIA DE RESPONSABILIDAD: En consonancia con lo estipulado por la ley correspondiente, Central Garden & Pet y la División de Jardinería garantizan este producto de calidad hasta la cantidad del precio de compra y solo cuando se usa como se indica.

30-0-4

ANÁLISIS GARANTIZADO

Total de nitrógeno (N)	30%
30% de nitrógeno ureico*	
Potasa soluble (K_2O)	4%
Calcio (Ca)	2%
Hierro (Fe)	5%
0,003% de hierro soluble en agua (Fe)	

Derivado de la urea, urea con recubrimiento de polímero, sucrato de hierro, cal natural y muriato de potasa.

***10,56% de nitrógeno ureico de liberación lenta a partir de la urea con recubrimiento de polímero.**

GARANTÍA DE DEVOLUCIÓN DE DINERO: Si no está completamente satisfecho con este producto, por favor envíe por correo su nombre, dirección, prueba de la compra (paquete CUP) y una breve explicación especificando que busca un reembolso.

CENTRAL Garden & Pet

Guaranteed & Distributed by
CENTRAL GARDEN & PET GARDEN DIVISION
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-884-90098
EPA Est. No. 59144-GA-1_A, 32802-MO-1_B, 37351-CO-1_C, 69759-OR-004_D, 41124-WI-1_E, 41124-OH-1_F, 9404-FL-1_G
Subscript used is the First letter of Run Code on Container

WWW.PENNINGTON.COM
1-800-265-0761 | GARDENDIVISION@CENTRAL.COM

Pennington and Pennington with design are registered trademarks of Pennington Seed, Inc. Central Garden & Pet, Central Garden & Pet with design and UltraGreen are trademarks of Central Garden & Pet Company. Scotts and Edgeguard are registered trademarks of OMS Investments, Inc. Sta-Green is a registered trademark of LF, LLC. Earthway and Ev-N-Spred are registered trademarks of Earthway Products, Inc.

©2018, 2019 Central Garden & Pet Company

0 21496 01994 9

UG237_V4 | 300527377 - 04/19