

# LESCO DIMENSION\* 0.21% Plus Fertilizer

**NOT FOR USE ON** turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

In New York State this product may only be used by commercial applicators and at no more than 500 lb (0.5 lb of active ingredient) per acre per year (or 1.5 lb product per 1,000 sq ft per year) and is prohibited from use in Nassau and Suffolk Counties.

<b>ACTIVE INGREDIENT:</b>	2-(dimethylamino)-4-(2-methylpropyl)-6-(methylamino)-3,5-dimethylcyclohexene
Dithiofou:	5.5% (5.5 lb/100 lb)
-3.5% (3.5 lb/100 lb)	0.21%
<b>INERT INGREDIENTS:</b>	88.78%
Total:	100.00%

Product protected by U.S. Patent No. 4,892,184. Other patents pending.  
Read the entire label before using this product. Use only according to label instructions.

**NOTICE:** Before using this product, read the Use Precautions, Warranty Statements, Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### FIRST AID

- If swallowed:
  - Call a poison control center or doctor immediately for treatment advice.
  - Have person sip 1 glass of water if able to swallow.
  - Do not induce vomiting; unless told by the poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- If in eyes:
  - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
  - Call a poison control center or doctor for treatment advice.
- If on skin or clothing:
  - Take off contaminated clothes.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- Inhalation:
  - Airway obstruction.
  - If person not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION!** CAUSE EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wear protective eyewear such as goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic areas. Do not apply directly to water, or to areas where surface water is present, or to areas where water below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

#### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses, and broadleaf weeds in established lawns and ornamental lawns, including golf courses, lawns, meadows, and weedy areas. This product will not harm newly established ornamentals when used according to label directions.

The product may be used on pastures, sodded, or sprigged lawns and ornamental lawns that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive application of this product. Use of this product on lawns and ornamental lawns that are not well-established, or on those that have been maintained by weeding, hand-weeding, hand-tilling, or cultural methods, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental lawns that are free of the following annual species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established subspecies are tolerant to the product:

#### Crabgrass, Grasses

Common Name	Scientific Name
Bermuda, couch*	Agrostis pallens
Bluegrass, Kentucky	Poa pratensis
Fescue, blue*	Festuca rubra
Fescue, tall	Festuca arundinacea
Fescue, perennial	Festuca rubra

Warm-Season Grasses	
Common Name	Scientific Name
Bermudagrass	Paspalum notatum
Bermudagrass	Cynodon dactylon
Buffalograss	Buffalo grass
Leymus	Leymus
Perennial ryegrass	Lolium perenne
Tall fescue	Festuca arundinacea
St. Augustinegrass	Zoysia japonica
Zoysia grass	Zoysia

DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.  
Use of this product on certain varieties of Creeping Bentgrass, such as 'Coloray', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of Creeping Bentgrass have been tested.  
Use of this product on certain varieties of Fine Fescue may result in undesirable turf injury. The following Fine Fescue varieties are known to be sensitive to this product: 'Adonis', 'Barber', 'Beauty', 'Bolero', 'CF-2', 'Endy', 'F-1', 'F-2', 'Highgate', 'Twink', 'Aegeanstar', 'Majestic', 'Mary', 'Penitentiary', 'Tanner', 'Tropicana', 'Whidbey', and 'Whitewell'. Not all varieties of Fine Fescue have been tested.  
DO NOT use this product on seedling Bahiagrass in the spring of the first year of establishment until the turfgrass is fully green and has established itself.

#### DIMENSION TURFGRASS REGIONS



#### Seedling, Overseeding, or Sprigging

Reseeding, overseeding or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practice such as soil cultivation, irrigation and fertilization should be followed.

However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeder equipment (if seeders) designed to give good seed to soil contact.

#### USE PRECAUTIONS

- Not for use on Golf Course Putting Greens.
- The early post-emergence crabgrass control described below is limited. Post-emergence applicators are only effective on crabgrass if applied before crabgrass has displayed its tiller (leaf or first tiller). For best results, cultural practices that destroy the soil, such as core, spike, or hydro-aerification, and fertilizing, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.
- Apply the product directly in established lawns or ornamental lawns only.
- DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.

#### PROGRAM

**Program 1** - pre-emergence (fall > 2.5 in.)  
**Program 2** - Post-emergence (fall > 1 in.)  
**Program 3** - Post-emergence (fall > 1 in.)

#### USE

**Program 1** - 1.15 lb/1,000 sq ft  
**Program 2** - 2.72 lb/1,000 sq ft  
**Program 3** - 4.54 lb/1,000 sq ft  
**Control Souls** - 4.64 lb/1,000 sq ft

#### Rate

Rate: 2.72 lb/1,000 sq ft is equal to 2.25 lb/1,000 sq ft per acre.

1. Paricle distribution at this rate may not provide adequate control or suppression.

\*DO NOT apply more than 5.48 lb of this product per 1,000 sq ft per application, and no more than 16.38 lb of this product per 1,000 sq ft per year.

\*\*Program 1 is not for use on golf courses.

Subsequent required pre- and/or post-emergence applications should be made where longer periods of control are desired.

When split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

#### APPLICATION DIRECTIONS - Control of Crabgrass

##### Pre-emergence and Early Post-emergence Control

This product provides "pre-emergence" control of crabgrass from the ground in established lawns and ornamental lawns. It can also provide "early post-emergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental sites. Post-emergence crabgrass control will be obtained only when the product is applied prior to the life stage of crabgrass, which corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the potential benefit of this product's additional, early post-emergence activity is that it (consequently) is slightly pre-emergence crabgrass protocol. Dimension controls crabgrass prior to and up to 4 weeks after germination.

##### Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 5.48 lb of this product per 1,000 sq ft per application, and no more than 16.38 lb of this product per 1,000 sq ft per year.

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the control location, the biology, growing weight, and whether the site is considered to be pre-emergence or early post-emergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

The Program 1 pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-6 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabgrass control.

The Program 2 pre-emergence control at sites where (a) Kentucky is cut low (e.g., golf courses) and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

The Program 3 pre-emergence control at sites where (a) Kentucky is cut low (e.g., golf courses) and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. The program may also be used for early post-emergence control at sites where Kentucky is cut low (e.g., golf courses).

Subsequent required pre- and/or post-emergence applications should be made where longer periods of control are desired.

When split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

##### Table 1 Recommended Single Application Use Rates\*

PROGRAM	I	II	III
(Bermuda cut)	Pre-emergence (fall > 2.5 in.)	Post-emergence (fall > 1 in.)	Post-emergence (fall > 1 in.)
Region	1.15 lb/1,000 sq ft	2.72 lb/1,000 sq ft	4.54 lb/1,000 sq ft
North	1.15 lb/1,000 sq ft	2.72 lb/1,000 sq ft	4.54 lb/1,000 sq ft
South	2.72 lb/1,000 sq ft	5.48 lb/1,000 sq ft	9.96 lb/1,000 sq ft
Central/Souls	4.64 lb/1,000 sq ft	10.96 lb/1,000 sq ft	19.92 lb/1,000 sq ft

Note: 2.72 lb/1,000 sq ft is equal to 2.25 lb/1,000 sq ft per acre.

1. Paricle distribution at this rate may not provide adequate control or suppression.

\*DO NOT apply more than 5.48 lb of this product per 1,000 sq ft per application, and no more than 16.38 lb of this product per 1,000 sq ft per year.

\*\*Program 1 is not for use on golf courses.

\*\*Program 2 includes Kentucky, NC, SC, TN, GA, AL, MS, AR, LA, OK, TX, and FL.

\*\*Program 3 includes KY, NC, SC, TN, GA, AL, MS, AR, LA, OK, TX, and FL.

\*\*Program 4 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 5 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 6 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 7 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 8 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 9 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 10 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 11 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 12 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 13 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 14 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 15 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 16 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 17 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 18 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 19 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 20 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 21 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 22 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 23 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 24 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 25 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 26 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 27 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 28 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 29 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 30 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 31 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 32 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 33 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 34 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 35 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 36 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 37 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 38 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 39 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 40 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 41 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 42 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 43 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 44 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 45 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 46 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 47 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 48 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 49 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 50 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 51 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 52 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 53 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 54 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 55 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 56 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 57 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 58 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 59 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 60 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 61 includes KS, MO, IL, WI, MI, VA, southern areas of DE, MD, NJ, NY, CT, RI, MA, NH, VT, and ME.

\*\*Program 62 includes KS, MO, IL, WI, MI, VA,