228-432-32802_HJE All Season Weed & Feed with Lazer 18-0-3_20230127_25_32802_.pdf



18

3

PERSIST
For short grass

PERSIST
Wait for growth to reach recommended

Guaranteed Analysis

Total Nitrogen (N)	75.00%
0.00% Water Insoluble Nitrogen (WIN)	
Soluble Potash (K2O)	7.00%
Derived From: Muriate of Potash, Urea	
Calories (Cl), not more than 0.20%	P2O5

Fertilizer guaranteed by Howard Johnson's Enterprises, Inc.

Information regarding the contents and kinds of metals in this product is available on the Internet at https://www.epa.gov/metals.htm

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Wash hands before eating, drinking, chewing gum or using tobacco.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Do not apply this product in rain weather when rain is not predicted for the next 24 hours as it helps to ensure that wind or rain does not blow or wash pesticide off the treatment area. Avoid applying when weather conditions favor runoff from heavy rain back onto the treated area of the lawn as it helps to prevent run off to water bodies or drainage systems. Do not contaminate water when disposing of equipment wash waters or rinsate.

FIRST AID

IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-425-1840 for emergency medical treatment information.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is a ready-to-use mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Bpx Grass, Centipede, Fescue, Rye, St. Augustine, Tall Fescue, Zoysia, Kikuyu, Seashore Paspalum and Buffalo grass. Do not use this product where desirable plants are present or allow the product to "wick" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

GENERAL PRECAUTIONS AND RESTRICTIONS: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

Do not apply more than 3 applications per year, including broadcast applications in combination with spot treatments.

MIXED CONTROLLED: This product is designed to provide fast green growth of bluets and control many broadleaf weeds such as: Alsike, Annual yellow rocket clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Bindweed, Black medic, Blackberry, Blessed thistle, Bracted plantain, Buckwheat species, Buffalo medic, Black medicine, Burweed species, Burweed species, Buttercup, Blue vervain, Blue dicks, Bramble, Brainbur, Brainbur, Brainbur, Broadcastbur, Bristly starbur, Broadleaf dock, Broadleaf plantain, Broadleaf signalgrass, Broadleaf species, Broadleaf species, Broadleaf species, Broadleaf species, Burclover, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Common chickweed, Common cocklebur, Common dandelion, Common groundsel, Common lambsquarters, Common mallow, Common mullein, Common plantain, Common ragweed, Corn chamomile, Creeping jenny, Crimson clover, Curlycup gumweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogbane, Dooryard, Dooryard, English daisy, Fall panicum, False dandelion, False flax, False sunflower, Field bindweed, Field pansy, Five leaf grass, Flixweed, Florida betony, Florida pusley, Forget-me-not, Garlic mustard, Goatsbeard, Goldenrod, Goldenrod species, Ground ivy, Gumweed, Hairy bittercress, Hairy buttercup, Hairy galinsoga, Hairy vetch, Hawkweed species, Hedge bindweed, Hedge mustard, Hemp, Henbit, Horse nettle, Hoary alyssum, Hoary vervain, Horsenettle, Hop clover, Horseweed, Horsetail, Indian mallow, Ironweed, Jerusalem goldenrod, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Longbeak buttercup, Lyre leaf sage, Mallow species, Marestail, Matchweed, Mayweed chamomile, Meadow buttercup, Mexican clover, Mexican fireweed, Mock mallow, Morningglory, Mouseear chickweed, Musk thistle, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Pale pea vetch, Parsley piert, Partridge pea, Pennsylvania smartweed, Peppergrass species, Perennial ryegrass, Pigweed species, Pineapple weed, Pitseed goosefoot, Plantain species, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Prairie, Prairie sunflower, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purple deadnettle, Purple loosestrife, Purslane, Quackgrass, Quackgrass, Quackgrass, Rough cinquefoil, Rough fleabane, Russian thistle, Russian thistle, Russian pigweed, Russian thistle, Sa, Sandvetch, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Shetleaf clover, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Smooth vetch, Southern wild rose, Southern sandbur, Spanishneedles, Speedwell, Spiny Amaranth, Spiny cocklebur, Spotted knapweed, Spotted spurge, Spurge species, Spurge species, Spurge species, Stinkweed, Stinkweed, Strawberry clover, Sumac, Sunflower species, Tall buttercup, Tall coreopsis, Tansy mustard, Tansy ragweed, Tarweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing crownvetch, Tumble pigweed, Tumble windmillgrass, Tumble windmillgrass, Tumble windmillgrass, Tumble windmillgrass, Vaseyi lupine, Velvetleaf, Virginia buttercup, Virginia creeper, Virginia pepperweed, Waterleaf, Wavyleaf, Western dichondra, Western salsify, White clover, White mustard, Wild mustard, Wild radish, Wild buckwheat, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild parsley, Wild parsnip, Wild pansy, Wild radish, Wild rose, Wild strawberry, Wild sweet potato, Wild vetch, Wild violet, Windmillgrass, Woodland, Wood sorrel, Wood violet, Woolly croton, Wormseed, Yarrow, Yellow rocket, Yellow sweetclover, Yellowflower pepperweed and other broadleaf weeds.

When to Apply

Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers exposed when walking with a spray-dampened type spreader. Do not exceed for at least 3 to 4 weeks after application. Do not apply to newly seeded areas until grass has been established for at least 3 months. Do not apply to newly sodded areas after application. Do not apply when grass is moist after a rain, sprinkling or dew. To avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weeds. Keep treated and do not exceed applying if rainfall is expected within 48 hours following watering.

Possible retreatment is approximately 30 days may be necessary if lawn is heavily overun with weeds or if environmental conditions restrict new conditions prevent and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

Do not apply to St. Augustine grass, Bahia grass, Centipede grass, Zoysiagrass or inadequate moisture, high temperature and high humidity, overapplication may cause moderate transient discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

How to Apply

This 1K pound bag should be used on 1,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application using the area at half the rate each time also is acceptable.

Use a drop spreader or rotary spreader. Fill hopper to no more than ½ full and operate spreader while walking forward at a steady pace (avoiding jogging or running). Use the following settings as a guide. Always calibrate your spreader before applying. Settings are based on an average walking speed. Reduce setting for slower speed. Increase setting for faster speeds. Check your spreader at 250 square feet and adjust to apply at the rate of 1LB pound per 250 square feet.

PATH A

Drop

First apply two strips of entire Feet apply Length of the lawn, overlapping very slightly. Then apply by walking perpendicular to the previous strips running.

PATH B

Rotary

Apply in a circular path, starting on the outside and working toward the center overlapping very slightly.

Spreader Settings

Broadcast Spreader		Drop Spreader	
Agway Co.	A2	Agway Co.	A2
Ames	A2	Ames	2.5
Ames Deluxe Dirt	A2	Ames Deluxe Dirt	2.5
Craftsman	A2	Craftsman	5.5
Earthway	11-D	Earthway	N/A
Gilson Home	N/A	Gilson Home	7.7
Precision GreenLine	5.75	Precision GreenLine	7.5
Precision Red Devil	5.75	Precision Red Devil	7.5
Precision Pro	5.75	Precision Pro	7.5
Scotts Speedy Green	5.75	Scotts Aero Green	6.5
Scotts Easy Green	5.5	Scotts Even Green	7.5
Scotts Basic	5	Scotts	7.7
Scotts Lawn Pro	5		
Scotts Deluxe EdgeGuard	5		
Solo	2.0		

Sweep on not watch all lights over materials from the spreader only after calibration, because certain fertilizer ingredients may cause permanent stains. Wash out spreader after using. Use a hose to dry these weights. Treat any exposed metal spreader areas with rust to reduce corrosion.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container to store pesticides and keep any bags containing unused product in a locked storage area. Remember to keep out of reach and access to children. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Empty: Place in trash or offer for recycling if available. Empty bags: Dispose in trash. Include in state agency for disposal instructions. Never allow children to play in or reuse in drinking, food or cooking. Never clean.

WARRANTY DISCLAIMER

Limitation of Liability

If you do not agree with or do not accept any of directions for use, the warranty disclaimer, or limitations on liability, do not use the product, and return it unopened to the Seller and the purchase price will be refunded.

LOT: 0000 EXP: 00/00/0000 MFG: 00/00/0000

Distributed By:

Howard Johnson's Enterprises, Inc.
9575 S 68th Street, Franklin, WI 53132

© Copyright 2021
Howard Johnson's Enterprises, Inc.