



All-Season

CRABGRASS CONTROL PLUS

With Prodiamine

Fertilizer that builds thick, green lawns. Stops many weed grasses before they grow.




| | |
|------------------------------|---------|
| ACTIVE INGREDIENTS: | |
| Prodiamine: (CAS 29091-21-2) | 0.29% |
| INERT INGREDIENTS: | 99.71% |
| Total | 100.00% |

EPA Reg. No. 32802-70
 EPA Est. No. 59144-GA-1 (A),
 EPA Est. No. 32802-MO-1 (B),
 EPA Est. No. 37351-CO-1 (C),
 EPA Est. No. 8378-IN-1 (D),
 EPA Est. No. 9404-FL-1 (E)
 First letter of bag code shows establishment where manufactured.

16·0·4

Covers 5,000 sq. ft. Net Weight: 15 lbs. (6.8 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR FURTHER PRECAUTIONS AND FIRST AID INFORMATION.



Guaranteed Analysis

| | |
|---|--------|
| Total Nitrogen (N) | 16.00% |
| 16.00% Urea Nitrogen | |
| Soluble Potash (K ₂ O) | 4.00% |
| Derived From: Muriate of Potash, Urea. | |
| Chlorine (Cl), not more than 6.00% | F315 |

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow this product off the treated area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

FIRST AID

| | |
|-------------------------------|--|
| If in eyes | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person. |
| If inhaled | <ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |

Hotline number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

General information

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens). It controls susceptible weeds by inhibiting weed seed germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed);

jungerlice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass²; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹ In many areas where not prohibited by fertilizer use restrictions³ a single application of 5.2 to 11.9 pounds / 1000 square feet (225 to 519 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application to 5.2 to 7.9 pounds / 1000 square feet (225 to 345 pounds/acre) to areas not prohibited by fertilizer use restrictions, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

² Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.

³ For this weed, apply in late summer, fall or winter prior to germination.

⁴ Suppression only.

When And How To Apply

WHEN TO APPLY TO ESTABLISHED TURF:

This product must be applied prior to germination of weed seed, as it will not control established weeds.


APPLICATION DIRECTIONS:

Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street back onto your lawn. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)

| Turf Species | Minimum Rate | Maximum Annual Rate ¹ |
|---|-----------------------|----------------------------------|
| Bahiagrass ² ; Bermudagrass ³ | 2.75 lbs / 1000 sq ft | 11.9 lbs / 1000 sq ft per year |
| Centipedegrass ² ; Zoysiagrass | | |
| Seashore Paspalum | | |
| St. Augustinegrass | | |
| Tall Fescue (including turf-type) | | |
| Buffalograss | 2.75 lbs / 1000 sq ft | 7.9 lbs / 1000 sq ft per year |
| Kentucky Bluegrass | | |
| Perennial Ryegrass | | |
| Creeping Red Fescue | 2.75 lbs / 1000 sq ft | 6 lbs / 1000 sq ft per year |
| Creeping Bentgrass | 2.75 lbs / 1000 sq ft | 5.2 lbs / 1000 sq ft per year |

¹ These are the maximum rates per calendar year.

² In Florida: Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the North. Do not exceed 3 applications per year to Centipedegrass in the Central and South regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.

³ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 4.1 pounds / 1000 square feet. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION:

This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseed results.

| USE RATE | MONTHS BEFORE OVERSEEDING | | |
|--------------------------|---------------------------|------------|-------|
| | North | Transition | South |
| 2.75 pounds / 1000 sq ft | 4 | 4 | 4 |
| 5.2 pounds / 1000 sq ft | 5 | 4 | 4 |
| 6 pounds / 1000 sq ft | 6 | 5 | 5 |
| 6.3 pounds / 1000 sq ft | --- | 6 | 6 |
| 7.9 pounds / 1000 sq ft | --- | 7 | 7 |
| 9 pounds / 1000 sq ft | --- | --- | 9 |
| 10.3 pounds / 1000 sq ft | --- | --- | 10 |
| 11.9 pounds / 1000 sq ft | --- | --- | 12 |

Pesticide Storage

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

IF EMPTY: Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available.

IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

WARRANTY: The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc. the manufacturer, or the seller.

To the extent consistent with applicable law, the products sold to you are furnished "as is" by Howard Johnson's Enterprises, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Howard Johnson's Enterprises, Inc., the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc. the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc., or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.

To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc., the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

Spreader Settings

| Broadcast Spreader | Drop Spreader |
|---------------------------|-------------------------|
| Agri-Fab 4.0 | Agri-Fab 6.0 |
| Ames 4.0 | Ames 2.5 |
| Ames Deluxe 2in1 3.0 | Ames Deluxe 2in1 1.5 |
| Craftsman 4.0 | Craftsman 5.5 |
| Earthway 11.5 | Earthway 9.5 |
| Plant Mate 8.0 | Plant Mate 1.7 |
| Precision GreenLine 3.75 | Precision GreenLine 1.5 |
| Precision Red Devil 3.75 | Precision Red Devil 1.5 |
| Precision Pro 3.75 | Precision Pro 1.5 |
| Scotts Speedy Green 5.75 | Scotts Accu Green 6.5 |
| Scotts Easy Green 26.0 | Scotts Even Green 7.5 |
| Scotts Basic 5 | Taylor 1.7 |
| Scotts Lawn Pro 5 | |
| Scotts Deluxe EdgeGuard 5 | |
| Taylor 8.0 | |

Fertilizer may stain driveways, walks, patios, pools, etc. Sweep off thoroughly immediately after each use.

Manufactured and Guaranteed By:



Howard Johnson's Enterprises, Inc.
9675 S 60th Street
Franklin, WI 53132

© Copyright 2017
Howard Johnson's Enterprises, Inc.

PKRHUY080
C70-02012010A 12/17
100206621

