33530

Hi-Yield Ant Killer Granules



For Outdoor Use Around Residential Sites Only

- Effective Home Barrier Treatment
- Kills Listed* Bugs Outdoors-Before They Come Inside
- Use on Lawns, Ornamental and Flower Gardens, and Listed Vegetables
- Treats 136 Fire Ant Mounds or up to 3,500 sq. ft.

*Please refer to the insect lists found in the Directions for Use

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements on back of bag.

Active Ingredient:



KEEP OUT OF REACH OF CHILDREN

Hi-Yield **Ant Killer Granules**

FIRST AID Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Pall a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatme In the event of a medical or chemical emergency contact ChemTel, Inc.

North America 1-800-255-3924 or Worldwide Intl. + 01-813-248-0585

If on skin or

If in eves

Note to Physician:
This product is a pyrethroid. If large amounts have been ingested, milk, cream, or other digestible fats and oils may increase absorption and so should be avoided.

PRECAUTIONARY STATEMENTS **Hazards to Humans and Domestic Animals**

CAUTION: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the

ENVIRONMENTAL HAZARDS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product directly to the lawn or garden area. Water treated area as directed on this label

General Instructions

Hi-Yield* Ant Killer Granules is for use on lawns, around home foundations (perimeter treatment), omamental and flower gardens, and vegetable gardens for the control of the insect pests listed in the table below.

Lawns
Apply Hi-Yeld® Ant Killer Granules when insect activity or damage is first observed (lawn thins, brown spots appear, or moths fly up from the grass when mowed).

Spread Hi-Yield® Ant Killer Granules uniformly over lawn using the suggested spreader settings. For best results,

Spread H-Yield® Ant Killer Granules uniformly over lawn using the suggested spreader settings. For best results, use a mechanical spreader.

Water- in product after application.

Do not use any other weed, disease, or insect control products for at least one week after application.

For best results and to prevent further damage, treat the entire lawn, not just the browned areas.

Repeat application if there are signs of renewed insect activity but not more than once every 7 days.

NOTE: Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring

Armyworms (including Lawn Armyworm, Fall Armyworm) Billbugs (including Bluegrass Billbug, Hunting Billbug) Black Turfgrass Ataenius (adults), Bristletali Chiggers Chinch Bugs Cockroaches Cockroaches Crickets (including House Cricket, Common Short-tailed Cricket, Arizona Cricket, Tropical Cricket, Arizona Cricket, Tropical Cricket, Arizona Cricket, Tropical Cricket, Mealybeute, Mealybug Ladybettle, Twice-stabbed Ladybeetle, Twice-s	PE	APPLICATION RATE	
	Armýworm, Fall Armýworm) Billbug (including Bluegrass Billbug, Hunting Billbug) Black Turfgrass Ataenius (adults), Bristletail Chiggers Chinch Bugs Cockroaches Crickets (including House Cricket, Common Short-tailed Cricket, Arizona Cricket, Tropical Cricket, Variable Field Cricket, Fall Field Cricket, Southeastern Field Cricket, Spring Field Cricket, Northern Wood Cricket, Western Striped Cricket, Eastern Striped Cricket, Eastern Striped Cricket, Eastern Striped Cricket, Eastern Striped Cricket, Sand Field	Fire Brat Fruit Fly Grasshoppers Greenbug Ladybeetles (including Convergent Ladybeetles, Even-spotted Ladybeetle, Two-spotted Ladybeetle, Thirteen-spotted Ladybeetle, Twice-stabbed Ladybeetle, Asian Ladybeetle, Mealybug Ladybeitly Leaf Hoppers Lucerne Moth Mealybugs Pillbugs Seed Maggots Silverfish Spiders	a 1,000 sq. ft. area of lawn using a spreader. For best results, thoroughly water the treated area

Cricket)
Cutworms (including Black Cutworm, Bronzed Cutworm, Granulate Cutworm, Variegated Cutworm)
Dichondra Flea Beetle
Earwigs
European Crane Fly

Ants (including Carpenter S

European Crane Fly

Ants (including Carpenter, Red
Harvester, Red Pavement, Odorous,
Pyramid, Black
Turf, White-footed, Crazy, Little
Black, Small Honey, Lawn, Cornfield,
Ghost, Thief, Acrobat, Big-Headed,
and Allegheny mound)
Annual Bluegrass Weevil (Hyperodes)
Chiggers

Crane Flies (larvae)
Fleas (larvae)
Hoas Pleas (larvae)
Mole Crickets (including Tawny Mole
Cricket, Southern Mole Cricket,
Short-Winged Mole Cricket,
Weevils (including Annual Bluegrass)

Ants (including Argentine, Pharaoh (sugar), Imported Fire Ants, and Foraging Fire Ants) Centipedes Millipedes

Ticks (including Brown Dog Tick) and ticks that may transmit Lyme Disease

Uniformly apply 1 to 2 lbs. per 500 sq. ft. or 2 to 4 lbs. per 1,000 sq. ft. of lawn. For best results, thoroughly water the treated area after application.

Uniformly apply 1 lb. per 250 sq ft. area of lawn or 4 lbs. per 1,000 sq. ft. For best results, thoroughly water the treated area after application.

For Fire Ant Mounds: Uniformly apply 1 tablespoon or 3 teaspoons of granules to the surface of each mound. For best results apply in cool weather, 65° to 80° F, in early morning or late evening hours. Avoid disturbing ants. Treat new mounds as they appear. Thoroughly

In New York State, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
 In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

PI-YIEId® Ant Killer Granules can be applied just before planting or after plants emerge.

Apply 0.5 to 1 b. evenly over each 500 sq. ft. area. Repeat if there are signs of renewed insect activity, but not more than every two weeks.

**Home Foundation / Barrier Treatment

Evenly apply a 2 ft. wide band of granules next to the house/home foundation. Use 0.5** to 2.0 lb. for each 500 sq. ft. (2 ft. x 250 ft.). If necessary, repeat applications every 2 weeks only if there are signs of renewed insect activity.

***O-5 lb/500 sq. ft. is equivalent to 1 lb./1,000 sq. ft.

Hi-Vield® Ant Killer Granules can be applied to the vegetables listed below.
Apply a single application before planting or after plants emerge and work into the top 4 to 6 inches of soil. Use 1.0 lb. of product for a 500 sn. ft. area

ib. or product for a 500 sq. it. area.						
CROP	DAYS TO WAIT AFTER APPLICATION TILL HARVEST	PESTS				
Tomatoes	1	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms				
Sweet Corn	1	Ants, armyworms, cutworms, crickets, earwigs, European com borer, fall armyworm, grasshoppers, southwestern com borer and weevils				
Green Peas Edible Pods (Sugar Snap Peas, Snow Peas), Green Beans (Wax Beans, Snap Beans), Dry Beans (Black Eyed Peas, Cow Pea)	3	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms				
Cucurbit Vegetables Chayote, Citron melon, Qucumber, Edible Gourd Muskmelon, (Cantaloupe, casaba, Crenshaw melon, Golden melon, Honeydew Melon, Pineapple melon, Santa Claus melon, Snake melon), Pumpkin, Summer squash (Crockneck, Scallop, Straithneck, Zucchini), Winter squash (Butternut, Catalapthneck, Tucchini), Acom and Spaghetti), Watermelon	3	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms				
Artichoke	5	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms				
Broccoli, Brussels Sprouts, Chinese Broccoli, Cabbage, Caulfflower, Cavalo Broccoli, Chinese Cabbage, Kohlrabi, Chinese Mustard, Eggplant, Head Lettuce, Peppers (Bell and Non-Bell)	7	Arts, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms				

Spreading Instructions

mendations. <mark>Spreader Settings:</mark>

Spreader	Spreader settings covers 5,000 sq. ft. Pounds of product per 1,000 sq. ft.			Application Directions
	1 lb.	2 lb.	4 lb.	
Scotts®/Drop Spreader EZ Drop, Republic (#530)	2	2 ½	4 1⁄4	Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.
Scotts® Easy Green® Rotary EZ Broadcast, Republic (#540)	2	2 ¾	4 ½	Application automatically starts and stops when you do. For complete coverage make each pass about 30 inches from the previous one, or some overlapping occurs.
ferti•lome®/Earthway Ev-N-Spred (Broadcast)	11	13	15	

STORAGE AND DISPOSAL

inate water, food or feed by storage or disposal

PESTICIDE STORAGE: Store in a cool, dry, secure place. Do not store near seeds, fertilizers, or foodstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Notice: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and use not in



230 FM 87 **BONHAM, TEXAS 75418** EPA Reg. No. 53883-194-7401 EPA Est. No. 7401-TX-01 Product Questions? 855-270-4776

