



Moletox **Mole Killer**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling, Apply this product only for the sites, pests and application methods specified on this label, Read this Label:
Read this entire label and follow all use directions and precautions.
IMPORTANT: Do not expose children, pels or other non-larget animals to rodenticides. To help prevent acidents:

1. Store product not in use in a location out of reach of children and pels.

2. Follow all application directions and Use Restrictions on this labet. Apply beit only in moles' underground runways. Keep pets out of treated areas and isolate treated areas using existing fending, if present.

3. Dispose of product container and unused or spoiled bait as specified on this labet.

USE RESTRICTIONS: This product may be used only

as specified on his label.

USE RESTRICTIONS: This product may be used only to control eastern moles (Scalopus aquaticus), starmose moles (Condylura cristata), histy-tailed moles (Parastalops therewin), coast moles (Scalopanus cararius), broad-footed moles (S. histmanus) and/or lownsend moles (S. histmanus) and/or lownsend moles (S. histmanus) and/or lownsend moles (S. hoveendi) or lawns, tut fraes, gof courses and other non-food grassy areas. This product may only be applied at the use states and by the application methods indicated on this shale. Bail must be applied directly into moles' burtow ayatems (man underground furnell or subsurface runways). Do not place bad on the ground surface.

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of surface nidges in the lurf or by a series of concial mounds of early bushed up from deep burrows. Conical mounds may be largy the location of main underground runways.

BATINGS: Prior to realment, determine which burrow systems are active. Using a round-ended wood rod, such as a broomstick, probe ground in violently of conical mounds or surface ndiges until main underground runways is detected by a sudden decrease in resistance against the probe. Mark open burrow systems and revisit them 2-3 days latel. For deeper tunnels, open system with a shovel and leave open. Burrow systems and revisit them 2-3 days latel. For deeper tunnels, open system with a shovel and leave open. Burrow systems that are active are little by to be flugged by moles. Treat only active burrow systems. To treat a burrow system, locate main runways by probing. Remove the cap from the lip of the applicator syringe and turn dosing ring to mark indicating half ounce. Inject half an ounce of ball (one-sixth of the contents of one syringe) through probe or shoveled opened hole at each ball placement location. Make six (6) such placements per burrow system. Cover holes made to enter tunnel with so and and; but do not allow dirt to sift in to foul or cover bait. FOLLOW-LP. Check effects of treatment 4-5 days after ball application by opening and marking burrows in the manner used to check for initial activity. Return 2-3 days after ball syringe/10,000 ft.

Slorage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Do not store used or partially used syringes. Pesticide Disposal: Wastes resulting from the use of this product may be disposal: Wastes resulting from the use of this product may be disposal. Wastes resulting from the use of this product may be disposed facility. Container Handling: Non-refitlable container; do not reuse or refill this container. Offer for recycling, if available; otherwise, dispose of empty container in a sanitary landfill; or by incineration.

PRECAUTIONARY STATEMENTS

Mazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed. Keep away from
children, domestic animals and pets, Anyone handling
balts or retireving carcasses must wear gloves. Wash
thoroughly with soap and water after handling and
before eating, drinking, chewing gum, or using tobacco.

HAVE LABEL WHEN DISTAINING TREATMENT ADVICE
II Swallowed: * Call a person control center, doctor, or National Pesticide Information Center (1-900-858-7379) immediately for treatment advice.

Have a person sip a glass of water if able to swallow,
 Do not induce vomiting unless told to do so

TREATMENT FOR PET POISONING
If arimal each suit, call eviderivation or 1-900-922-1222 at once.
NOTE TO PHYSICIAN OR VETERINATION.
Contains the anticoagulant Warfarin. If swallowed, this
material may reduce the clotting ability of the blood and
cause bleeding. For humans or arimals ingesting ball and/or
howing poisoning signs (bleeding or elevated prothrombin
times), give Vitamin K, orally or intramuscularly.

Environmental Hazards
This product is extremely toxic to mammals and birds.
Doss, cats and other predatory and scavenging mammals
and birds might be poisoned if they feed upon animals
that have eaten this bait. Do not apply directly to water.

that have eaten this bait. Do not apply directly to water.

WARRANTY: To the extent consistent with applicable law,
Manufacturer and Seller make no warranty, expressed or implied,
concerning the use of this product other than indicated on the
label. Buyer assumes afrist of use and/or handling of this material
when such use and/or handling is contrary to label instructions.
Notice: This product contains a chemical know to the state of
California to cause birth defects or other reproductive harm.

Distributed by BONIDE PRODUCTS, INC.

5301 SUTLIFF ROAD - ORISKANY, NY 13424



REVENGE

Moletox **Mole Killer**

- Treats an entire burrow system
- Poison Bait with syringe applicator
- Use on lawns, turf areas, golf courses & other non-food grassy areas

Ge tc natu.



Keep Out CAUTI **Net Weig**

200000000

FIRST AID: HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If Swallowed: Call a poison control center, doctor, or National Pesticide Information Center (1-800-858-7378) immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian or 1-800-222-1222 at once.

NOTE TO PHYSICIAN OR VETERINARIAN: Contains the anticoagulant Warfarin. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K, orally or intramuscularly.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Do not store used or partially used syringes. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Handling: Non-refillable container; do not reuse or refill this container. Offer for recycling, if available; otherwise, dispose of empty container in a sanitary landfill: or by incineration.



Moletox® Mole Killer

This product may only be used to control moles in manual, below-ground applications.

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

Read complete label and instructions on packaging before using this product.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. Anyone handling baits or retrieving carcasses must wear gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

XXXXXXX

693