

GREENVIEW® Fairway Formula

Spring Fertilizer

Weed & Feed and Crabgrass Preventer

**Kills lawn weeds -
crabgrass, dandelions, viney
weeds, even clover**

**Extended time release feeding
of nitrogen lasts into the summer**

ACTIVE INGREDIENTS:

2,4-D	0.64%
Mecoprop-p	0.14%
Dicamba	0.06%
Dithiopyr	0.16%
OTHER INGREDIENTS:	99.00%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

12.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
2.8 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid
equivalent per ton.
1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
3.2 lbs. dithiopyr per ton.

Isomer Specific by AOAC Methods

EPA Reg. No. 2217-859-961

EPA Est. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

Covers 10,000 sq. ft.

24-0-6

Net Weight 36 lbs. (16.32 kg.)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for complete Precautionary Statements, including First Aid,
and Directions for Use, including Storage and Disposal.

NOTICE: Read the entire label. Use only according to label directions.

In New York State: Do not exceed one application at the recommended rate of
3.6 lbs. of product per 1,000 sq. ft. (0.25 lb active ingredient of dithiopyr per acre).

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. Avoid contact with skin, eyes or clothing. Wear long sleeved shirt and long pants, shoes plus socks, and waterproof gloves. Wash thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- If incorporation is accomplished by watering-in, do not enter or allow entry to the treated areas (except those involved in watering) until the watering-in is complete and the surface is dry.
- Read all precautions and directions for use carefully before applying this product.
- Do not use this product where desirable clovers are present or spread (apply) the granules onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosage for any area; be particularly careful within dripline of trees and other ornamental species.
- Do not use clippings from treated areas for mulching, vegetables, or fruit trees.
- Do not contaminate domestic or irrigation waters.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRODUCT INFORMATION

GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer will control your lawn's weeds in one application. GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer will kill lawn weeds as listed, which are commonly found in home lawns including dandelion, chickweed, thistle and clover, and will prevent crabgrass and other undesirable annual grasses. GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer also contains a long-lasting, timed release fertilizer that has all the nutrients your lawn needs for 2 to 3 months, depending on grass variety.

Use of GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer is recommended for the following established turf species:

Established Cool Season Turfgrasses - Kentucky bluegrass, perennial ryegrass, fine and tall fescues.

Established Warm Season Turfgrasses - Bahiagrass, bermudagrass, buffalograss, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floratum').

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer on colonial bentgrass, dichondra, carpetgrass, or 'Floratum' St. Augustinegrass.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

Weeds Controlled

When applied as directed, GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer will control the following weeds:

Annual Grass Weeds: Annual Bluegrass (Poa annua), Annual Ryegrass, Barnyardgrass, Crabgrass, Crowfootgrass, Foxtail (green and yellow), Goosegrass, Kikuyugrass (seedling), Perennial Ryegrass, Smutgrass (seedling).

Established Broadleaf Weeds: Annual yellow sweetclover; Aster; Austrian fieldcress; Bedstraw; Beggarticks; Betony, Florida; Bindweed, field; Bird vetch; Bitter wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black medic; Black mustard; Blackseed plantain; Blessed thistle; Bloodflower milkweed; Blue lettuce; Blue vervain; Bracted plantain; Brassbuttons; Bristly cottongue; Broadleaf dock; Broadleaf plantain; Broomweed; Buckhorn plantain; Bulbous buttercup; Bull thistle; Bullnettle; Burclover; Burdock; Burning nettle; Burweed; Buttercup; Buttonweed; Canada thistle; Carolina geranium; Carpetweed; Catchweed bedstraw; Catnip; Catsear; Chickweed, common; Chickweed, mouseear; Chicory; Cinquefoil; Clover, crimson; Clover, hop; Clover, red; Clover, strawberry; Clover, sweet; Clover, white; Cockle; Cocklebur; Common mullein; Creeping jenny; Cudweed; Curly dock; Daisy, English; Daisy fleabane; Daisy, oxeeye; Dandelion; Dichondra; Dogfennel; Dollarweed; False dandelion; Falseflax; False sunflower; Fiddleneck; Florida pusley; Frenchweed; Galinsoga; Goathead; Goldenrod; Ground ivy; Gumweed; Hairy fleabane; Hawkweed; Healall; Heartleaf drymary; Heath aster; Hedge bindweed; Hedge mustard; Henbit; Hoary cress; Hoary plantain; Hoary vervain; Horsenettle; Jimsonweed; Knawel; Knotweed; Kochia; Lambsquarters; Lespedeza; Mallow; Matchweed; Mexicanweed; Milk vetch; Morningglory; Mouseear hawkweed; Mugwort; Musk thistle; Mustard; Narrowleaf plantain; Narrowleaf vetch; Nettle; Orange hawkweed; Oriental cocklebur; Oxalis; Parsley-piert; Parsnip; Pearlwort; Pennycress; Pepperweed; Pigweed; Pineappleweed; Pineywoods bedstraw; Plains tickseed; Plantain; Poison ivy; Poison oak; Pokeweed; Poorjoe; Prairie sunflower; Prickly lettuce; Prickly sida; Prostrate knotweed; Prostrate pigweed; Prostrate spurge; Prostrate vervain; Puncturevine; Purslane, common; Ragweed; Red sorrel; Redroot pigweed; Redstem filaree; Rough cinquefoil; Rough fleabane; Russian pigweed; Scarlet pimpernel; Scotch thistle; Sheep sorrel; Shepherdspurse; Slender plantain; Smallflower galinsoga; Smooth dock; Smooth pigweed; Sorrel; Sowthistle; Spanisneedles; Speedwell; Spiny amaranth; Spiny cocklebur; Spiny sowthistle; Spotted catsear; Spotted spurge; Spurweed; Stinging nettle; Strawberry, india mock; Tall nettle; Tall vervain; Tansy ragwort; Tansy mustard; Tanweed; Thistle; Trailing crownvetch; Tumble mustard; Tumble pigweed; Velvetleaf; Venice mallow; Virginia buttonweed; Virginia creeper; Virginia pepperweed; Wavyleaf bullthistle; Western clematis; Western salsify; White mustard; Wild aster; Wild buckwheat; Wild carrot; Wild four-o'clock; Wild garlic; Wild geranium; Wild lettuce; Wild marigold; Wild mustard; Wild onion; Wild parsnip; Wild radish; Wild rape; Wild sweet potato; Wild vetch; Woodsorrel; Woolly croton; Woolly morningglory; Woolly plantain; Wormseed; Yarrow; Yellow rocket; Yellowflower pepperweed.


When To Apply

GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer can be used either in the spring or fall. This product should be used when the grass and weeds are actively growing and the soil is moist.

Do not apply under drought conditions.

In the spring, the ideal time to apply GreenView Fairway Formula Spring Fertilizer Weed & Feed and Crabgrass Preventer is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application of this product in 6-8 weeks for best weed and crabgrass control and for full season fertilization.

In the fall, apply this product when the weather cools and both the grass and weeds are actively growing, generally around Labor Day.



Drop Spreader Settings

Apply in a single pass at a rate of approximately 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working toward the center. Overlap the coverage slightly, and shut off the spreader before making hard turns.

Drop Spreader	Setting
Earthway®	14 1/2
Republic® Drop	5
Scotts® AccuGreen	5
Sta-Green®	4



If your spreader model is not listed, go to our website at www.lebsea.com, or call toll-free 800-233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

How To Apply

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Do not apply this product onto desirable plants or shrubs as damage may result. If the product gets onto other plants immediately brush or wash it off the foliage.

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. nitrogen per 1,000 sq. ft.

How Much To Use

Apply 3.6 pounds of product per 1,000 sq. ft. of turfgrass.

In New York State: Do not exceed one application at the recommended rate of 3.6 lb of product per 1,000 sq. ft. (0.25 lb active ingredient of diithiopyr per acre).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq. ft.


Calibration Instructions

- For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- Check equipment frequently to verify calibration.

Broadcast Spreader Settings

Apply in a single pass at a rate of approximately 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. A little overlap coverage is desirable.

Broadcast Spreader	Setting
Agri-Fab® Rotary	4
Earthway® Rotary	15
Republic® Rotary	10
Scotts® Deluxe Edgeguard	4 1/2
Scotts® EasyGreen	25
Sta-Green®	3 1/2



To calibrate your spreader: 1 lb. of material should cover approximately 275 sq. ft.

GUARANTEED ANALYSIS

Total Nitrogen (N)	24%
2.8% Ammoniacal Nitrogen	
6.4% Water Insoluble Nitrogen*	
5.4% Urea Nitrogen	
9.4% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	6%
Sulfur (S)	5.6%
5.6% Combined Sulfur (S)	
Derived from: Ammonium Sulfate, Urea, Methylene Ureas, Muriate of Potash.	
*15.8% Slowly Available Nitrogen from Methylene Ureas.	
F699	

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.Regulatory-info-lebsea.com>

GUARANTEED SATISFACTION

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

Limited Warranty and Disclaimer

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Distributed by:

Lebanon Seaboard Corporation
1600 East Cumberland Street, Lebanon, PA 17042

If your spreader model is not listed, go to our website at www.lebsea.com, or call toll-free 800-233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.



21-29173