

LESCO DIMENSION* 0.21% Plus Fertilizer

NOT FOR USE on turf being grown for sale or other commercial use as seed, or for commercial sale production, or for research purposes. In New York State this product may only be used by commercial applicators and at no more than 500 lb (0.5 lb of active ingredient) per acre per year (or 11.5 lb product per 1,000 ft² per year) and its prohibition from use in New and Suburb Counties.

Contains LESCO® Poly Plant Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.	
ACTIVE INGREDIENT	
Diuretic, 5,5-Dimethyl-2-thiazolidinone N-ethyl-2-thiohydantoin	0.21%
2-thiohydantoin	1.23%
INERT INGREDIENTS	99.76%

Product protected by U.S. Patent No. 4,932,184. Other patents pending. Read the entire label before using this product. Use only according to label instructions. **NOTICE:** Before using this product, read the Use Precautions, Warranty Statement, Directions for Use, and the Storage and Disposal Instructions. If the Warranty Statement is not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
Inhalation	Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the poison control center or doctor. • Do not get product or liquid in eyes.
Eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
On skin or clothing	• Take off contaminated clothing. • Wash immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
On face	• Wipe product from face. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Wash thoroughly with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor for treatment. You may also contact Chem-Treat at 1-800-241-3330 for emergency medical treatment information.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Causes eye irritation. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wear protective eyewear such as goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS
This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to riparian areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not discharge water when disposing of equipment washwaters.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION
This product is a herbicide that provides control of weeds, other annual grasses, and broadleaf weeds in established lawns and ornamental lawns, including golf courses, tennis courts, and lawns. This product will not harm newly established ornamentals when used according to label directions. This product may be used on seeded, sodded, or sprigged lawns and ornamental lawns that are well-established. The grass roots have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental lawns that are not well-established, or on lawns that have been weakened by weather, pests, diseases, chemical, or mechanical-related stresses, may increase the chances for turf injury. This product should only be applied to lawns and ornamental lawns that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established turfgrasses are tolerant to this product:

Common Name	
Aztecgrass	Agrostis pallens
Beardgrass	Poa pratensis
Bluegrass	Poa annua
Creolegrass	Pennisetum clandestinum
St. Augustine	Stenotaphrum secundatum
Turfgrass	Cynodon dactylon
Yarrowgrass	Poa annua

Common Name	Scientific Name
Aztecgrass	Agrostis pallens
Beardgrass	Poa pratensis
Bluegrass	Poa annua
Creolegrass	Pennisetum clandestinum
St. Augustine	Stenotaphrum secundatum
Turfgrass	Cynodon dactylon
Yarrowgrass	Poa annua

DO NOT apply this product to Colonial Turfgrasses (Agrostis tenuis) varieties. Use of this product on certain varieties of Colonial Turfgrasses, such as "Colonial", "Carmen", "Seaside", and "Washington" may result in undesirable turfgrass injury. Not all varieties of Colonial Turfgrasses have been tested. Use of this product on certain varieties of Fine Fescue may result in undesirable turf injury. The following Fine Fescue varieties have been tested to be sensitive to this product: "Maverik", "Banner", "Majesty", "Majesty", "F.C.", "Turf", "YF-67", "Nightjar", "Imbi", "Jamaican", "Nobel", "Napoli", "Mary", "Pentagon", "Tanger", "Vigoro", "Vigoro", and "Wild Owl". Not all varieties of Fine Fescue have been tested.



Re seeding, Overseeding, or Sprigging
Re seeding, overseeding or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Re seeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When re seeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed.

USE PRECAUTIONS
• Use on lawns on Golf Courses Putting Greens.
• The early post-emergence crabgrass control described below is best. Post-emergence applications are only effective on crabgrass if applied before crabgrass has developed its 10th leaf or six tillers. For best results, do not apply to areas where crabgrass is already established.
• Do not apply to areas where crabgrass is already established.
• Do not use dipping from treated turf for mulching around vegetables or fruit trees.
• Do not apply this product to wet lawns, roofs or other wet surfaces.
• Do not apply this product to wet lawns, roofs and parts of treated areas until dust has settled.

18-0-10

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	18.00%
18.00% Urea Nitrogen	
0.00% Water Insoluble Nitrogen	
0.00% Other Water Soluble Nitrogen	
SOLUBLE POTASH (K ₂ O)	10.00%
DERIVED FROM: Polymer Coated Urea, Urea, Nitrate of Potash.	
CHLORINE (Cl) Max.	7.50%
18.00% Slowly Available Urea Nitrogen from Polymer Coated Urea.	

APPLICATION DIRECTIONS - Control of Crabgrass

Pre-emergence and Early Post-emergence Control
This product provides "pre-emergence" control of crabgrass including the large, smooth, and southern species when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental lawns. It can also provide "early post-emergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in established lawns and ornamental lawns. Post-emergence crabgrass control will be obtained only when this product is applied prior to the flowering of crabgrass, which generally corresponds to the time when you can first see it in the crabgrass plants in the lawn or turf. See the product benefits of this product's additional early post-emergence activity in the label compared to other pre-emergence application products. Dimension controls crabgrass prior to and up to 4 weeks after germination.

Application Frequency and Timing
This product may be applied as a single application, as a split application, or as a repeated application for crabgrass control at the spring, summer, or fall.
DO NOT apply more than 3.46 lb of this product per 1,000 ft² per application, and no more than 18.28 lb of this product per 1,000 ft² per year.

Spring Applications
For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user location, the turfgrass mowing height, and whether the use is intended to be pre-emergence or early post-emergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Mid-Treatment (M) for pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 10 months of pre-emergence control. This program may also be used for early post-emergence control at sites where the turfgrass is cut low (e.g., golf courses), and (B) low grass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence control. This program may also be used for early post-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns).

Mid-Treatment (M) for pre-emergence control at sites where the turfgrass is cut low (e.g., golf courses) and (B) maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence control. This program may also be used for early post-emergence control at sites where the turfgrass is cut low (e.g., golf courses).

Subsequent pre- and/or post-emergence applications should be made when longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1. Recommended Single Application Use Rates*

PROGRAM	Pre-emergence Control	Mid-Treatment (M)	Early Post-emergence Control
Rate (lb/1,000 ft ²)	3.46	3.46	3.46
Rate (lb/100 ft ²)	0.346	0.346	0.346
Rate (lb/1,000 sq yd)	3.46	3.46	3.46
Rate (lb/100 sq yd)	0.346	0.346	0.346

Note: 2,7,27 lb per 1,000 ft² is the equivalent of 0.28, 2.8, 28.0 lb active ingredient per acre.

* If applied in the fall, this product may not provide adequate control or application, and may require more than 18.28 lb of this product per 1,000 ft² per year. Post-emergence control is limited. Use "Crabgrass Control" and "Preseason" sections above. Reports indicate areas have been lost. **Do not use on the following turfgrasses:**
- Do not apply to areas where crabgrass is already established.
- Do not apply to areas where crabgrass is already established.

Fall Applications
This product can also be applied in the late summer or early fall (late August through November) at the "Preseason" rate used above in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

The Pre-season Control
This product may be applied in the late summer or early fall (late August through November) at the "Preseason" rate used above in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Spring Application
Using an directed for crabgrass control in the spring, this product will also control (at the "Preseason" rate) the following weeds when applied prior to their emergence:
- Annual weeds (annual)
- Perennial weeds (perennial)
- Broadleaf weeds (broadleaf)
- Yellow foxtail (perennial)
- Yellow foxtail (annual)
- Yellow foxtail (perennial)
- Yellow foxtail (annual)

Not controlled at the "Preseason" rate:
- Annual weeds (annual)
- Perennial weeds (perennial)
- Broadleaf weeds (broadleaf)
- Yellow foxtail (perennial)
- Yellow foxtail (annual)
- Yellow foxtail (perennial)
- Yellow foxtail (annual)

SUGGESTED SPREADER SETTINGS*

	Recommended Rate in lb/1,000 ft ²
1.5 lb	1.5 lb
2.7 lb	2.7 lb
4.5 lb	4.5 lb
11" 17"	11" 17"
LESCO Calibration Gauge	
G00709818A	1 1/2 2 1/2 3 1/2 4 1/2
Crabgrass or Sprigged	2 1/2 3 3 1/2 4 1/2
LESCO Product	1 1/2 1 1/2 2 2 1/2 3 1/2

*IMPORTANT: These settings are only approximate. Always consult your spreader and operator manual on these settings. Be sure to always read your spreader and operator manual for full instructions. An optional (Optional) setting is shown on the back of this label.

APPLICATION EQUIPMENT AND INSTRUCTIONS
Apply this product with the appropriate application equipment for your site. Read the label and operator manual for full instructions. This product is not to be applied to areas where it is not intended to be applied. Follow the application instructions for each application. Do not apply to areas where it is not intended to be applied. Follow the application instructions for each application. Do not apply to areas where it is not intended to be applied. Follow the application instructions for each application.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secure location.
PESTICIDE DISPOSAL: Whatever remains from the use of this product that cannot be used as intended, or is otherwise unsuitable for use, should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local provisions. Do not use or reuse this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of properly (e.g., in a sanitary landfill or by incineration), if allowed by state and local authorities. By burning, if burning is the only method available for disposal.
SPEL: In case of spill, soak up material and dispose of material according to "Produce Disposal" instructions above.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY
NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before using this product. If the terms are not acceptable, return the product to the retailer. The Directions for Use of this product must be followed carefully and completely. It is the responsibility of the user to determine whether the use of this product is appropriate for the site and conditions. The user assumes all liability for any damage or injury resulting from the use of this product, whether or not caused in whole or in part by the use of this product. The user is responsible for the selection of the site and conditions of use. The user is responsible for the selection of the site and conditions of use. The user is responsible for the selection of the site and conditions of use.

LESKO and Plus are registered trademarks and the remaining design is a trademark of LESKO Technology, LLC. IN LESKO is a registered trademark of Sun Applications. G00711 is a registered trademark of the LESKO Company. All other trademarks are the property of their respective owners. L14518-18-0-10 (R) 11/15
Net Weight: 50 lb (22.7 kg) SG# 248
EPA Reg. No. 145-1044-10 (R) 11/15
EPA Reg. No. 145-1044-10 (R) 11/15
EPA Reg. No. 145-1044-10 (R) 11/15
EPA Reg. No. 145-1044-10 (R) 11/15
EPA Reg. No. 145-1044-10 (R) 11/15
EPA Reg. No. 145-1044-10 (R) 11/15

#702032 PP